


## (2.1) The Bisection Method

MATH 4701 Numerical Analysis

# Intermediate Value Theorem


**Intermediate value theorem (special case):** If  $f(x) : [a, b] \rightarrow \mathbb{R}$  is **continuous** and  $f(a)f(b) < 0$  ( $f(a)$  and  $f(b)$  have opposite signs) then for some  $a < \alpha < b$  we have  $f(\alpha) = 0$ .



# Intermediate Value Theorem

**Intermediate value theorem (special case):** If  $f(x) : [a, b] \rightarrow \mathbb{R}$  is **continuous** and  $f(a)f(b) < 0$  ( $f(a)$  and  $f(b)$  have opposite signs) then for some  $a < \alpha < b$  we have  $f(\alpha) = 0$ .

Notice that the theorem states the existence of one solution to  $f(x) = 0$  while there may be multiple such solutions.



# Intermediate Value Theorem

**Intermediate value theorem (special case):** If  $f(x) : [a, b] \rightarrow \mathbb{R}$  is **continuous** and  $f(a)f(b) < 0$  ( $f(a)$  and  $f(b)$  have opposite signs) then for some  $a < \alpha < b$  we have  $f(\alpha) = 0$ .

Notice that the theorem states the existence of one solution to  $f(x) = 0$  while there may be multiple such solutions.

While the theorem seems to only have theoretical implications on existence of a solution without giving any information on how that solution can be found, one can in fact approximate a solution  $f(x) = 0$  using **bisection method** which is based on the intermediate value theorem.

# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.

# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.
- If  $f(x_{n+1})f(B) < 0$  then we let  $A = x_{n+1}$  and we keep the value of  $B$  unchanged.



# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.
- If  $f(x_{n+1})f(B) < 0$  then we let  $A = x_{n+1}$  and we keep the value of  $B$  unchanged.
- If  $f(x_{n+1}) = 0$  we announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  and we terminate the process.

# Bisection Method

Suppose  $f(x)$  is a **continuous** function which satisfies  $f(x_0)f(x_1) < 0$  and we ask for a value  $\epsilon > 0$  of maximum **error** we are willing to tolerate.

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.
- If  $f(x_{n+1})f(B) < 0$  then we let  $A = x_{n+1}$  and we keep the value of  $B$  unchanged.
- If  $f(x_{n+1}) = 0$  we announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  and we terminate the process.

Notice that in every step  $f(A)$  and  $f(B)$  will remain two non zero numbers with opposite signs and the above three options are the only possibilities.

# Bisection Method

We start by letting  $A = x_0$  and  $B = x_1$ .

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.
- If  $f(x_{n+1})f(B) < 0$  then we let  $A = x_{n+1}$  and we keep the value of  $B$  unchanged.
- If  $f(x_{n+1}) = 0$  we announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  and we terminate the process.

Notice that in every step  $f(A)$  and  $f(B)$  will remain two non zero numbers with opposite signs and the above three options are the only possibilities.

**Termination:** We stop the algorithm and announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  once  $\frac{|x_1 - x_0|}{2^n} < \epsilon$ .

# Bisection Method

At each step of the algorithm, we let  $x_{n+1} = \frac{A+B}{2}$  (midpoint of  $A$  and  $B$ ).

- If  $f(x_{n+1})f(A) < 0$  then we let  $B = x_{n+1}$  and we keep the value of  $A$  unchanged.
- If  $f(x_{n+1})f(B) < 0$  then we let  $A = x_{n+1}$  and we keep the value of  $B$  unchanged.
- If  $f(x_{n+1}) = 0$  we announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  and we terminate the process.

Notice that in every step  $f(A)$  and  $f(B)$  will remain two non zero numbers with opposite signs and the above three options are the only possibilities.

**Termination:** We stop the algorithm and announce  $\alpha = x_{n+1}$  as the solution of  $f(x) = 0$  once  $\frac{|x_1 - x_0|}{2^n} < \epsilon$ . It is because  $x_{n+1}$  is the midpoint of an interval of length  $\frac{|x_1 - x_0|}{2^{n-1}}$  where a solution of  $f(x) = 0$  exists.

# Examples

- Use the bisection method for  $f(x) = \sin x + x^2 - 2$  with  $x_0 = -2$  and  $x_1 = -1$  to approximate a solution of  $f(x) = 0$  accurate within an error of at most  $\frac{1}{50}$ .

# Examples

- Use the bisection method for  $f(x) = \sin x + x^2 - 2$  with  $x_0 = -2$  and  $x_1 = -1$  to approximate a solution of  $f(x) = 0$  accurate within an error of at most  $\frac{1}{50}$ .
- If we use the bisection method for  $f(x) = x^3 - 2x + 0.5$  with  $x_0 = -2$  and  $x_1 = 2$  what is the smallest index  $n$  for which we know  $x_n$  is within 0.001 of an actual solution of  $f(x) = 0$ .

# Examples

- Use the bisection method for  $f(x) = \sin x + x^2 - 2$  with  $x_0 = -2$  and  $x_1 = -1$  to approximate a solution of  $f(x) = 0$  accurate within an error of at most  $\frac{1}{50}$ .
- If we use the bisection method for  $f(x) = x^3 - 2x + 0.5$  with  $x_0 = -2$  and  $x_1 = 2$  what is the smallest index  $n$  for which we know  $x_n$  is within 0.001 of an actual solution of  $f(x) = 0$ .
- For  $F(x) = (x + 1)x^3(x - 1)(x - 2)^2(x - 3)(x - 5)$  if we start with  $x_0 = -2$  and  $x_1 = 7$  which of the solutions of  $F(x) = 0$  will the bisection method find.