

(2.5) Accelerating Convergence

MATH 4701 Numerical Analysis

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**. This means there is a positive number $\lambda < 1$ such that

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lambda$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**. This means there is a positive number $\lambda < 1$ such that

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lambda$$

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\frac{p_{n+1} - p}{p_n - p} \approx \frac{p_{n+2} - p}{p_{n+1} - p} \approx \lambda$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**. This means there is a positive number $\lambda < 1$ such that

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lambda$$

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\frac{p_{n+1} - p}{p_n - p} \approx \frac{p_{n+2} - p}{p_{n+1} - p} \approx \lambda$$

$$\Rightarrow (p_{n+1} - p)^2 \approx (p_n - p)(p_{n+2} - p)$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**. This means there is a positive number $\lambda < 1$ such that

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lambda$$

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\frac{p_{n+1} - p}{p_n - p} \approx \frac{p_{n+2} - p}{p_{n+1} - p} \approx \lambda$$

$$\Rightarrow (p_{n+1} - p)^2 \approx (p_n - p)(p_{n+2} - p)$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} + p^2 \approx p_n p_{n+2} - p(p_n + p_{n+2}) + p^2$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\frac{p_{n+1} - p}{p_n - p} \approx \frac{p_{n+2} - p}{p_{n+1} - p} \approx \lambda$$

$$\Rightarrow (p_{n+1} - p)^2 \approx (p_n - p)(p_{n+2} - p)$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} + p^2 \approx p_n p_{n+2} - p(p_n + p_{n+2}) + p^2$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} \approx p_n p_{n+2} - p(p_n + p_{n+2})$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\Rightarrow (p_{n+1} - p)^2 \approx (p_n - p)(p_{n+2} - p)$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} + p^2 \approx p_n p_{n+2} - p(p_n + p_{n+2}) + p^2$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} \approx p_n p_{n+2} - p(p_n + p_{n+2})$$

$$\Rightarrow p(p_n + p_{n+2}) - 2pp_{n+1} \approx p_n p_{n+2} - p_{n+1}^2$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} + p^2 \approx p_n p_{n+2} - p(p_n + p_{n+2}) + p^2$$

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} \approx p_n p_{n+2} - p(p_n + p_{n+2})$$

$$\Rightarrow p(p_n + p_{n+2}) - 2pp_{n+1} \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p(p_n + p_{n+2} - 2p_{n+1}) \approx p_n p_{n+2} - p_{n+1}^2$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\Rightarrow p_{n+1}^2 - 2pp_{n+1} \approx p_n p_{n+2} - p(p_n + p_{n+2})$$

$$\Rightarrow p(p_n + p_{n+2}) - 2pp_{n+1} \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p(p_n + p_{n+2} - 2p_{n+1}) \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p \approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}}$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\Rightarrow p(p_n + p_{n+2}) - 2pp_{n+1} \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p(p_n + p_{n+2} - 2p_{n+1}) \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p \approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}}$$

$$= \frac{p_n p_{n+2} - 2p_n p_{n+1} + p_n^2 - p_{n+1}^2 + 2p_n p_{n+1} - p_n^2}{p_n + p_{n+2} - 2p_{n+1}}$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\Rightarrow p(p_n + p_{n+2} - 2p_{n+1}) \approx p_n p_{n+2} - p_{n+1}^2$$

$$\Rightarrow p \approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}}$$

$$= \frac{p_n p_{n+2} - 2p_n p_{n+1} + p_n^2 - p_{n+1}^2 + 2p_n p_{n+1} - p_n^2}{p_n + p_{n+2} - 2p_{n+1}}$$

$$= \frac{p_n(p_{n+2} - 2p_{n+1} + p_n) - (p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\begin{aligned}\Rightarrow p &\approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}} \\&= \frac{p_n p_{n+2} - 2p_n p_{n+1} + p_n^2 - p_{n+1}^2 + 2p_n p_{n+1} - p_n^2}{p_n + p_{n+2} - 2p_{n+1}} \\&= \frac{p_n(p_{n+2} - 2p_{n+1} + p_n) - (p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} \\&= p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}\end{aligned}$$

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\begin{aligned}\Rightarrow p &\approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}} \\ &= \frac{p_n(p_{n+2} - 2p_{n+1} + p_n) - (p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} \\ &= p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}\end{aligned}$$

Aitken's Δ^2 method assumes that the sequence (\widehat{p}_n) defined by

$\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}$ converges to p faster than the original sequence (p_n) .

Aitken's Δ^2 Method

Suppose (p_n) is a sequence converging to p **linearly**.

If we assume for large values of n , the signs of $p_n - p$, $p_{n+1} - p$, $p_{n+2} - p$ are all the same (or they alternate), we have

$$\begin{aligned}\Rightarrow p &\approx \frac{p_n p_{n+2} - p_{n+1}^2}{p_n + p_{n+2} - 2p_{n+1}} \\ &= p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}\end{aligned}$$

Aitken's Δ^2 method assumes that the sequence (\widehat{p}_n) defined by $\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}$ converges to p faster than the original sequence (p_n) .

Theorem: Suppose $\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lambda$ such that $0 < \lambda < 1$. Then $\lim_{n \rightarrow \infty} \frac{|\widehat{p}_n - p|}{|p_n - p|} = 0$.

Examples

If $p_n = \frac{1}{2^n}$, then

$$\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}}$$

Examples

If $p_n = \frac{1}{2^n}$, then

$$\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} = \frac{1}{2^n} - \frac{\left(\frac{1}{2^{n+1}} - \frac{1}{2^n}\right)^2}{\frac{1}{2^n} + \frac{1}{2^{n+2}} - \frac{2}{2^{n+1}}}$$

Examples

If $p_n = \frac{1}{2^n}$, then

$$\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} = \frac{1}{2^n} - \frac{\left(\frac{1}{2^{n+1}} - \frac{1}{2^n}\right)^2}{\frac{1}{2^n} + \frac{1}{2^{n+2}} - \frac{2}{2^{n+1}}} = \frac{1}{2^n} - \frac{\frac{1}{2^{2n+2}}}{\frac{1}{2^{n+2}}} = \frac{1}{2^n} - \frac{1}{2^{n+2}}$$

Examples

If $p_n = \frac{1}{2^n}$, then

$$\begin{aligned}\widehat{p}_n &= p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} = \frac{1}{2^n} - \frac{\left(\frac{1}{2^{n+1}} - \frac{1}{2^n}\right)^2}{\frac{1}{2^n} + \frac{1}{2^{n+2}} - \frac{2}{2^{n+1}}} = \frac{1}{2^n} - \frac{\frac{1}{2^{2n+2}}}{\frac{1}{2^{n+2}}} \\ &= \frac{1}{2^n} - \frac{1}{2^n} = 0\end{aligned}$$

Examples

If $p_n = \frac{1}{2^n}$, then

$$\begin{aligned}\widehat{p}_n &= p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} = \frac{1}{2^n} - \frac{\left(\frac{1}{2^{n+1}} - \frac{1}{2^n}\right)^2}{\frac{1}{2^n} + \frac{1}{2^{n+2}} - \frac{2}{2^{n+1}}} = \frac{1}{2^n} - \frac{\frac{1}{2^{2n+2}}}{\frac{1}{2^{n+2}}} \\ &= \frac{1}{2^n} - \frac{1}{2^n} = 0\end{aligned}$$

So for $p_n = \frac{1}{2^n}$ which converges to zero linearly, the sequence (\widehat{p}_n) is constant zero sequence.

Examples

Suppose $g(x) = \frac{x^2+2}{2x}$. Let $p_0 = 4$ and $p_{n+1} = g(p_n)$. Find the first five entries in the sequence $(\widehat{p_n})$.

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(\Delta a_n)$$

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(\Delta a_n) = \Delta(a_{n+1} - a_n)$$

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(\Delta a_n) = \Delta(a_{n+1} - a_n) = (a_{n+2} - a_{n+1}) - (a_{n+1} - a_n)$$

Δ Notation

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(a_{n+1} - a_n) = (a_{n+2} - a_{n+1}) - (a_{n+1} - a_n) = a_{n+2} - 2a_{n+1} + a_n$$

Δ Notation

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(a_{n+1} - a_n) = (a_{n+2} - a_{n+1}) - (a_{n+1} - a_n) = a_{n+2} - 2a_{n+1} + a_n$$

On the other hand,

$$(\Delta a_n)^2 = (a_{n+1} - a_n)^2$$

Δ Notation

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(a_{n+1} - a_n) = (a_{n+2} - a_{n+1}) - (a_{n+1} - a_n) = a_{n+2} - 2a_{n+1} + a_n$$

On the other hand,

$$(\Delta a_n)^2 = (a_{n+1} - a_n)^2 = a_{n+1}^2 - 2a_{n+1}a_n + a_n^2$$

Δ Notation

If (a_n) is a sequence of numbers, Δa_n is the sequence of **forward differences** of consecutive terms of (a_n) defined by

$$\Delta a_n = a_{n+1} - a_n$$

In particular,

$$\Delta^2 a_n = \Delta(a_{n+1} - a_n) = (a_{n+2} - a_{n+1}) - (a_{n+1} - a_n) = a_{n+2} - 2a_{n+1} + a_n$$

On the other hand,

$$(\Delta a_n)^2 = (a_{n+1} - a_n)^2 = a_{n+1}^2 - 2a_{n+1}a_n + a_n^2$$

Therefore, Aitken's Δ^2 accelerated sequence can be described by

$$\widehat{p}_n = p_n - \frac{(p_{n+1} - p_n)^2}{p_n + p_{n+2} - 2p_{n+1}} = p_n - \frac{(\Delta p_n)^2}{\Delta^2 p_n}$$

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Steffensen's Method

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Steffensen's Method

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

Steffensen's Method

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

Steffensen's Method

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

We then use $p_0^{(1)}$ as an initial guess to generate $p_1^{(1)} = g(p_0^{(1)})$ and $p_2^{(1)} = g(p_1^{(1)})$.

Steffensen's Method

Steffensen's method is a variation of Aitken's Δ^2 acceleration applied to iterative processes.

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

We then use $p_0^{(1)}$ as an initial guess to generate $p_1^{(1)} = g(p_0^{(1)})$ and $p_2^{(1)} = g(p_1^{(1)})$.

Using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0^{(1)}}$ and we call it $p_0^{(2)}$.

Steffensen's Method

Suppose $g(x)$ is a function and the sequence (p_n) defined by $p_{n+1} = g(p_n)$ converges linearly to a fixed point of g .

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

We then use $p_0^{(1)}$ as an initial guess to generate $p_1^{(1)} = g(p_0^{(1)})$ and $p_2^{(1)} = g(p_1^{(1)})$.

Using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0^{(1)}}$ and we call it $p_0^{(2)}$.

In every step, once $p_0^{(n)}$ is generated we use it as an initial guess to generate $p_1^{(n)} = g(p_0^{(n)})$ and $p_2^{(n)} = g(p_1^{(n)})$.

Steffensen's Method

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

We then use $p_0^{(1)}$ as an initial guess to generate $p_1^{(1)} = g(p_0^{(1)})$ and $p_2^{(1)} = g(p_1^{(1)})$.

Using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0^{(1)}}$ and we call it $p_0^{(2)}$.

In every step, once $p_0^{(n)}$ is generated we use it as an initial guess to generate $p_1^{(n)} = g(p_0^{(n)})$ and $p_2^{(n)} = g(p_1^{(n)})$.

Then using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0^{(n)}}$ and we call it $p_0^{(n+1)}$.

Steffensen's Method

Starting with an initial guess p_0 we find $p_1 = g(p_0)$ and $p_2 = g(p_1)$.

At this point, using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0}$ and we call it $p_0^{(1)}$.

In every step, once $p_0^{(n)}$ is generated we use it as an initial guess to generate $p_1^{(n)} = g(p_0^{(n)})$ and $p_2^{(n)} = g(p_1^{(n)})$.

Then using these three entries we find the Aitken's Δ^2 entry $\widehat{p_0^{(n)}}$ and we call it $p_0^{(n+1)}$.

Example: Suppose $g(x) = \frac{x^2+2}{2x}$. Let $p_0 = 4$ and use the Steffensen's method to approximate a fixed point of $g(x)$ accurate to an error of at most 10^{-6} .