

(3.4) Hermite Interpolation

MATH 4701 Numerical Analysis

Hermite Interpolation

Suppose x_0, x_1, \dots, x_n are $n + 1$ **distinct** numbers, and y_0, y_1, \dots, y_n and z_0, z_1, \dots, z_n are arbitrary numbers.

Hermite Interpolation

Suppose x_0, x_1, \dots, x_n are $n + 1$ **distinct** numbers, and y_0, y_1, \dots, y_n and z_0, z_1, \dots, z_n are arbitrary numbers.

We would like to find a polynomial $H(x)$ of degree at most $2n + 1$ such that

$$\textcircled{H} \quad H(x_k) = y_k \quad \text{and} \quad H'(x_k) = z_k \quad \text{for all } 0 \leq k \leq n.$$

Hermite Interpolation

Suppose x_0, x_1, \dots, x_n are $n + 1$ **distinct** numbers, and y_0, y_1, \dots, y_n and z_0, z_1, \dots, z_n are arbitrary numbers.

We would like to find a polynomial $H(x)$ of degree at most $2n + 1$ such that

$$\textcircled{H} \quad H(x_k) = y_k \quad \text{and} \quad H'(x_k) = z_k \quad \text{for all } 0 \leq k \leq n.$$

Although we already know how to find such polynomial using divided differences, we want to find a description of such polynomial in a way similar to that of Lagrange interpolating polynomial as it turns out to be useful later.

Hermite Interpolation

Suppose x_0, x_1, \dots, x_n are $n + 1$ **distinct** numbers, and y_0, y_1, \dots, y_n and z_0, z_1, \dots, z_n are arbitrary numbers.

We would like to find a polynomial $H(x)$ of degree at most $2n + 1$ such that

$$\textcircled{H} \quad H(x_k) = y_k \quad \text{and} \quad H'(x_k) = z_k \quad \text{for all } 0 \leq k \leq n.$$

For each $0 \leq k \leq n$, we find a polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Hermite Interpolation

We would like to find a polynomial $H(x)$ of degree at most $2n + 1$ such that

$$\textcircled{H} \quad H(x_k) = y_k \quad \text{and} \quad H'(x_k) = z_k \quad \text{for all } 0 \leq k \leq n.$$

For each $0 \leq k \leq n$, we find a polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

For each $0 \leq k \leq n$, we find a polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}'_k(x_k) = 1 \end{cases}$$

Hermite Interpolation

We would like to find a polynomial $H(x)$ of degree at most $2n + 1$ such that

$$\textcircled{H} \quad H(x_k) = y_k \quad \text{and} \quad H'(x_k) = z_k \quad \text{for all } 0 \leq k \leq n.$$

For each $0 \leq k \leq n$, we find a polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

For each $0 \leq k \leq n$, we find a polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}'_k(x_k) = 1 \end{cases}$$

Then $H(x) = y_0 H_0(x) + \dots + y_n H_n(x) + z_0 \hat{H}_0(x) + \dots + z_n \hat{H}_n(x)$ is a polynomial of degree at most $2n + 1$ satisfying the conditions \textcircled{H} above.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 \\ H'_k(x_j) = 0 \end{cases} \quad \begin{array}{l} \text{for all } 0 \leq j \leq n, j \neq k \\ \text{for all } 0 \leq j \leq n \end{array}$$

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)}{(x_k-x_j)} = \frac{(x-x_0)\dots(x-x_{k-1})(x-x_{k+1})\dots(x-x_n)}{(x_k-x_0)\dots(x_k-x_{k-1})(x_k-x_{k+1})\dots(x_k-x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $L_k^2(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)^2}{(x_k-x_j)^2} = \frac{(x-x_0)^2\dots(x-x_{k-1})^2(x-x_{k+1})^2\dots(x-x_n)^2}{(x_k-x_0)^2\dots(x_k-x_{k-1})^2(x_k-x_{k+1})^2\dots(x_k-x_n)^2}$, of degree $2n$, satisfies $L_k^2(x_k) = 1$ and $L_k^2(x_j) = 0$ and $(L_k^2)'(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$ (all $x_j, j \neq k$ are double roots of $L_k^2(x)$).

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)}{(x_k-x_j)} = \frac{(x-x_0)\dots(x-x_{k-1})(x-x_{k+1})\dots(x-x_n)}{(x_k-x_0)\dots(x_k-x_{k-1})(x_k-x_{k+1})\dots(x_k-x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $L_k^2(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)^2}{(x_k-x_j)^2} = \frac{(x-x_0)^2\dots(x-x_{k-1})^2(x-x_{k+1})^2\dots(x-x_n)^2}{(x_k-x_0)^2\dots(x_k-x_{k-1})^2(x_k-x_{k+1})^2\dots(x_k-x_n)^2}$, of degree $2n$, satisfies $L_k^2(x_k) = 1$ and $L_k^2(x_j) = 0$ and $(L_k^2)'(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$ (all $x_j, j \neq k$ are double roots of $L_k^2(x)$).

Polynomial $H_k(x) = (A(x-x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k-x_k) + 1)L_k^2(x_k) = 1$

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)}{(x_k-x_j)} = \frac{(x-x_0)\dots(x-x_{k-1})(x-x_{k+1})\dots(x-x_n)}{(x_k-x_0)\dots(x_k-x_{k-1})(x_k-x_{k+1})\dots(x_k-x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $L_k^2(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x-x_j)^2}{(x_k-x_j)^2} = \frac{(x-x_0)^2 \dots (x-x_{k-1})^2 (x-x_{k+1})^2 \dots (x-x_n)^2}{(x_k-x_0)^2 \dots (x_k-x_{k-1})^2 (x_k-x_{k+1})^2 \dots (x_k-x_n)^2}$, of degree $2n$, satisfies $L_k^2(x_k) = 1$ and $L_k^2(x_j) = 0$ and $(L_k^2)'(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$ (all $x_j, j \neq k$ are double roots of $L_k^2(x)$).

Polynomial $H_k(x) = (A(x-x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k-x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j-x_k) + 1)L_k^2(x_j) = 0$.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $L_k^2(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)^2}{(x_k - x_j)^2} = \frac{(x - x_0)^2 \dots (x - x_{k-1})^2 (x - x_{k+1})^2 \dots (x - x_n)^2}{(x_k - x_0)^2 \dots (x_k - x_{k-1})^2 (x_k - x_{k+1})^2 \dots (x_k - x_n)^2}$, of degree $2n$, satisfies $L_k^2(x_k) = 1$ and $L_k^2(x_j) = 0$ and $(L_k^2)'(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$ (all $x_j, j \neq k$ are double roots of $L_k^2(x)$).

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.
 $H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x)$.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.
 $H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x)$.

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.

$$H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x).$$

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Additionally, $H'_k(x_k) = AL_k^2(x_k) + (A(x_k - x_k) + 1)2L_k(x_k)L'_k(x_k)$

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.

$$H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x).$$

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Additionally,

$$H'_k(x_k) = AL_k^2(x_k) + (A(x_k - x_k) + 1)2L_k(x_k)L'_k(x_k) = A(1)^2 + (A(0) + 1)2(1)L'_k(x_k)$$

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.
 $H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x)$.

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Additionally, $H'_k(x_k) = AL_k^2(x_k) + (A(x_k - x_k) + 1)2L_k(x_k)L'_k(x_k) = A(1)^2 + (A(0) + 1)2(1)L'_k(x_k) = A + 2L'_k(x_k)$.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\left\{ \begin{array}{ll} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{array} \right.$$

Polynomial $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$, of degree n , satisfies $L_k(x_k) = 1$ and $L_k(x_j) = 0$ for all $0 \leq j \leq n, j \neq k$.

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.
 $H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x)$.

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Additionally, $H'_k(x_k) = AL_k^2(x_k) + (A(x_k - x_k) + 1)2L_k(x_k)L'_k(x_k) = A(1)^2 + (A(0) + 1)2(1)L'_k(x_k) = A + 2L'_k(x_k)$. Therefore, to make sure $H'_k(x_k) = 0$, we choose $A = -2L'_k(x_k)$.

Construction of $H_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $H_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} H_k(x_k) = 1 \\ H_k(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ H'_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \end{cases}$$

Polynomial $H_k(x) = (A(x - x_k) + 1)L_k^2(x)$, of degree $2n + 1$, satisfies $H_k(x_k) = (A(x_k - x_k) + 1)L_k^2(x_k) = 1$ and $H_k(x_j) = (A(x_j - x_k) + 1)L_k^2(x_j) = 0$.
 $H'_k(x) = AL_k^2(x) + (A(x - x_k) + 1)2L_k(x)L'_k(x)$.

For all $0 \leq j \leq n, j \neq k$, $H'_k(x_j) = AL_k^2(x_j) + (A(x_j - x_k) + 1)2L_k(x_j)L'_k(x_j) = 0$

Additionally, $H'_k(x_k) = AL_k^2(x_k) + (A(x_k - x_k) + 1)2L_k(x_k)L'_k(x_k) = A(1)^2 + (A(0) + 1)2(1)L'_k(x_k) = A + 2L'_k(x_k)$. Therefore, to make sure $H'_k(x_k) = 0$, we choose $A = -2L'_k(x_k)$.

Therefore, $H_k(x) = (-2L'_k(x_k)(x - x_k) + 1)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \left\{ \begin{array}{ll} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{array} \right.$$

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

$\hat{H}_k(x)$ has one factor of $x - x_k$ and two factors of $(x - x_j)$ for all $j \neq k$.

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

$\hat{H}_k(x)$ has one factor of $x - x_k$ and two factors of $(x - x_j)$ for all $j \neq k$.
Therefore, $\hat{H}_k(x_j) = 0$ for all $1 \leq j \leq n$.

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

$\hat{H}_k(x)$ has one factor of $x - x_k$ and two factors of $(x - x_j)$ for all $j \neq k$.
Therefore, $\hat{H}_k(x_j) = 0$ for all $1 \leq j \leq n$.

$$\hat{H}_k'(x) = L_k^2(x) + 2(x - x_k)L_k(x)L_k'(x).$$

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

$\hat{H}_k(x)$ has one factor of $x - x_k$ and two factors of $(x - x_j)$ for all $j \neq k$.
Therefore, $\hat{H}_k(x_j) = 0$ for all $1 \leq j \leq n$.

$$\hat{H}_k'(x) = L_k^2(x) + 2(x - x_k)L_k(x)L_k'(x).$$

$$\hat{H}_k'(x_j) = L_k^2(x_j) + 2(x_j - x_k)L_k(x_j)L_k'(x_j) = 0, \text{ when } j \neq k.$$

Construction of $\hat{H}_k(x)$

For each $0 \leq k \leq n$, we would like to find the polynomial $\hat{H}_k(x)$ of degree $2n + 1$ satisfying

$$\blacksquare \begin{cases} \hat{H}_k(x_j) = 0 & \text{for all } 0 \leq j \leq n \\ \hat{H}_k'(x_j) = 0 & \text{for all } 0 \leq j \leq n, j \neq k \\ \hat{H}_k'(x_k) = 1 \end{cases}$$

$\hat{H}_k(x) = (x - x_k)L_k^2(x)$, where $L_k(x) = \prod_{\substack{1 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$ is the polynomial of degree $2n + 1$ satisfying the conditions \blacksquare .

$\hat{H}_k(x)$ has one factor of $x - x_k$ and two factors of $(x - x_j)$ for all $j \neq k$.
Therefore, $\hat{H}_k(x_j) = 0$ for all $1 \leq j \leq n$.

$$\hat{H}_k'(x) = L_k^2(x) + 2(x - x_k)L_k(x)L_k'(x).$$

$$\hat{H}_k'(x_j) = L_k^2(x_j) + 2(x_j - x_k)L_k(x_j)L_k'(x_j) = 0, \text{ when } j \neq k.$$

$$\hat{H}_k'(x_k) = L_k^2(x_k) + 2(x_k - x_k)L_k(x_k)L_k'(x_k) = (1)^2 + 0 = 1.$$

Example

Find the Hermite interpolation polynomial for the following data:

x	1.1	1.3	1.5
$f(x)$	0.45	0.27	0.07
$f'(x)$	-0.89	-0.96	-1

Error Estimate For Hermite Interpolation

Suppose $f \in C^{2n+1}[a, b]$ and $f^{(2n+2)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$.

Error Estimate For Hermite Interpolation

Suppose $f \in C^{2n+1}[a, b]$ and $f^{(2n+2)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Let $H_n(x)$ be the Hermite polynomial of degree at most $2n + 1$ satisfying

$$H_n(x_k) = f(x_k) \quad \text{and} \quad H'_n(x_k) = f'(x_k) \quad \text{for all } 0 \leq k \leq n.$$

Error Estimate For Hermite Interpolation

Suppose $f \in C^{2n+1}[a, b]$ and $f^{(2n+2)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Let $H_n(x)$ be the Hermite polynomial of degree at most $2n + 1$ satisfying

$$H_n(x_k) = f(x_k) \quad \text{and} \quad H'_n(x_k) = f'(x_k) \quad \text{for all } 0 \leq k \leq n.$$

Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - H_n(x) = \frac{f^{(2n+2)}(\zeta(x))}{(2n+2)!} (x - x_0)^2 (x - x_1)^2 \dots (x - x_n)^2$$