

# Linear Difference Equations and Bernoulli's Method

MATH 4701 Numerical Analysis

# Linear Difference Equations

A **linear difference equation of order  $n$**  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

A solution of this equation is a **sequence**  $(u_i)$  of numbers.

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

A solution of this equation is a **sequence**  $(u_i)$  of numbers.

A solution of the equation is uniquely determined by any choice of  $n$  (the order of the equation) **initial conditions**  $u_0, u_1, \dots, u_{n-1}$ .

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

A solution of this equation is a **sequence**  $(u_i)$  of numbers.

A solution of the equation is uniquely determined by any choice of  $n$  (the order of the equation) **initial conditions**  $u_0, u_1, \dots, u_{n-1}$ .

**Example (1):** The **Fibonacci** difference equation

$$u_k - u_{k-1} - u_{k-2} = 0$$

is a difference equation of **order two**.

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

A solution of this equation is a **sequence**  $(u_i)$  of numbers.

A solution of the equation is uniquely determined by any choice of  $n$  (the order of the equation) **initial conditions**  $u_0, u_1, \dots, u_{n-1}$ .

**Example (1):** The **Fibonacci** difference equation

$$u_k - u_{k-1} - u_{k-2} = 0 \Rightarrow u_k = u_{k-1} + u_{k-2}$$

is a difference equation of **order two**.

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (1):** The **Fibonacci** difference equation

$$u_k - u_{k-1} - u_{k-2} = 0 \Rightarrow u_k = u_{k-1} + u_{k-2}$$

is a difference equation of **order two**.

With initial conditions  $u_0 = 1, u_1 = 1$  we get the solution

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, \dots$$

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (1):** The **Fibonacci** difference equation

$$u_k - u_{k-1} - u_{k-2} = 0 \Rightarrow u_k = u_{k-1} + u_{k-2}$$

is a difference equation of **order two**.

With initial conditions  $u_0 = 1, u_1 = 1$  we get the solution

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, \dots$$

With initial conditions  $u_0 = -1, u_1 = 4$  we get the solution

$$-1, 4, 3, 7, 10, 17, 27, 44, 71, 115, 186, \dots$$



# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$$

is a linear difference equation of **order three**.

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n+1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0 \Rightarrow u_k = 4u_{k-1} - 3u_{k-2} - u_{k-3}$$

is a linear difference equation of **order three**.

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0 \Rightarrow u_k = 4u_{k-1} - 3u_{k-2} - u_{k-3}$$

is a linear difference equation of **order three**.

With initial conditions  $u_0 = 1$ ,  $u_1 = 3$ ,  $u_2 = -1$  we get the solution

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0 \Rightarrow u_k = 4u_{k-1} - 3u_{k-2} - u_{k-3}$$

is a linear difference equation of **order three**.

With initial conditions  $u_0 = 1, u_1 = 3, u_2 = -1$  we get the solution

$$1, 3, -1, -14, -56, -181, \dots$$

# Linear Difference Equations

A **linear difference equation of order  $n$**  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0 \Rightarrow u_k = 4u_{k-1} - 3u_{k-2} - u_{k-3}$$

is a linear difference equation of **order three**.

With initial conditions  $u_0 = 1, u_1 = 3, u_2 = -1$  we get the solution

$$1, 3, -1, -14, -56, -181, \dots$$

With initial conditions  $u_0 = -1, u_1 = 1, u_2 = -1$  we get the solution

# Linear Difference Equations

A **linear difference equation** of **order**  $n$  is an equation of the form

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

where  $a_0, a_1, \dots, a_n$  are  $n + 1$  constants and  $a_n \neq 0$ .

**Example (2):** The difference equation

$$u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0 \Rightarrow u_k = 4u_{k-1} - 3u_{k-2} - u_{k-3}$$

is a linear difference equation of **order three**.

With initial conditions  $u_0 = 1, u_1 = 3, u_2 = -1$  we get the solution

$$1, 3, -1, -14, -56, -181, \dots$$

With initial conditions  $u_0 = -1, u_1 = 1, u_2 = -1$  we get the solution

$$-1, 1, -1, -6, -22, -69, \dots$$

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.



# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

All entries of a sequence can be multiplied by a number to define the  
scalar product of the vectors with numbers.

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

All entries of a sequence can be multiplied by a number to define the scalar product of the vectors with numbers. For example if  $\mathbf{u} = (\frac{1}{n}) = (1, \frac{1}{2}, \frac{1}{3}, \dots)$ , then  $2\mathbf{u} = (\frac{2}{n}) = (2, 1, \frac{2}{3}, \dots)$ .

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

All entries of a sequence can be multiplied by a number to define the scalar product of the vectors with numbers. For example if

$$\mathbf{u} = \left(\frac{1}{n}\right) = \left(1, \frac{1}{2}, \frac{1}{3}, \dots\right), \text{ then } 2\mathbf{u} = \left(\frac{2}{n}\right) = \left(2, 1, \frac{2}{3}, \dots\right).$$

The vector addition is defined by adding the two sequences term by term.

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

All entries of a sequence can be multiplied by a number to define the scalar product of the vectors with numbers. For example if  $\mathbf{u} = (\frac{1}{n}) = (1, \frac{1}{2}, \frac{1}{3}, \dots)$ , then  $2\mathbf{u} = (\frac{2}{n}) = (2, 1, \frac{2}{3}, \dots)$ .

The vector addition is defined by adding the two sequences term by term. For example if  $\mathbf{u}_1 = (2^n) = (2, 4, 8, 16, \dots)$  and  $\mathbf{u}_2 = ((-1)^n) = (-1, 1, -1, 1, \dots)$ , then  $\mathbf{u}_1 + \mathbf{u}_2 = (2^n + (-1)^n) = (1, 5, 7, 17, \dots)$ .

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**.

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**. vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  
vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$   
(associative property)

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**. vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$  where  $\mathbf{0}$  is the **constant zero sequence**.



# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**. vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$   
where  $\mathbf{0}$  is the **constant zero sequence**.
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$   
where  $-\mathbf{u}$  is the sequence with each entry equal to the opposite of the corresponding entry of  $\mathbf{u}$ .

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**. vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$  where  $\mathbf{0}$  is the **constant zero sequence**.
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$  where  $-\mathbf{u}$  is the sequence with each entry equal to the opposite of the corresponding entry of  $\mathbf{u}$ . For example if  $\mathbf{u} = (\frac{1}{n}) = (1, \frac{1}{2}, \frac{1}{3}, \dots)$ , then  $-\mathbf{u} = (-\frac{1}{n}) = (-1, -\frac{1}{2}, -\frac{1}{3}, \dots)$ .

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**. vectors in this space are the sequences.

$\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$  where  $\mathbf{0}$  is the **constant zero sequence**.
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$  where  $-\mathbf{u}$  is the sequence with each entry equal to the opposite of the corresponding entry of  $\mathbf{u}$ .
- For every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v}$

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  $\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$  where  $\mathbf{0}$  is the **constant zero sequence**.
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$  where  $-\mathbf{u}$  is the sequence with each entry equal to the opposite of the corresponding entry of  $\mathbf{u}$ .
- For every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v}$
- For every scalars  $\alpha$  and  $\beta$  and every vector  $\mathbf{u}$ ,  $(\alpha + \beta)\mathbf{u} = \alpha\mathbf{u} + \beta\mathbf{u}$

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  $\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$
- For every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v}$
- For every scalars  $\alpha$  and  $\beta$  and every vector  $\mathbf{u}$ ,  $(\alpha + \beta)\mathbf{u} = \alpha\mathbf{u} + \beta\mathbf{u}$
- For every scalars  $\alpha$  and  $\beta$  and every vector  $\mathbf{u}$ ,  $(\alpha\beta)\mathbf{u} = \alpha(\beta\mathbf{u})$

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  $\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**. This means

- For every vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  (commutative property)
- For every vectors  $\mathbf{u}$ ,  $\mathbf{v}$ , and  $\mathbf{w}$ ,  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  (associative property)
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
- For every vector  $\mathbf{u}$ ,  $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$
- For every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\alpha(\mathbf{u} + \mathbf{v}) = \alpha\mathbf{u} + \alpha\mathbf{v}$
- For every scalars  $\alpha$  and  $\beta$  and every vector  $\mathbf{u}$ ,  $(\alpha + \beta)\mathbf{u} = \alpha\mathbf{u} + \beta\mathbf{u}$
- For every scalars  $\alpha$  and  $\beta$  and every vector  $\mathbf{u}$ ,  $(\alpha\beta)\mathbf{u} = \alpha(\beta\mathbf{u})$
- For every vector  $\mathbf{u}$ ,  $1\mathbf{u} = \mathbf{u}$

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  $\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**.

A subset of  $\mathcal{S}$  is called a **linear subspace** of  $\mathcal{S}$  if with the scalar product and vector addition of  $\mathcal{S}$  it also forms a vector space.

# Linear Subspace of Solutions of a Linear Difference Eq.

The space of sequences  $\mathcal{S}$  of numbers can be regarded as a **vector space**.  $\mathcal{S}$  together with the scalar product and vector addition we defined forms a **vector space**.

A subset of  $\mathcal{S}$  is called a **linear subspace** of  $\mathcal{S}$  if with the scalar product and vector addition of  $\mathcal{S}$  it also forms a vector space.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .



# Linear Subspace of Solutions of a Linear Difference Eq.

A subset of  $\mathcal{S}$  is called a **linear subspace** of  $\mathcal{S}$  if with the scalar product and vector addition of  $\mathcal{S}$  it also forms a vector space.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then

$$a_n x_k + a_{n-1} x_{k-1} + \dots + a_1 x_{k-n+1} + a_0 x_{k-n} = 0 \text{ and}$$

$$a_n y_k + a_{n-1} y_{k-1} + \dots + a_1 y_{k-n+1} + a_0 y_{k-n} = 0$$

# Linear Subspace of Solutions of a Linear Difference Eq.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then

$$a_n x_k + a_{n-1} x_{k-1} + \dots + a_1 x_{k-n+1} + a_0 x_{k-n} = 0 \text{ and}$$

$$a_n y_k + a_{n-1} y_{k-1} + \dots + a_1 y_{k-n+1} + a_0 y_{k-n} = 0$$

Multiplying the first equation by  $\alpha$  and regrouping we get

$$a_n(\alpha x_k) + a_{n-1}(\alpha x_{k-1}) + \dots + a_1(\alpha x_{k-n+1}) + a_0(\alpha x_{k-n}) = 0$$

# Linear Subspace of Solutions of a Linear Difference Eq.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then

$$a_n x_k + a_{n-1} x_{k-1} + \dots + a_1 x_{k-n+1} + a_0 x_{k-n} = 0 \text{ and}$$

$$a_n y_k + a_{n-1} y_{k-1} + \dots + a_1 y_{k-n+1} + a_0 y_{k-n} = 0$$

Multiplying the first equation by  $\alpha$  and regrouping we get

$$a_n(\alpha x_k) + a_{n-1}(\alpha x_{k-1}) \dots + a_1(\alpha x_{k-n+1}) + a_0(\alpha x_{k-n}) = 0$$

Adding the two equations and regrouping we get

$$a_n(x_k + y_k) + a_{n-1}(x_{k-1} + y_{k-1}) \dots + a_0(x_{k-n} + y_{k-n}) = 0$$

# Linear Subspace of Solutions of a Linear Difference Eq.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then Multiplying the first equation by  $\alpha$  and regrouping we get

$$a_n(\alpha x_k) + a_{n-1}(\alpha x_{k-1}) \dots + a_1(\alpha x_{k-n+1}) + a_0(\alpha x_{k-n}) = 0$$

Adding the two equations and regrouping we get

$$a_n(x_k + y_k) + a_{n-1}(x_{k-1} + y_{k-1}) \dots + a_0(x_{k-n} + y_{k-n}) = 0$$

This means for solutions  $(x_i)$  and  $(y_i)$  of the equation both  $\alpha(x_i)$  and  $(x_i) + (y_i)$  are also solutions of the difference equation.

# Linear Subspace of Solutions of a Linear Difference Eq.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then This means for solutions  $(x_i)$  and  $(y_i)$  of the equation both  $\alpha(x_i)$  and  $(x_i) + (y_i)$  are also solutions of the difference equation.

Moreover, since with any choice of  $n$  initial conditions a solution can be generated, the set of solutions to the difference equation is not empty.

# Linear Subspace of Solutions of a Linear Difference Eq.

To check if a subset  $\mathcal{A}$  of  $\mathcal{S}$  is a linear subspace it is sufficient to verify that it is **non-empty** and for every scalar  $\alpha$  and vectors  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathcal{A}$ ,  $\alpha\mathbf{u}$  and  $\mathbf{u} + \mathbf{v}$  are also in  $\mathcal{A}$ .

For any linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

if  $(x_i)$  and  $(y_i)$  are two solutions of the equation and  $\alpha$  is any number then This means for solutions  $(x_i)$  and  $(y_i)$  of the equation both  $\alpha(x_i)$  and  $(x_i) + (y_i)$  are also solutions of the difference equation.

Moreover, since with any choice of  $n$  initial conditions a solution can be generated, the set of solutions to the difference equation is not empty.

Hence, **the set of solutions of a linear difference equation forms a linear subspace of  $\mathcal{S}$ .**

# A Finite Basis For the Subspace of Solutions

For every linear difference equation of order  $n$

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

a polynomial of degree  $n$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

can be considered.

# A Finite Basis For the Subspace of Solutions

For every linear difference equation of order  $n$

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

a polynomial of degree  $n$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

can be considered.

This polynomial is called the **characteristic polynomial** of the difference equation.



# A Finite Basis For the Subspace of Solutions

For every linear difference equation of order  $n$

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

a polynomial of degree  $n$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

can be considered.

This polynomial is called the **characteristic polynomial** of the difference equation.

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

# A Finite Basis For the Subspace of Solutions

For every linear difference equation of order  $n$

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

a polynomial of degree  $n$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

can be considered.

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$

# A Finite Basis For the Subspace of Solutions

For every linear difference equation of order  $n$

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

a polynomial of degree  $n$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

can be considered.

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ .

# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation.

# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

For example, the solution with initial conditions  $u_0 = 1$  and  $u_1 = 2$  is the sequence

# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

For example, the solution with initial conditions  $u_0 = 1$  and  $u_1 = 2$  is the sequence

$$1, 2, 4, 8, 16, 32, \dots$$

which is  $\mathbf{u}_1$  itself and can be obtained by letting  $\alpha = 1$  and  $\beta = 0$  in  $u_i = \alpha 2^i + \beta (-3)^i$ .

# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence



# A Finite Basis For the Subspace of Solutions

If characteristic polynomial  $P(x)$  has  $n$  distinct roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , then the  $n$  sequences  $\mathbf{u}_1 = (\lambda_1^i)_{i \in \mathbb{N}}$ ,  $\mathbf{u}_2 = (\lambda_2^i)_{i \in \mathbb{N}}$ , ...,  $\mathbf{u}_n = (\lambda_n^i)_{i \in \mathbb{N}}$  form a basis for the linear subspace of all solutions of the difference equation.

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta (-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta (-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

$$\begin{cases} i = 0 \Rightarrow \alpha 2^0 + \beta (-3)^0 = -1 \\ i = 1 \Rightarrow \alpha 2^1 + \beta (-3)^1 = 1 \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta (-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

$$\begin{cases} i = 0 \Rightarrow \alpha 2^0 + \beta (-3)^0 = -1 \\ i = 1 \Rightarrow \alpha 2^1 + \beta (-3)^1 = 1 \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . The sequences  $\mathbf{u}_1 = (2^i)_{i \in \mathbb{N}}$  and  $\mathbf{u}_2 = ((-3)^i)_{i \in \mathbb{N}}$  form a basis for all solutions of the equation. As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta (-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta (-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

$$\begin{cases} i = 0 \Rightarrow \alpha 2^0 + \beta (-3)^0 = -1 \\ i = 1 \Rightarrow \alpha 2^1 + \beta (-3)^1 = 1 \end{cases} \Rightarrow \begin{cases} \alpha + \beta = -1 \\ 2\alpha - 3\beta = 1 \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta(-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta(-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

$$\Rightarrow \begin{cases} \alpha + \beta = -1 \\ 2\alpha - 3\beta = 1 \end{cases}$$

Adding  $-2$  times the first equation to the second equation we get  $-5\beta = 3$  or  $\beta = -\frac{3}{5}$ .

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta(-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta(-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

$$\Rightarrow \begin{cases} \alpha + \beta = -1 \\ 2\alpha - 3\beta = 1 \end{cases}$$

Adding  $-2$  times the first equation to the second equation we get  $-5\beta = 3$  or  $\beta = -\frac{3}{5}$ . Letting  $\beta = -\frac{3}{5}$  we get  $\alpha = -\frac{2}{5}$ .

# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta(-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta(-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

Adding  $-2$  times the first equation to the second equation we get  $-5\beta = 3$  or  $\beta = -\frac{3}{5}$ . Letting  $\beta = -\frac{3}{5}$  we get  $\alpha = -\frac{2}{5}$ . The particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 2$  can be expressed explicitly by  $u_i = -\frac{2}{5}2^i - \frac{3}{5}(-3)^i$ .



# A Finite Basis For the Subspace of Solutions

**Example (1):** The difference equation  $u_k + u_{k-1} - 6u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 + x - 6$  which has roots  $\lambda_1 = 2$  and  $\lambda_2 = -3$ . As a result, the general solution of this difference equation is in the form  $u_i = \alpha 2^i + \beta(-3)^i$ .

On the other hand, the solution with initial conditions  $u_0 = -1$  and  $u_1 = 1$  is the sequence

$$-1, 1, -7, 13, -55, 133, \dots$$

To see which values of  $\alpha$  and  $\beta$  in  $u_i = \alpha 2^i + \beta(-3)^i$  generate this solution we let  $i = 0$  and  $i = 1$  to get the following system of two linear equations in two variables.

Adding  $-2$  times the first equation to the second equation we get  $-5\beta = 3$  or  $\beta = -\frac{3}{5}$ . Letting  $\beta = -\frac{3}{5}$  we get  $\alpha = -\frac{2}{5}$ . The particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 2$  can be expressed explicitly by  $u_i = -\frac{2}{5}2^i - \frac{3}{5}(-3)^i$ . For example, we have  $u_5 = -\frac{2}{5}2^5 - \frac{3}{5}(-3)^5 = 133$ .

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ .

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ .

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\begin{cases} i = 0 \Rightarrow \alpha\lambda_1^0 + \beta\lambda_2^0 = 1 \\ \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\begin{cases} i = 0 \Rightarrow \alpha\lambda_1^0 + \beta\lambda_2^0 = 1 \\ i = 1 \Rightarrow \alpha\lambda_1^1 + \beta\lambda_2^1 = 1 \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\begin{cases} i = 0 \Rightarrow \alpha\lambda_1^0 + \beta\lambda_2^0 = 1 \\ i = 1 \Rightarrow \alpha\lambda_1^1 + \beta\lambda_2^1 = 1 \end{cases} \Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1$



# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5}$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5}$$

$$1 - \lambda_1 = 1 - \frac{1+\sqrt{5}}{2} = \frac{1-\sqrt{5}}{2} = \lambda_2$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1 \Rightarrow -\sqrt{5}\beta = \lambda_2$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5}$$

$$1 - \lambda_1 = 1 - \frac{1+\sqrt{5}}{2} = \frac{1-\sqrt{5}}{2} = \lambda_2$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1 \Rightarrow -\sqrt{5}\beta = \lambda_2 \Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5}$$

$$1 - \lambda_1 = 1 - \frac{1+\sqrt{5}}{2} = \frac{1-\sqrt{5}}{2} = \lambda_2$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1 \Rightarrow -\sqrt{5}\beta = \lambda_2 \Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5} \Rightarrow -\frac{\lambda_2}{\sqrt{5}} + \frac{\lambda_1}{\sqrt{5}} = 1$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1 \Rightarrow -\sqrt{5}\beta = \lambda_2 \Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5} \Rightarrow -\frac{\lambda_2}{\sqrt{5}} + \frac{\lambda_1}{\sqrt{5}} = 1 \Rightarrow \beta + \frac{\lambda_1}{\sqrt{5}} = 1$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

Multiplying the first equation by  $-\lambda_1$  and adding it to the second equation we get  $(\lambda_2 - \lambda_1)\beta = 1 - \lambda_1 \Rightarrow -\sqrt{5}\beta = \lambda_2 \Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$

$$\lambda_2 - \lambda_1 = \frac{1-\sqrt{5}}{2} - \frac{1+\sqrt{5}}{2} = -\sqrt{5} \Rightarrow -\frac{\lambda_2}{\sqrt{5}} + \frac{\lambda_1}{\sqrt{5}} = 1 \Rightarrow \beta + \frac{\lambda_1}{\sqrt{5}} = 1$$

Therefore, from the first equation, we have  $\alpha = 1 - \beta = \frac{\lambda_1}{\sqrt{5}}$ .

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

$$\Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$$

$$\alpha = 1 - \beta = \frac{\lambda_1}{\sqrt{5}}.$$

This means the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  can be expressed by the explicit formula

$$u_i = \alpha\lambda_1^i + \beta\lambda_2^i = \frac{\lambda_1}{\sqrt{5}}\lambda_1^i - \frac{\lambda_2}{\sqrt{5}}\lambda_2^i$$



# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \begin{cases} \alpha + \beta = 1 \\ \lambda_1\alpha + \lambda_2\beta = 1 \end{cases}$$

$$\Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$$

$$\alpha = 1 - \beta = \frac{\lambda_1}{\sqrt{5}}.$$

This means the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  can be expressed by the explicit formula

$$u_i = \alpha\lambda_1^i + \beta\lambda_2^i = \frac{\lambda_1}{\sqrt{5}}\lambda_1^i - \frac{\lambda_2}{\sqrt{5}}\lambda_2^i = \frac{\lambda_1^{i+1} - \lambda_2^{i+1}}{\sqrt{5}}$$

# A Finite Basis For the Subspace of Solutions

**Example (2):** The **Fibonacci** difference equation  $u_k - u_{k-1} - u_{k-2} = 0$  has characteristic polynomial  $P(x) = x^2 - x - 1$  which has roots  $\lambda_1 = \frac{1+\sqrt{5}}{2}$  and  $\lambda_2 = \frac{1-\sqrt{5}}{2}$ . This means every solution of the Fibonacci difference equation is in the form  $u_i = \alpha\lambda_1^i + \beta\lambda_2^i$ . To find which  $\alpha$  and  $\beta$  generate the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  we let  $i = 0$  and  $i = 1$  to get

$$\Rightarrow \beta = -\frac{\lambda_2}{\sqrt{5}}$$

$$\alpha = 1 - \beta = \frac{\lambda_1}{\sqrt{5}}.$$

This means the particular solution with initial conditions  $u_0 = 1$  and  $u_1 = 1$  can be expressed by the explicit formula

$$u_i = \frac{\lambda_1^{i+1} - \lambda_2^{i+1}}{\sqrt{5}}$$

For example  $u_7 = \frac{\lambda_1^8 - \lambda_2^8}{\sqrt{5}} = 21$  which is the 7th entry in the Fibonacci sequence.

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

$$i = 0 \Rightarrow \alpha + \beta + \gamma = 2$$

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

$$i = 0 \Rightarrow \alpha + \beta + \gamma = 2$$

$$i = 1 \Rightarrow -\alpha - 2\beta + 3\gamma = -7$$

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

$$i = 0 \Rightarrow \alpha + \beta + \gamma = 2$$

$$i = 1 \Rightarrow -\alpha - 2\beta + 3\gamma = -7$$

$$i = 2 \Rightarrow \alpha + 4\beta + 9\gamma = -3$$



# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

$$i = 0 \Rightarrow \alpha + \beta + \gamma = 2$$

$$i = 1 \Rightarrow -\alpha - 2\beta + 3\gamma = -7$$

$$i = 2 \Rightarrow \alpha + 4\beta + 9\gamma = -3$$

$$\Rightarrow \alpha = 2, \beta = 1, \text{ and } \gamma = -1$$

# A Finite Basis For the Subspace of Solutions

**Example (3):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k - 7u_{k-2} - 6u_{k-3} = 0$$

with initial conditions  $u_0 = 2$ ,  $u_1 = -7$ ,  $u_2 = -3$ .

Characteristic polynomial  $P(x) = x^3 - 7x - 6 = (x + 1)(x + 2)(x - 3)$ .

General solution can be represented by  $u_i = \alpha(-1)^i + \beta(-2)^i + \gamma 3^i$ .

$$i = 0 \Rightarrow \alpha + \beta + \gamma = 2$$

$$i = 1 \Rightarrow -\alpha - 2\beta + 3\gamma = -7$$

$$i = 2 \Rightarrow \alpha + 4\beta + 9\gamma = -3$$

$$\Rightarrow \alpha = 2, \beta = 1, \text{ and } \gamma = -1$$

$$u_i = 2(-1)^i + (-2)^i - 3^i.$$

# Multiple Roots of Characteristic Polynomial

When characteristic polynomial  $P(x)$  of a linear difference equation has a root  $\delta$  of multiplicity  $m$  then this root generates  $m$  basis solutions in the form  $\mathbf{x}_k = (i^k \delta^i)$  for  $k = 0, 1, \dots, m - 1$  (assuming  $0^0 = 1$ ).

# Multiple Roots of Characteristic Polynomial

When characteristic polynomial  $P(x)$  of a linear difference equation has a root  $\delta$  of multiplicity  $m$  then this root generates  $m$  basis solutions in the form  $\mathbf{x}_k = (i^k \delta^i)$  for  $k = 0, 1, \dots, m - 1$  (assuming  $0^0 = 1$ ).

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

# Multiple Roots of Characteristic Polynomial

When characteristic polynomial  $P(x)$  of a linear difference equation has a root  $\delta$  of multiplicity  $m$  then this root generates  $m$  basis solutions in the form  $\mathbf{x}_k = (i^k \delta^i)$  for  $k = 0, 1, \dots, m - 1$  (assuming  $0^0 = 1$ ).

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3, u_1 = 6, u_2 = 17, u_3 = 42, u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

# Multiple Roots of Characteristic Polynomial

When characteristic polynomial  $P(x)$  of a linear difference equation has a root  $\delta$  of multiplicity  $m$  then this root generates  $m$  basis solutions in the form  $\mathbf{x}_k = (i^k \delta^i)$  for  $k = 0, 1, \dots, m - 1$  (assuming  $0^0 = 1$ ).

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3, u_1 = 6, u_2 = 17, u_3 = 42, u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

General solution can be represented by

$$u_i = a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i$$

# Multiple Roots of Characteristic Polynomial

When characteristic polynomial  $P(x)$  of a linear difference equation has a root  $\delta$  of multiplicity  $m$  then this root generates  $m$  basis solutions in the form  $\mathbf{x}_k = (i^k \delta^i)$  for  $k = 0, 1, \dots, m - 1$  (assuming  $0^0 = 1$ ).

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3, u_1 = 6, u_2 = 17, u_3 = 42, u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$



# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$

$$i = 1 \Rightarrow a + b + c - 2d - 2e = 6$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

Characteristic polynomial:

$$P(x) = x^5 + x^4 - 5x^3 - x^2 + 8x - 4 = (x - 1)^3(x + 2)^2.$$

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$

$$i = 1 \Rightarrow a + b + c - 2d - 2e = 6$$

$$i = 2 \Rightarrow a + 2b + 4c + 4d + 8e = 17$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$

$$i = 1 \Rightarrow a + b + c - 2d - 2e = 6$$

$$i = 2 \Rightarrow a + 2b + 4c + 4d + 8e = 17$$

$$i = 3 \Rightarrow a + 3b + 9c - 8d - 24e = 42$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$

$$i = 1 \Rightarrow a + b + c - 2d - 2e = 6$$

$$i = 2 \Rightarrow a + 2b + 4c + 4d + 8e = 17$$

$$i = 3 \Rightarrow a + 3b + 9c - 8d - 24e = 42$$

$$i = 4 \Rightarrow a + 4b + 16c + 16d + 64e = 95$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$i = 0 \Rightarrow a + d = 3$$

$$i = 1 \Rightarrow a + b + c - 2d - 2e = 6$$

$$i = 2 \Rightarrow a + 2b + 4c + 4d + 8e = 17$$

$$i = 3 \Rightarrow a + 3b + 9c - 8d - 24e = 42$$

$$i = 4 \Rightarrow a + 4b + 16c + 16d + 64e = 95$$

$$\Rightarrow a = \frac{113}{27}, b = -\frac{163}{27}, c = \frac{58}{9}, d = -\frac{32}{27}, e = \frac{13}{27}.$$

# Multiple Roots of Characteristic Polynomial

**Example (4):** Find an explicit description for terms of the solution of the linear difference equation

$$u_k + u_{k-1} - 5u_{k-2} - u_{k-3} + 8u_{k-4} - 4u_{k-5} = 0$$

with initial conditions  $u_0 = 3$ ,  $u_1 = 6$ ,  $u_2 = 17$ ,  $u_3 = 42$ ,  $u_4 = 95$ .

General solution can be represented by

$$\begin{aligned} u_i &= a(1)^i + bi(1)^i + ci^2(1)^i + d(-2)^i + ei(-2)^i \\ &= a + bi + ci^2 + d(-2)^i + ei(-2)^i \end{aligned}$$

$$\Rightarrow a = \frac{113}{27}, b = -\frac{163}{27}, c = \frac{58}{9}, d = -\frac{32}{27}, e = \frac{13}{27}.$$

$$\Rightarrow u_i = \frac{113}{27} - \frac{163}{27}i + \frac{58}{9}i^2 - \frac{32}{27}(-2)^i + \frac{13}{27}i2^i.$$

# Bernoulli's Method

Suppose  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  is a polynomial with  $n$  distinct real roots  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$$|\alpha_1| > |\alpha_2| \geq |\alpha_3| \geq \dots \geq |\alpha_n|$$

# Bernoulli's Method

Suppose  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  is a polynomial with  $n$  distinct real roots  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$$|\alpha_1| > |\alpha_2| \geq |\alpha_3| \geq \dots \geq |\alpha_n|$$

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .



# Bernoulli's Method

Suppose  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  is a polynomial with  $n$  distinct real roots  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$$|\alpha_1| > |\alpha_2| \geq |\alpha_3| \geq \dots \geq |\alpha_n|$$

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ . Therefore, every solution of it can be expressed as

$$u_i = c_1 \alpha_1^i + c_2 \alpha_2^i + \dots + c_n \alpha_n^i$$

# Bernoulli's Method

Suppose  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  is a polynomial with  $n$  distinct real roots  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$$|\alpha_1| > |\alpha_2| \geq |\alpha_3| \geq \dots \geq |\alpha_n|$$

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ . Therefore, every solution of it can be expressed as

$$u_i = c_1 \alpha_1^i + c_2 \alpha_2^i + \dots + c_n \alpha_n^i$$

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k}$$

# Bernoulli's Method

Suppose  $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  is a polynomial with  $n$  distinct real roots  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$$|\alpha_1| > |\alpha_2| \geq |\alpha_3| \geq \dots \geq |\alpha_n|$$

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ . Therefore, every solution of it can be expressed as

$$u_i = c_1 \alpha_1^i + c_2 \alpha_2^i + \dots + c_n \alpha_n^i$$

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} = \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} + c_2 \alpha_2^{k+1} + \dots + c_n \alpha_n^{k+1}}{c_1 \alpha_1^k + c_2 \alpha_2^k + \dots + c_n \alpha_n^k}$$

# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\begin{aligned}\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} &= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} + c_2 \alpha_2^{k+1} + \dots + c_n \alpha_n^{k+1}}{c_1 \alpha_1^k + c_2 \alpha_2^k + \dots + c_n \alpha_n^k} \\ &= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} \left(1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^{k+1} + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^{k+1}\right)}{c_1 \alpha_1^k \left(1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^k + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^k\right)}\end{aligned}$$

# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\begin{aligned}\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} &= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} + c_2 \alpha_2^{k+1} + \dots + c_n \alpha_n^{k+1}}{c_1 \alpha_1^k + c_2 \alpha_2^k + \dots + c_n \alpha_n^k} \\&= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} \left(1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^{k+1} + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^{k+1}\right)}{c_1 \alpha_1^k \left(1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^k + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^k\right)} \\&= \lim_{k \rightarrow \infty} \frac{\alpha_1 \left(1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^{k+1} + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^{k+1}\right)}{1 + \frac{c_2}{c_1} \left(\frac{\alpha_2}{\alpha_1}\right)^k + \dots + \frac{c_n}{c_1} \left(\frac{\alpha_n}{\alpha_1}\right)^k}\end{aligned}$$

Assuming  $c_1 \neq 0$ .

# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} &= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} + c_2 \alpha_2^{k+1} + \dots + c_n \alpha_n^{k+1}}{c_1 \alpha_1^k + c_2 \alpha_2^k + \dots + c_n \alpha_n^k} \\ &= \lim_{k \rightarrow \infty} \frac{\alpha_1 \left( 1 + \frac{c_2}{c_1} \left( \frac{\alpha_2}{\alpha_1} \right)^{k+1} + \dots + \frac{c_n}{c_1} \left( \frac{\alpha_n}{\alpha_1} \right)^{k+1} \right)}{1 + \frac{c_2}{c_1} \left( \frac{\alpha_2}{\alpha_1} \right)^k + \dots + \frac{c_n}{c_1} \left( \frac{\alpha_n}{\alpha_1} \right)^k} = \frac{\alpha_1 (1 + 0 + \dots + 0)}{1 + 0 + \dots + 0} \end{aligned}$$

For all  $j = 2, \dots, n$ , since  $|\alpha_1| > |\alpha_j|$ , we have  $\lim_{k \rightarrow \infty} \left( \frac{\alpha_j}{\alpha_1} \right)^k = 0$ .

# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} &= \lim_{k \rightarrow \infty} \frac{c_1 \alpha_1^{k+1} + c_2 \alpha_2^{k+1} + \dots + c_n \alpha_n^{k+1}}{c_1 \alpha_1^k + c_2 \alpha_2^k + \dots + c_n \alpha_n^k} \\ &= \lim_{k \rightarrow \infty} \frac{\alpha_1 \left( 1 + \frac{c_2}{c_1} \left( \frac{\alpha_2}{\alpha_1} \right)^{k+1} + \dots + \frac{c_n}{c_1} \left( \frac{\alpha_n}{\alpha_1} \right)^{k+1} \right)}{1 + \frac{c_2}{c_1} \left( \frac{\alpha_2}{\alpha_1} \right)^k + \dots + \frac{c_n}{c_1} \left( \frac{\alpha_n}{\alpha_1} \right)^k} = \frac{\alpha_1 (1 + 0 + \dots + 0)}{1 + 0 + \dots + 0} = \alpha_1 \end{aligned}$$

For all  $j = 2, \dots, n$ , since  $|\alpha_1| > |\alpha_j|$ , we have  $\lim_{k \rightarrow \infty} \left( \frac{\alpha_j}{\alpha_1} \right)^k = 0$ .

# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} = \alpha_1$$



# Bernoulli's Method

The linear difference equation

$$a_n u_k + a_{n-1} u_{k-1} + \dots + a_1 u_{k-n+1} + a_0 u_{k-n} = 0$$

has characteristic polynomial  $P(x)$ .

Then the limit of the ratio of consecutive terms of the sequence  $(u_i)$  is

$$\lim_{k \rightarrow \infty} \frac{u_{k+1}}{u_k} = \alpha_1$$

Therefore, if a randomly generated solution of the difference equation associated with the polynomial  $P(x)$  has non-zero  $c_1$  coefficient, for large values of  $k$ ,  $\frac{u_{k+1}}{u_k}$  is an approximation of the root of  $P(x)$  with largest absolute value.

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

$$u_3 = 0 \quad u_4 = -4 \quad u_5 = -17 \quad u_6 = -56 \quad \dots$$

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

$$u_3 = 0 \quad u_4 = -4 \quad u_5 = -17 \quad u_6 = -56 \quad \dots$$

$$u_{13} = -87892 \quad u_{14} = -246427 \quad u_{15} = -690703$$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = -4 \quad u_5 = -17 \quad u_6 = -56 \quad \dots$$

$$u_{13} = -87892 \quad u_{14} = -246427 \quad u_{15} = -690703$$

At every step we also check the ratio of the last two numbers

$$\frac{u_{14}}{u_{13}} \approx 2.803747781 \qquad \frac{u_{15}}{u_{14}} \approx 2.802870627$$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = -4 \quad u_5 = -17 \quad u_6 = -56 \quad \dots$$

$$u_{13} = -87892 \quad u_{14} = -246427 \quad u_{15} = -690703$$

At every step we also check the ratio of the last two numbers

$$\frac{u_{14}}{u_{13}} \approx 2.803747781 \qquad \frac{u_{15}}{u_{14}} \approx 2.802870627$$

These ratios are approximations to the root of largest absolute value.



## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of largest absolute value.

We form the linear difference equation  $u_k - 4u_{k-1} + 3u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = -4 \quad u_5 = -17 \quad u_6 = -56 \quad \dots$$

$$u_{13} = -87892 \quad u_{14} = -246427 \quad u_{15} = -690703$$

At every step we also check the ratio of the last two numbers

$$\frac{u_{14}}{u_{13}} \approx 2.803747781 \qquad \frac{u_{15}}{u_{14}} \approx 2.802870627$$

These ratios are approximations to the root of largest absolute value. In this example, the actual root of largest absolute value is

$$\alpha_1 \approx 2.801937736.$$

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1$

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1 = \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1$ .

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1 = \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1$ .

We have  $P(\frac{1}{x}) = 0 \Leftrightarrow \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1 = 0$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1 = \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1$ .

We have  $P(\frac{1}{x}) = 0 \Leftrightarrow \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1 = 0 \Leftrightarrow 1 - 4x + 3x^2 + x^3 = 0$

(Multiplying both sides by  $x^3$ )

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1 = \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1$ .

We have  $P(\frac{1}{x}) = 0 \Leftrightarrow \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1 = 0 \Leftrightarrow 1 - 4x + 3x^2 + x^3 = 0$

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ .

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

If we consider  $P(\frac{1}{x}) = (\frac{1}{x})^3 - 4(\frac{1}{x})^2 + 3(\frac{1}{x}) + 1 = \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1$ .

We have  $P(\frac{1}{x}) = 0 \Leftrightarrow \frac{1}{x^3} - \frac{4}{x^2} + \frac{3}{x} + 1 = 0 \Leftrightarrow 1 - 4x + 3x^2 + x^3 = 0$

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.



# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

# Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

$$u_3 = 0 \quad u_4 = 3 \quad u_5 = -10 \quad u_6 = 42 \quad \dots$$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

We randomly chose initial conditions  $u_0 = 1$ ,  $u_1 = 1$ ,  $u_2 = 1$  and use the difference equation to form terms of the sequence.

$$u_3 = 0 \quad u_4 = 3 \quad u_5 = -10 \quad u_6 = 42 \quad \dots$$

$$u_{13} = -745276 \quad u_{14} = 3017561 \quad u_{15} = -12217855$$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

Therefore, roots of  $Q(x) = x^3 + 3x^2 - 4x + 1$  are reciprocals of roots of  $P(x)$ . In particular, the root of  $Q(x)$  with largest absolute value is reciprocal of the root of  $P(x)$  with smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = 3 \quad u_5 = -10 \quad u_6 = 42 \quad \dots$$

$$u_{13} = -745276 \quad u_{14} = 3017561 \quad u_{15} = -12217855$$

The ratios  $\frac{u_k}{u_{k+1}} = \frac{1}{\frac{u_{k+1}}{u_k}}$  approximate the reciprocal of the root of largest absolute value for  $Q(x)$  which is the root of smallest absolute value for  $P(x)$ .

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = 3 \quad u_5 = -10 \quad u_6 = 42 \quad \dots$$

$$u_{13} = -745276 \quad u_{14} = 3017561 \quad u_{15} = -12217855$$

The ratios  $\frac{u_k}{u_{k+1}} = \frac{1}{\frac{u_{k+1}}{u_k}}$  approximate the reciprocal of the root of largest absolute value for  $Q(x)$  which is the root of smallest absolute value for  $P(x)$ .

$$\frac{u_{13}}{u_{14}} \approx -0.2469795971 \qquad \frac{u_{14}}{u_{15}} \approx -0.2469796048$$

## Example

For  $P(x) = x^3 - 4x^2 + 3x + 1$  approximate the root of smallest absolute value.

We consider linear difference equation  $u_k + 3u_{k-1} - 4u_{k-2} + u_{k-3} = 0$ .

$$u_3 = 0 \quad u_4 = 3 \quad u_5 = -10 \quad u_6 = 42 \quad \dots$$

$$u_{13} = -745276 \quad u_{14} = 3017561 \quad u_{15} = -12217855$$

The ratios  $\frac{u_k}{u_{k+1}} = \frac{1}{\frac{u_{k+1}}{u_k}}$  approximate the reciprocal of the root of largest absolute value for  $Q(x)$  which is the root of smallest absolute value for  $P(x)$ .

$$\frac{u_{13}}{u_{14}} \approx -0.2469795971 \quad \frac{u_{14}}{u_{15}} \approx -0.2469796048$$

In this example, the actual root of smallest absolute value is  $\alpha_3 \approx -0.2469796037$ .

It can be shown that even if  $P(x)$  has non-real complex solutions or multiple solutions the Bernoulli method converges to the root of largest absolute value as long as there is only one root of largest absolute value which is simple (multiplicity one) and real.



It can be shown that even if  $P(x)$  has non-real complex solutions or multiple solutions the Bernoulli method converges to the root of largest absolute value as long as there is only one root of largest absolute value which is simple (multiplicity one) and real.

When the root of largest absolute value is real but not simple the algorithm converges but very slowly.

# Final Comments

It can be shown that even if  $P(x)$  has non-real complex solutions or multiple solutions the Bernoulli method converges to the root of largest absolute value as long as there is only one root of largest absolute value which is simple (multiplicity one) and real.

When the root of largest absolute value is real but not simple the algorithm converges but very slowly.

When there are non-real roots of largest absolute value, the algorithm requires significant modifications and even with those modifications it will be preferred to use alternative methods.