

## (2.4) Error Analysis for Iterative Methods

MATH 4701 Numerical Analysis

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ .

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

A higher order of convergence and a smaller asymptotic error constant both mean a faster convergence.

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

A higher order of convergence and a smaller asymptotic error constant both mean a faster convergence. The impact of a higher order of convergence, however, is much more significant than the impact of smaller asymptotic error constant.

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

A higher order of convergence and a smaller asymptotic error constant both mean a faster convergence. The impact of a higher order of convergence, however, is much more significant than the impact of smaller asymptotic error constant.

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.

# Order of Convergence

Suppose  $(p_n)_{n \in \mathbb{N}}$  is a sequence with  $\lim_{n \rightarrow \infty} p_n = p$ . Moreover, assume  $p_n \neq p$  for all  $n \in \mathbb{N}$ .

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

A higher order of convergence and a smaller asymptotic error constant both mean a faster convergence. The impact of a higher order of convergence, however, is much more significant than the impact of smaller asymptotic error constant.

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.



# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

For example,  $p_n = \frac{1}{2^n}$  converges to zero linearly with  $\lambda = \frac{1}{2}$ .

$$\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \dots$$

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ .

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ . If we start with  $q_1 = \frac{1}{2}$ , we get  $q_2 = \frac{1}{2^3}$ ,

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ . If we start with  $q_1 = \frac{1}{2}$ , we get  $q_2 = \frac{1}{2^3}$ ,  $q_3 = \frac{1}{2^7}$ ,

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ . If we start with  $q_1 = \frac{1}{2}$ , we get  $q_2 = \frac{1}{2^3}$ ,  $q_3 = \frac{1}{2^7}$ ,  $q_4 = \frac{1}{2^{15}}$ ,

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ . If we start with  $q_1 = \frac{1}{2}$ , we get  $q_2 = \frac{1}{2^3}$ ,  $q_3 = \frac{1}{2^7}$ ,  $q_4 = \frac{1}{2^{15}}$ , and in general,  $p_n = \frac{1}{2^{2^n - 1}}$ .

# Order of Convergence

We say  $(p_n)_{n \in \mathbb{N}}$  converges to  $p$  of **order**  $\alpha$ , with **asymptotic error constant**  $\lambda$  if for **positive** constants  $\alpha$  and  $\lambda$  we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^\alpha} = \lambda$$

- If  $\alpha = 1$  and  $\lambda < 1$  the sequence is said to **converge linearly**.
- If  $\alpha = 2$  the sequence is said to **converge quadratically**.

To make an example of a sequence  $(q_n)$  that converges to 0 quadratically, with  $\lambda = \frac{1}{2}$ , we let  $q_{n+1} = \frac{1}{2}q_n^2$ . If we start with  $q_1 = \frac{1}{2}$ , we get  $q_2 = \frac{1}{2^3}$ ,  $q_3 = \frac{1}{2^7}$ ,  $q_4 = \frac{1}{2^{15}}$ , and in general,  $p_n = \frac{1}{2^{2^n - 1}}$ .

$$\frac{1}{2}, \frac{1}{8}, \frac{1}{128}, \frac{1}{32768}, \frac{1}{2147483648}, \dots$$

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ .



# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ .

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ . Then by the **fixed point iteration theorem**  $g$  has a unique fixed point  $p$  inside  $[a, b]$  and for any initial guess  $p_0 \in [a, b]$ , The sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$ .

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ . Then by the **fixed point iteration theorem**  $g$  has a unique fixed point  $p$  inside  $[a, b]$  and for any initial guess  $p_0 \in [a, b]$ , The sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$ .

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ . Then by the **fixed point iteration theorem**  $g$  has a unique fixed point  $p$  inside  $[a, b]$  and for any initial guess  $p_0 \in [a, b]$ , The sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$ .

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$g(p_n) - g(p) = g'(\zeta_n)(p_n - p)$$

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ . Then by the **fixed point iteration theorem**  $g$  has a unique fixed point  $p$  inside  $[a, b]$  and for any initial guess  $p_0 \in [a, b]$ , The sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$ .

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p)$$

# Linear Convergence of Fixed Point Iteration

Suppose  $g \in C[a, b]$ ,  $g([a, b]) \subset [a, b]$ . Moreover, suppose for some  $0 < K < 1$  and for all  $x \in [a, b]$ ,  $g'(x)$  exists and  $|g'(x)| \leq K$ . Then by the **fixed point iteration theorem**  $g$  has a unique fixed point  $p$  inside  $[a, b]$  and for any initial guess  $p_0 \in [a, b]$ , The sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$ .

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p) \Rightarrow \frac{p_{n+1} - p}{p_n - p} = g'(\zeta_n)$$

# Linear Convergence of Fixed Point Iteration

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p) \Rightarrow \frac{p_{n+1} - p}{p_n - p} = g'(\zeta_n)$$

By continuity of  $g'(x)$  on  $(a, b)$  and since  $\zeta_n$  between  $p_n$  and  $p$  and  $\lim_{n \rightarrow \infty} p_n = p$ , we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lim_{n \rightarrow \infty} |g'(\zeta_n)|$$

# Linear Convergence of Fixed Point Iteration

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p) \Rightarrow \frac{p_{n+1} - p}{p_n - p} = g'(\zeta_n)$$

By continuity of  $g'(x)$  on  $(a, b)$  and since  $\zeta_n$  between  $p_n$  and  $p$  and  $\lim_{n \rightarrow \infty} p_n = p$ , we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lim_{n \rightarrow \infty} |g'(\zeta_n)| = |g'(p)|$$



# Linear Convergence of Fixed Point Iteration

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p) \Rightarrow \frac{p_{n+1} - p}{p_n - p} = g'(\zeta_n)$$

By continuity of  $g'(x)$  on  $(a, b)$  and since  $\zeta_n$  between  $p_n$  and  $p$  and  $\lim_{n \rightarrow \infty} p_n = p$ , we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lim_{n \rightarrow \infty} |g'(\zeta_n)| = |g'(p)|$$

But  $\lambda = |g'(p)| \leq K < 1$  and  $\lambda = |g'(p)| > 0$ .

# Linear Convergence of Fixed Point Iteration

**Theorem:** If  $g'(p) \neq 0$  and  $g'(x)$  is continuous on  $(a, b)$ , then the sequence  $(p_n)$  of iterations converges only linearly to the fixed point  $p$ .

**Proof:** By the **mean value theorem**, for each  $n$ , there is a number  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} - p = g(p_n) - g(p) = g'(\zeta_n)(p_n - p) \Rightarrow \frac{p_{n+1} - p}{p_n - p} = g'(\zeta_n)$$

By continuity of  $g'(x)$  on  $(a, b)$  and since  $\zeta_n$  between  $p_n$  and  $p$  and  $\lim_{n \rightarrow \infty} p_n = p$ , we have

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|} = \lim_{n \rightarrow \infty} |g'(\zeta_n)| = |g'(p)|$$

But  $\lambda = |g'(p)| \leq K < 1$  and  $\lambda = |g'(p)| > 0$ . Therefore,  $(p_n)$  converges to  $p$  linearly.

# Taylor's Theorem

Although implementation of Newton's method only requires the existence of the first derivative of the function, its convergence depends of the function to have a continuous second derivative.

# Taylor's Theorem

Although implementation of Newton's method only requires the existence of the first derivative of the function, its convergence depends of the function to have a continuous second derivative.

To understand the conditions that guarantee the convergence of Newton's method, we need Taylor's theorem.

# Taylor's Theorem

Although implementation of Newton's method only requires the existence of the first derivative of the function, its convergence depends of the function to have a continuous second derivative.

To understand the conditions that guarantee the convergence of Newton's method, we need Taylor's theorem.

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$

# Taylor's Theorem

Although implementation of Newton's method only requires the existence of the first derivative of the function, its convergence depends of the function to have a continuous second derivative.

To understand the conditions that guarantee the convergence of Newton's method, we need Taylor's theorem.

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and

# Taylor's Theorem

Although implementation of Newton's method only requires the existence of the first derivative of the function, its convergence depends of the function to have a continuous second derivative.

To understand the conditions that guarantee the convergence of Newton's method, we need Taylor's theorem.

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and
- $x_0 \in [a, b]$

# Taylor's Theorem

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and
- $x_0 \in [a, b]$

Let  $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$



# Taylor's Theorem

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and
- $x_0 \in [a, b]$

Let  $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$

Then for every  $x \in [a, b]$ , there is a number  $\zeta = \zeta(x)$ , between  $x_0$  and  $x$ , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

# Taylor's Theorem

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and
- $x_0 \in [a, b]$

Let  $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$

Then for every  $x \in [a, b]$ , there is a number  $\zeta = \zeta(x)$ , between  $x_0$  and  $x$ , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

The polynomial  $P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)(x - x_0)^k}{k!}$  is called the **Taylor polynomial of degree  $n$  about  $x_0$** .

# Taylor's Theorem

**Taylor's Theorem:** Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$  (the  $n + 1$ th derivative of  $f$ ) exists on  $[a, b]$ , and
- $x_0 \in [a, b]$

Let  $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$

Then for every  $x \in [a, b]$ , there is a number  $\zeta = \zeta(x)$ , between  $x_0$  and  $x$ , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

The polynomial  $P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)(x - x_0)^k}{k!}$  is called the **Taylor polynomial of degree  $n$  about  $x_0$** .

The expression  $R_{n+1}(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n+1)!}$  is called the **remainder term** and represents the **error** for using  $P_n(x)$  to calculate the value of  $f(x)$ .

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}}$$



# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}} \Rightarrow f^{(3)}(1) = \frac{3}{8}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}} \Rightarrow f^{(3)}(1) = \frac{3}{8}$$

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}} \Rightarrow f^{(3)}(1) = \frac{3}{8}$$

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}} \Rightarrow f^{(4)}(1) = -\frac{15}{16}.$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}} \Rightarrow f^{(3)}(1) = \frac{3}{8}$$

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}} \Rightarrow f^{(4)}(1) = -\frac{15}{16}.$$

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2 + \frac{3}{8}(x-1)^3 - \frac{15}{16}(x-1)^4$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f(x) = x^{\frac{1}{2}} \Rightarrow f(1) = \sqrt{1} = 1$$

$$f'(x) = \left(\frac{1}{2}\right)x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \Rightarrow f'(1) = \frac{1}{2}$$

$$f''(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)x^{-\frac{3}{2}} \Rightarrow f''(1) = -\frac{1}{4}$$

$$f^{(3)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^{-\frac{5}{2}} \Rightarrow f^{(3)}(1) = \frac{3}{8}$$

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}} \Rightarrow f^{(4)}(1) = -\frac{15}{16}.$$

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2 + \frac{3}{8}(x-1)^3 - \frac{15}{16}(x-1)^4$$

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}} \Rightarrow f^{(4)}(1) = -\frac{15}{16}.$$
$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .



# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$f^{(4)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^{-\frac{7}{2}} \Rightarrow f^{(4)}(1) = -\frac{15}{16}.$$
$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$$f^{(5)}(x) = \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)\left(-\frac{7}{2}\right)x^{-\frac{9}{2}} = \left(\frac{105}{32}\right)x^{-\frac{9}{2}}.$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$R_5(0.5) = \frac{f^{(5)}(\zeta)(0.5-1)^5}{5!}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$R_5(0.5) = \frac{f^{(5)}(\zeta)(0.5-1)^5}{5!} = \frac{(\frac{105}{32})\zeta^{-\frac{9}{2}}(-0.5)^5}{5!}$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$R_5(0.5) = \frac{f^{(5)}(\zeta)(0.5-1)^5}{5!} = \frac{(\frac{105}{32})\zeta^{-\frac{9}{2}}(-0.5)^5}{5!} = -\frac{7}{8192\zeta^{\frac{9}{2}}}$$

## Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \leq R_5(0.5) = -\frac{7}{8192\zeta^{\frac{9}{2}}} \leq -\frac{7}{8192(1)^{\frac{9}{2}}}$$

$h(\zeta) = -\frac{7}{8192\zeta^{\frac{9}{2}}}$  is an **increasing** function.

## Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \leq R_5(0.5) = -\frac{7}{8192\zeta^{\frac{9}{2}}} \leq -\frac{7}{8192(1)^{\frac{9}{2}}}$$

Therefore, the error in calculating

$$f(0.5) = \sqrt{\frac{1}{2}} \approx 0.7071067810 \text{ using } P_4(0.5)$$

## Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \leq R_5(0.5) = -\frac{7}{8192\zeta^{\frac{9}{2}}} \leq -\frac{7}{8192(1)^{\frac{9}{2}}}$$

Therefore, the error in calculating

$f(0.5) = \sqrt{\frac{1}{2}} \approx 0.7071067810$  using

$$P_4(0.5) = 1 + \frac{1}{2}(0.5-1) - \frac{1}{2}(0.5-1)^2 + \frac{3}{8}(0.5-1)^3 - \frac{15}{24}(0.5-1)^4$$

## Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \leq R_5(0.5) = -\frac{7}{8192\zeta^{\frac{9}{2}}} \leq -\frac{7}{8192(1)^{\frac{9}{2}}}$$

Therefore, the error in calculating

$$f(0.5) = \sqrt{\frac{1}{2}} \approx 0.7071067810 \text{ using } P_4(0.5) \approx 0.7084960938$$



## Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

$f^{(5)}(x) = (\frac{105}{32})x^{-\frac{9}{2}}$ . Therefore, for  $0.5 \leq \zeta \leq 1$ , we have

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \leq R_5(0.5) = -\frac{7}{8192\zeta^{\frac{9}{2}}} \leq -\frac{7}{8192(1)^{\frac{9}{2}}}$$

Therefore, the error in calculating

$f(0.5) = \sqrt{\frac{1}{2}} \approx 0.7071067810$  using  $P_4(0.5) \approx 0.7084960938$

is between  $-\frac{7}{8192(1)^{\frac{9}{2}}} \approx -0.00085449$  and

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \approx -0.019335.$$

# Example

For  $f(x) = \sqrt{x}$ ,

- (1) Find the fourth degree Taylor polynomial  $P_4(x)$  for  $f(x)$  centered at (about)  $c = 1$ .

$$P_4(x) = 1 + \frac{1}{2}(x-1) - \frac{1}{8}(x-1)^2 + \frac{1}{16}(x-1)^3 + \frac{5}{128}(x-1)^4$$

- (2) Find the maximum and minimum possible error when we use  $P_4(x)$  to approximate  $f(0.5) = \sqrt{\frac{1}{2}}$ .

Therefore, the error in calculating

$$f(0.5) = \sqrt{\frac{1}{2}} \approx 0.7071067810 \text{ using } P_4(0.5) \approx 0.7084960938$$

is between  $-\frac{7}{8192(1)^{\frac{9}{2}}} \approx -0.00085449$  and

$$-\frac{7}{8192(0.5)^{\frac{9}{2}}} \approx -0.019335.$$

(In fact,  $f(0.5) - P_4(0.4) \approx -0.0013893128$ )

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume  $g'(p) = 0$ .**

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume  $g'(p) = 0$ .** Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume**  $g'(p) \neq 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

If  $x \in [p - \delta, p + \delta]$ , then  $|g(x) - p| = |g(x) - g(p)|$



# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

If  $x \in [p - \delta, p + \delta]$ , then  $|g(x) - p| = |g(x) - g(p)| \leq K|x - p|$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

If  $x \in [p - \delta, p + \delta]$ , then  $|g(x) - p| = |g(x) - g(p)| \leq K|x - p| \leq K\delta < \delta$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

If  $x \in [p - \delta, p + \delta]$ , then

$$|g(x) - p| = |g(x) - g(p)| \leq K|x - p| \leq K\delta < \delta \Rightarrow g(x) \in [p - \delta, p + \delta].$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

If  $x \in [p - \delta, p + \delta]$ , then

$$|g(x) - p| = |g(x) - g(p)| \leq K|x - p| \leq K\delta < \delta \Rightarrow g(x) \in [p - \delta, p + \delta].$$

Therefore, if  $p_0$  is in  $[p - \delta, p + \delta]$  all the values of  $p_n$  are in  $[p - \delta, p + \delta]$  as well.

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

By taking a possibly smaller value of  $\delta$  we can ensure that  $[p - \delta, p + \delta] \subset (a, b)$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$ .

Therefore, if  $p_0$  is in  $[p - \delta, p + \delta]$  all the values of  $p_n$  are in  $[p - \delta, p + \delta]$  as well.

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:** Since  $g'(x)$  is continuous on an open interval containing  $p$ , for any  $0 < K < 1$ , there is  $\delta > 0$  such that for all  $x \in [p - \delta, p + \delta]$ ,  $|g'(x)| \leq K$ .

Therefore, if  $p_0$  is in  $[p - \delta, p + \delta]$  all the values of  $p_n$  are in  $[p - \delta, p + \delta]$  as well.

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2 = p + \frac{g''(\zeta_x)}{2}(x - p)^2$$

Since  $g(p) = p$  and  $g'(p) = 0$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

Therefore, if  $p_0$  is in  $[p - \delta, p + \delta]$  all the values of  $p_n$  are in  $[p - \delta, p + \delta]$  as well.

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2 = p + \frac{g''(\zeta_x)}{2}(x - p)^2$$

In particular, since all the values of  $p_n$  are in  $[p - \delta, p + \delta]$ , there is  $\zeta_n$  between  $p_n$  and  $p$  such that

$$g(p_n) = p + \frac{g''(\zeta_n)}{2}(p_n - p)^2$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

Therefore, if  $p_0$  is in  $[p - \delta, p + \delta]$  all the values of  $p_n$  are in  $[p - \delta, p + \delta]$  as well.

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2 = p + \frac{g''(\zeta_x)}{2}(x - p)^2$$

In particular, since all the values of  $p_n$  are in  $[p - \delta, p + \delta]$ , there is  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} = g(p_n) = p + \frac{g''(\zeta_n)}{2}(p_n - p)^2$$



# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2 = p + \frac{g''(\zeta_x)}{2}(x - p)^2$$

In particular, since all the values of  $p_n$  are in  $[p - \delta, p + \delta]$ , there is  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} = g(p_n) = p + \frac{g''(\zeta_n)}{2}(p_n - p)^2 \Rightarrow \frac{p_{n+1} - p}{(p_n - p)^2} = \frac{g''(\zeta_n)}{2}$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

By **Taylor's Theorem** for all  $x \in [p - \delta, p + \delta]$ , there is  $\zeta_x$  between  $x$  and  $p$  such that

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\zeta_x)}{2}(x - p)^2 = p + \frac{g''(\zeta_x)}{2}(x - p)^2$$

In particular, since all the values of  $p_n$  are in  $[p - \delta, p + \delta]$ , there is  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} = g(p_n) = p + \frac{g''(\zeta_n)}{2}(p_n - p)^2 \Rightarrow \frac{p_{n+1} - p}{(p_n - p)^2} = \frac{g''(\zeta_n)}{2}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|g''(\zeta_n)|}{2}$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

In particular, since all the values of  $p_n$  are in  $[p - \delta, p + \delta]$ , there is  $\zeta_n$  between  $p_n$  and  $p$  such that

$$p_{n+1} = g(p_n) = p + \frac{g''(\zeta_n)}{2}(p_n - p)^2 \Rightarrow \frac{p_{n+1} - p}{(p_n - p)^2} = \frac{g''(\zeta_n)}{2}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|g''(\zeta_n)|}{2} = \frac{|g''(p)|}{2}$$

Since  $\zeta_n$  is between  $p_n$  and  $p$ ,  $\lim_{n \rightarrow \infty} p_n = p$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$  which contains  $p$  and all values of  $p_n$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume  $g'(p) = 0$ .** Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

**Proof:**

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|g''(\zeta_n)|}{2} = \frac{|g''(p)|}{2}$$

Since  $\zeta_n$  is between  $p_n$  and  $p$ ,  $\lim_{n \rightarrow \infty} p_n = p$  and  $g''(x)$  is continuous on  $[p - \delta, p + \delta]$  which contains  $p$  and all values of  $p_n$ .

This means  $(p_n)$  converges to  $p$  quadratically if  $g''(p) \neq 0$  and of higher convergence order if  $g''(p) = 0$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume  $g'(p) = 0$ .** Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$  and

$$g'(p) = 1 + \phi'(p)f(p) + \phi(p)f'(p)$$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$  and

$$g'(p) = 1 + \phi'(p)f(p) + \phi(p)f'(p) = 1 + \phi(p)f'(p)$$



# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$  and

$$g'(p) = 1 + \phi'(p)f(p) + \phi(p)f'(p) = 1 + \phi(p)f'(p)$$

Therefore,  $g'(p) = 0 \Leftrightarrow 1 + \phi(p)f'(p) = 0$

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$  and

$$g'(p) = 1 + \phi'(p)f(p) + \phi(p)f'(p) = 1 + \phi(p)f'(p)$$

Therefore,  $g'(p) = 0 \Leftrightarrow 1 + \phi(p)f'(p) = 0 \Leftrightarrow \phi(p) = \frac{-1}{f'(p)}$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

In attempting to create a fixed point method that helps us find root  $p$  of  $f(x) = 0$ , we can try to work with  $g(x) = x + \phi(x)f(x)$ , where  $\phi(x)f(x)$  as a functional multiple of  $f(x)$  will also have a root at  $p$  and adding  $x$  will change the root finding problem to a fixed point problem.

Then  $g'(x) = 1 + \phi'(x)f(x) + \phi(x)f'(x)$  and

$$g'(p) = 1 + \phi'(p)f(p) + \phi(p)f'(p) = 1 + \phi(p)f'(p)$$

Therefore,  $g'(p) = 0 \Leftrightarrow 1 + \phi(p)f'(p) = 0 \Leftrightarrow \phi(p) = \frac{-1}{f'(p)}$ .

In particular, if we take  $\phi(x) = \frac{-1}{f'(x)}$  we can accomplish  $g'(p) = 0$  for  $g(x) = x - \frac{f(x)}{f'(x)}$  which gives us Newton's method.

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ . **Additionally, assume  $g'(p) = 0$ .** Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

If we want to use the theorem above, we would need for  $f$  to be in  $C^3[a, b]$  (so that  $g \in C^2[a, b]$ ).

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .

**Additionally, assume**  $g'(p) \neq 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

If we want to use the theorem above, we would need for  $f$  to be in  $C^3[a, b]$  (so that  $g \in C^2[a, b]$ ). We can, however, show that Newton's method converges at least quadratically only using the conditions stated in the convergence of Newton's method theorem and only assuming  $f \in C^2[a, b]$ .

# Quadratic Convergence of Newton's Method

**Theorem:** Suppose  $g \in C^2[a, b]$  and  $g(p) = p$  for some  $p \in (a, b)$ .  
**Additionally, assume**  $g'(p) = 0$ . Then there is  $\delta > 0$  such that for all  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $(p_n)$  defined by  $p_{n+1} = g(p_n)$  converges to  $p$  **at least quadratically**.

If we want to use the theorem above, we would need for  $f$  to be in  $C^3[a, b]$  (so that  $g \in C^2[a, b]$ ). We can, however, show that Newton's method converges at least quadratically only using the conditions stated in the convergence of Newton's method theorem and only assuming  $f \in C^2[a, b]$ .

If  $f \in C^1[a, b]$  and  $f''(x)$  exists, by **Taylor's theorem**,

$$f(x) = f(p) + f'(p)(x - p) + f''(\zeta) \frac{(x - p)^2}{2}$$

for some  $\delta > 0$  and all  $x$  in  $[p - \delta, p + \delta]$  and some  $\zeta$  between  $x$  and  $p$ .

# Quadratic Convergence of Newton's Method

We can, however, show that Newton's method converges at least quadratically only using the conditions stated in the convergence of Newton's method theorem and only assuming  $f \in C^2[a, b]$ .

If  $f \in C^1[a, b]$  and  $f''(x)$  exists, by **Taylor's theorem**,

$$f(x) = f(p) + f'(p)(x - p) + f''(\zeta) \frac{(x - p)^2}{2}$$

for some  $\delta > 0$  and all  $x$  in  $[p - \delta, p + \delta]$  and some  $\zeta$  between  $x$  and  $p$ .

We have already seen for a choice of  $\delta > 0$  all points  $p_n$  are in  $[p - \delta, p + \delta]$  and letting the center  $c = p_n$  and  $x = p$ , we get

$$f(p) = f(p_n) + f'(p_n)(p - p_n) + f''(\zeta_n) \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

# Quadratic Convergence of Newton's Method

We have already seen for a choice of  $\delta > 0$  all points  $p_n$  are in  $[p - \delta, p + \delta]$  and letting the center  $c = p_n$  and  $x = p$ , we get

$$f(p) = f(p_n) + f'(p_n)(p - p_n) + f''(\zeta_n) \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

Since  $f(p) = 0$  we have

$$0 = f(p_n) + f'(p_n)(p - p_n) + f''(\zeta_n) \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .



# Quadratic Convergence of Newton's Method

Since  $f(p) = 0$  we have

$$0 = f(p_n) + f'(p_n)(p - p_n) + f''(\zeta_n) \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

Dividing both sides by  $f'(p_n) \neq 0$  we have

$$0 = \frac{f(p_n)}{f'(p_n)} + p - p_n + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

# Quadratic Convergence of Newton's Method

Dividing both sides by  $f'(p_n) \neq 0$  we have

$$0 = \frac{f(p_n)}{f'(p_n)} + p - p_n + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

Since  $p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)}$  we get

$$p_n - \frac{f(p_n)}{f'(p_n)} = p + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

# Quadratic Convergence of Newton's Method

Dividing both sides by  $f'(p_n) \neq 0$  we have

$$0 = \frac{f(p_n)}{f'(p_n)} + p - p_n + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

Since  $p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)}$  we get

$$p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)} = p + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

# Quadratic Convergence of Newton's Method

Since  $p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)}$  we get

$$p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)} = p + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

$$\Rightarrow p_{n+1} - p = \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

# Quadratic Convergence of Newton's Method

Since  $p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)}$  we get

$$p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)} = p + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

$$\Rightarrow p_{n+1} - p = \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

$$\Rightarrow \frac{p_{n+1} - p}{(p - p_n)^2} = \frac{f''(\zeta_n)}{2f'(p_n)}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

# Quadratic Convergence of Newton's Method

Since  $p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)}$  we get

$$p_{n+1} = p_n - \frac{f(p_n)}{f'(p_n)} = p + \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

$$\Rightarrow p_{n+1} - p = \frac{f''(\zeta_n)}{f'(p_n)} \frac{(p - p_n)^2}{2}$$

$$\Rightarrow \frac{p_{n+1} - p}{(p - p_n)^2} = \frac{f''(\zeta_n)}{2f'(p_n)}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p - p_n|^2} = \lim_{n \rightarrow \infty} \frac{|f''(\zeta_n)|}{2|f'(p_n)|}$$

# Quadratic Convergence of Newton's Method

$$\Rightarrow \frac{p_{n+1} - p}{(p - p_n)^2} = \frac{f''(\zeta_n)}{2f'(p_n)}$$

for some  $\zeta_n$  between  $p_n$  and  $p$ .

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p - p_n|^2} = \lim_{n \rightarrow \infty} \frac{|f''(\zeta_n)|}{2|f'(p_n)|} = \frac{|f''(p)|}{2|f'(p)|}$$

If  $f''(x)$  is continuous on  $[p - \delta, p + \delta]$ , since  $\zeta_n$  between  $p_n$  and  $p$  and  $p_n$  are in  $[p - \delta, p + \delta]$ .

# Quadratic Convergence of Newton's Method

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p - p_n|^2} = \lim_{n \rightarrow \infty} \frac{|f''(\zeta_n)|}{2|f'(p_n)|} = \frac{|f''(p)|}{2|f'(p)|}$$

If  $f''(x)$  is continuous on  $[p - \delta, p + \delta]$ , since  $\zeta_n$  between  $p_n$  and  $p$  and  $p$  and  $p_n$  are in  $[p - \delta, p + \delta]$ .

This means Newton's method converges quadratically if  $f''(p) \neq 0$  and it converges faster than quadratically if  $f''(p) = 0$ .



# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  
 $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  
 $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .  
 $p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:** If  $p$  is a simple zero of  $f(x)$  then  $f(x) = (x - p)q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:** If  $p$  is a simple zero of  $f(x)$  then  $f(x) = (x - p)q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ .

Using product rule we get  $f'(x) = q(x) + (x - p)q'(x)$  for all  $x \neq p$  and since  $f \in C^1[a, b]$ ,

$$f'(p) = \lim_{x \rightarrow p} f'(x) = q(p) + (p - p)q'(p) = q(p) \neq 0.$$

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:** If  $p$  is a simple zero of  $f(x)$  then  $f(x) = (x - p)q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ .

Using product rule we get  $f'(x) = q(x) + (x - p)q'(x)$  for all  $x \neq p$  and since  $f \in C^1[a, b]$ ,

$f'(p) = \lim_{x \rightarrow p} f'(x) = q(p) + (p - p)q'(p) = q(p) \neq 0$ . Therefore, a simple zero  $p$  satisfies  $f(p) = 0$  and  $f'(p) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:**

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity  $m$**  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

## Proof:

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ . Then for all  $x \in [a, b]$ ,  $f(x) = f(p) + f'(\zeta(x))(x - p)$  where  $\zeta(x)$  is between  $p$  and  $x$ .



# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

## Proof:

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ . Then for all  $x \in [a, b]$ ,  $f(x) = f(p) + f'(\zeta(x))(x - p)$  where  $\zeta(x)$  is between  $p$  and  $x$ . Letting  $q(x) = f'(\zeta(x))$ , using  $f(p) = 0$  we get  $f(x) = q(x)(x - p)$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if

$f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:**

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ . Then for all  $x \in [a, b]$ ,  
 $f(x) = f(p) + f'(\zeta(x))(x - p)$  where  $\zeta(x)$  is between  $p$  and  $x$ . Letting  
 $q(x) = f'(\zeta(x))$ , using  $f(p) = 0$  we get  $f(x) = q(x)(x - p)$ . Moreover,  
 $\lim_{x \rightarrow p} q(x) = \lim_{x \rightarrow p} f'(\zeta(x))$

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if

$f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

**Proof:**

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ . Then for all  $x \in [a, b]$ ,  
 $f(x) = f(p) + f'(\zeta(x))(x - p)$  where  $\zeta(x)$  is between  $p$  and  $x$ . Letting  
 $q(x) = f'(\zeta(x))$ , using  $f(p) = 0$  we get  $f(x) = q(x)(x - p)$ . Moreover,  
 $\lim_{x \rightarrow p} q(x) = \lim_{x \rightarrow p} f'(\zeta(x)) = f'(p) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

## Proof:

Now assume  $f(p) = 0$  and  $f'(p) \neq 0$ . Then for all  $x \in [a, b]$ ,  $f(x) = f(p) + f'(\zeta(x))(x - p)$  where  $\zeta(x)$  is between  $p$  and  $x$ . Letting  $q(x) = f'(\zeta(x))$ , using  $f(p) = 0$  we get  $f(x) = q(x)(x - p)$ . Moreover,  $\lim_{x \rightarrow p} q(x) = \lim_{x \rightarrow p} f'(\zeta(x)) = f'(p) \neq 0$ . Therefore,  $p$  is a simple zero of  $f(x)$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

$p$  is called a **simple zero** of  $f(x)$  if  $p$  is a zero of **multiplicity one**.

**Theorem:** For  $f \in C^1[a, b]$ ,  $p \in (a, b)$  is a simple zero of  $f(x)$  if and only if  $f(p) = 0$  and  $f'(p) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0$ ,  $f'(p) = 0, \dots$ ,  $f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0$ ,  $f'(p) = 0, \dots$ ,  $f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

**Example:** For  $f(x) = e^x - 1 - x - \frac{x^2}{2}$  we have  
 $f(0) = 1 - 1 - 0 - 0 = 0$ ,

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if  $f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0, f'(p) = 0, \dots, f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

**Example:** For  $f(x) = e^x - 1 - x - \frac{x^2}{2}$  we have

$$f(0) = 1 - 1 - 0 - 0 = 0,$$

$$f'(x) = e^x - 1 - x \Rightarrow f'(0) = 1 - 1 - 0 = 0,$$



# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if

$f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0$ ,  $f'(p) = 0, \dots$ ,  $f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

**Example:** For  $f(x) = e^x - 1 - x - \frac{x^2}{2}$  we have

$$f(0) = 1 - 1 - 0 - 0 = 0,$$

$$f'(x) = e^x - 1 - x \Rightarrow f'(0) = 1 - 1 - 0 = 0,$$

$$f''(x) = e^x - 1 \Rightarrow f''(0) = 1 - 1 = 0,$$

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if

$f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0$ ,  $f'(p) = 0, \dots$ ,  $f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

**Example:** For  $f(x) = e^x - 1 - x - \frac{x^2}{2}$  we have

$$f(0) = 1 - 1 - 0 - 0 = 0,$$

$$f'(x) = e^x - 1 - x \Rightarrow f'(0) = 1 - 1 - 0 = 0,$$

$$f''(x) = e^x - 1 \Rightarrow f''(0) = 1 - 1 = 0,$$

$$\text{and } f^{(3)}(x) = e^x \Rightarrow f^{(3)}(0) = 1 \neq 0.$$

# Multiple Roots

A solution  $p$  of  $f(x) = 0$  is a **zero of multiplicity**  $m$  if

$f(x) = (x - p)^m q(x)$  for a function  $q(x)$  satisfying  $\lim_{x \rightarrow p} q(x) \neq 0$ .

This more general version of the above theorem can be proven in a similar way.

**Theorem:** For  $f \in C^m[a, b]$ ,  $p \in (a, b)$  is a zero of multiplicity  $m$  of  $f(x)$  if and only if  $f(p) = 0$ ,  $f'(p) = 0, \dots$ ,  $f^{(m-1)}(p) = 0$ , but  $f^{(m)}(p) \neq 0$ .

**Example:** For  $f(x) = e^x - 1 - x - \frac{x^2}{2}$  we have

$$f(0) = 1 - 1 - 0 - 0 = 0,$$

$$f'(x) = e^x - 1 - x \Rightarrow f'(0) = 1 - 1 - 0 = 0,$$

$$f''(x) = e^x - 1 \Rightarrow f''(0) = 1 - 1 = 0,$$

$$\text{and } f^{(3)}(x) = e^x \Rightarrow f^{(3)}(0) = 1 \neq 0.$$

Therefore,  $p = 0$  is a zero of **multiplicity three** for  $f(x)$ .

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$



# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)}$$

# Modified Newton's Method

When  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , our proof of convergence of Newton's method and its quadratic rate of convergence break down.

As a result, instead we try to find zeros of a function for which all the zeros of  $f(x)$  appear as simple zeros.

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)}$$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)} = 0$$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)} = 0$$

Also if we let  $Q(x) = \frac{q(x)}{m q(x) + (x - p) q'(x)}$  then  $h(x) = (x - p) Q(x)$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)} = 0$$

Also if we let  $Q(x) = \frac{q(x)}{m q(x) + (x - p) q'(x)}$  then  $h(x) = (x - p) Q(x)$  and  $\lim_{x \rightarrow p} Q(x) = \lim_{x \rightarrow p} \frac{q(x)}{m q(x) + (x - p) q'(x)}$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)} = 0$$

Also if we let  $Q(x) = \frac{q(x)}{m q(x) + (x - p) q'(x)}$  then  $h(x) = (x - p) Q(x)$  and  $\lim_{x \rightarrow p} Q(x) = \lim_{x \rightarrow p} \frac{q(x)}{m q(x) + (x - p) q'(x)} = \frac{q(p)}{m q(p) + (p - p) q'(p)}$

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

To see this notice that if  $f(x) = (x - p)^m q(x)$  such that  $\lim_{x \rightarrow p} q(x) \neq 0$ , then  $f'(x) = m(x - p)^{m-1} q(x) + (x - p)^m q'(x)$  and

$$h(x) = \frac{(x - p)^m q(x)}{m(x - p)^{m-1} q(x) + (x - p)^m q'(x)} = \frac{(x - p) q(x)}{m q(x) + (x - p) q'(x)}$$

$$h(p) = \frac{(p - p) q(p)}{m q(p) + (p - p) q'(p)} = 0$$

Also if we let  $Q(x) = \frac{q(x)}{m q(x) + (x - p) q'(x)}$  then  $h(x) = (x - p) Q(x)$  and  $\lim_{x \rightarrow p} Q(x) = \frac{q(p)}{m q(p) + (p - p) q'(p)} = \frac{q(p)}{m q(p)} = \frac{1}{m} \neq 0$ .



# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

Therefore, we can apply Newton's method to  $h(x)$  to find its simple root  $p$  instead of applying it to  $f(x)$  for which the conditions of theorems we proved are not satisfied.

# Modified Newton's Method

If  $p$  is a zero of  $f(x)$  with multiplicity  $m \geq 2$ , then  $p$  is a zero of  $f'(x)$  with multiplicity  $m - 1 \geq 1$ .

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

Therefore, we can apply Newton's method to  $h(x)$  to find its simple root  $p$  instead of applying it to  $f(x)$  for which the conditions of theorems we proved are not satisfied.

calculating  $h'(x) = \frac{[f'(x)]^2 - f(x)f''(x)}{[f'(x)]^2}$  we have

$$g(x) = x - \frac{h(x)}{h'(x)}$$

# Modified Newton's Method

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

Therefore, we can apply Newton's method to  $h(x)$  to find its simple root  $p$  instead of applying it to  $f(x)$  for which the conditions of theorems we proved are not satisfied.

calculating  $h'(x) = \frac{[f'(x)]^2 - f(x)f''(x)}{[f'(x)]^2}$  we have

$$g(x) = x - \frac{h(x)}{h'(x)} = x - \frac{\frac{f(x)}{f'(x)}}{\frac{[f'(x)]^2 - f(x)f''(x)}{[f'(x)]^2}}$$

Therefore, in modified Newton's method, instead of iterations of  $g(x) = x - \frac{f(x)}{f'(x)}$ , we consider iterations of  $g(x) = x - \frac{f(x)f'(x)}{[f'(x)]^2 - f(x)f''(x)}$ .

# Modified Newton's Method

If we define  $h(x) = \frac{f(x)}{f'(x)}$ , then  $p$  is a simple zero of  $h(x)$ .

Therefore, we can apply Newton's method to  $h(x)$  to find its simple root  $p$  instead of applying it to  $f(x)$  for which the conditions of theorems we proved are not satisfied.

calculating  $h'(x) = \frac{[f'(x)]^2 - f(x)f''(x)}{[f'(x)]^2}$  we have

$$g(x) = x - \frac{h(x)}{h'(x)} = x - \frac{\frac{f(x)}{f'(x)}}{\frac{[f'(x)]^2 - f(x)f''(x)}{[f'(x)]^2}} = x - \frac{f(x)f'(x)}{[f'(x)]^2 - f(x)f''(x)}$$

Therefore, in modified Newton's method, instead of iterations of  $g(x) = x - \frac{f(x)}{f'(x)}$ , we consider iterations of  $g(x) = x - \frac{f(x)f'(x)}{[f'(x)]^2 - f(x)f''(x)}$ .