

Gräffe's Root Squaring Method

MATH 4701 Numerical Analysis

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

Since

$$-P(x)P(-x) = -(x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(-x - \alpha_1)(-x - \alpha_2)(-x - \alpha_3)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example

$$-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1 \text{ and } P_1(w) = w^3 - 10w^2 + 17w - 1.$$

Since

$$\begin{aligned} -P(x)P(-x) &= -(x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(-x - \alpha_1)(-x - \alpha_2)(-x - \alpha_3) \\ &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(x + \alpha_1)(x + \alpha_2)(x + \alpha_3) \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned}P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\&= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)\end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

Since

$$\begin{aligned}-P(x)P(-x) &= -(x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(-x - \alpha_1)(-x - \alpha_2)(-x - \alpha_3) \\&= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(x + \alpha_1)(x + \alpha_2)(x + \alpha_3) \\&= (x^2 - \alpha_1^2)(x^2 - \alpha_2^2)(x^2 - \alpha_3^2)\end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned}P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\&= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)\end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

Since

$$\begin{aligned}-P(x)P(-x) &= -(x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(-x - \alpha_1)(-x - \alpha_2)(-x - \alpha_3) \\&= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)(x + \alpha_1)(x + \alpha_2)(x + \alpha_3) \\&= (x^2 - \alpha_1^2)(x^2 - \alpha_2^2)(x^2 - \alpha_3^2)\end{aligned}$$

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Roots of $P_1(w)$ are squares of roots of $P(x)$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Roots of $P_1(w)$ are squares of roots of $P(x)$. In particular,

$$P_1(x) = x^3 - (\alpha_1^2 + \alpha_2^2 + \alpha_3^2)x^2 + (\alpha_1^2\alpha_2^2 + \alpha_1^2\alpha_3^2 + \alpha_2^2\alpha_3^2)x - (\alpha_1^2\alpha_2^2\alpha_3^2)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_1(w) = -P(x)P(-x)$ where $w = x^2$. For this example $-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1$ and $P_1(w) = w^3 - 10w^2 + 17w - 1$.

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Roots of $P_1(w)$ are squares of roots of $P(x)$. In particular,

$$P_1(x) = x^3 - (\alpha_1^2 + \alpha_2^2 + \alpha_3^2)x^2 + (\alpha_1^2\alpha_2^2 + \alpha_1^2\alpha_3^2 + \alpha_2^2\alpha_3^2)x - (\alpha_1^2\alpha_2^2\alpha_3^2)$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

For this example

$$-P(x)P(-x) = x^6 - 10x^4 + 17x^2 - 1 \text{ and } P_1(w) = w^3 - 10w^2 + 17w - 1.$$

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Roots of $P_1(w)$ are squares of roots of $P(x)$. In particular,

$$P_1(x) = x^3 - (\alpha_1^2 + \alpha_2^2 + \alpha_3^2)x^2 + (\alpha_1^2\alpha_2^2 + \alpha_1^2\alpha_3^2 + \alpha_2^2\alpha_3^2)x - (\alpha_1^2\alpha_2^2\alpha_3^2)$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$-P_1(x)P_1(-x) = x^6 - 66x^4 + 269x^2 - 1$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

$$\text{and } P_1(w) = w^3 - 10w^2 + 17w - 1.$$

$$\Rightarrow P_1(w) = (w - \alpha_1^2)(w - \alpha_2^2)(w - \alpha_3^2)$$

Roots of $P_1(w)$ are squares of roots of $P(x)$. In particular,

$$P_1(x) = x^3 - (\alpha_1^2 + \alpha_2^2 + \alpha_3^2)x^2 + (\alpha_1^2\alpha_2^2 + \alpha_1^2\alpha_3^2 + \alpha_2^2\alpha_3^2)x - (\alpha_1^2\alpha_2^2\alpha_3^2)$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$-P_1(x)P_1(-x) = x^6 - 66x^4 + 269x^2 - 1 \text{ and}$$

$$P_2(w) = w^3 - 66w^2 + 269w - 1.$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$-P_1(x)P_1(-x) = x^6 - 66x^4 + 269x^2 - 1 \text{ and}$$

$$P_2(w) = w^3 - 66w^2 + 269w - 1.$$

Since

$$-P_1(x)P_1(-x) = -(x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(-x - \alpha_1^2)(-x - \alpha_2^2)(-x - \alpha_3^2)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$\begin{aligned} -P_1(x)P_1(-x) &= x^6 - 66x^4 + 269x^2 - 1 \text{ and} \\ P_2(w) &= w^3 - 66w^2 + 269w - 1. \end{aligned}$$

Since

$$\begin{aligned} -P_1(x)P_1(-x) &= -(x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(-x - \alpha_1^2)(-x - \alpha_2^2)(-x - \alpha_3^2) \\ &= (x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(x + \alpha_1^2)(x + \alpha_2^2)(x + \alpha_3^2) \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$\begin{aligned} -P_1(x)P_1(-x) &= x^6 - 66x^4 + 269x^2 - 1 \text{ and} \\ P_2(w) &= w^3 - 66w^2 + 269w - 1. \end{aligned}$$

Since

$$\begin{aligned} -P_1(x)P_1(-x) &= -(x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(-x - \alpha_1^2)(-x - \alpha_2^2)(-x - \alpha_3^2) \\ &= (x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(x + \alpha_1^2)(x + \alpha_2^2)(x + \alpha_3^2) \\ &= (x^2 - \alpha_1^4)(x^2 - \alpha_2^4)(x^2 - \alpha_3^4) \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$-P_1(x)P_1(-x) = x^6 - 66x^4 + 269x^2 - 1 \text{ and}$$

$$P_2(w) = w^3 - 66w^2 + 269w - 1.$$

Since

$$\begin{aligned} -P_1(x)P_1(-x) &= -(x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(-x - \alpha_1^2)(-x - \alpha_2^2)(-x - \alpha_3^2) \\ &= (x - \alpha_1^2)(x - \alpha_2^2)(x - \alpha_3^2)(x + \alpha_1^2)(x + \alpha_2^2)(x + \alpha_3^2) \\ &= (x^2 - \alpha_1^4)(x^2 - \alpha_2^4)(x^2 - \alpha_3^4) \end{aligned}$$

$$\Rightarrow P_2(w) = (w - \alpha_1^4)(w - \alpha_2^4)(w - \alpha_3^4)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example

$$\begin{aligned} -P_1(x)P_1(-x) &= x^6 - 66x^4 + 269x^2 - 1 \text{ and} \\ P_2(w) &= w^3 - 66w^2 + 269w - 1. \end{aligned}$$

$$\Rightarrow P_2(w) = (w - \alpha_1^4)(w - \alpha_2^4)(w - \alpha_3^4)$$

Roots of $P_2(w)$ are roots of $P(x)$ to the power four.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Now we form $P_2(w) = -P_1(x)P_1(-x)$ where $w = x^2$. For this example $-P_1(x)P_1(-x) = x^6 - 66x^4 + 269x^2 - 1$ and $P_2(w) = w^3 - 66w^2 + 269w - 1$.

$$\Rightarrow P_2(w) = (w - \alpha_1^4)(w - \alpha_2^4)(w - \alpha_3^4)$$

Roots of $P_2(w)$ are roots of $P(x)$ to the power four. In particular,

$$P_2(x) = x^3 - (\alpha_1^4 + \alpha_2^4 + \alpha_3^4)x^2 + (\alpha_1^4\alpha_2^4 + \alpha_1^4\alpha_3^4 + \alpha_2^4\alpha_3^4)x - (\alpha_1^4\alpha_2^4\alpha_3^4)$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

$$\Rightarrow P_2(w) = (w - \alpha_1^4)(w - \alpha_2^4)(w - \alpha_3^4)$$

Roots of $P_2(w)$ are roots of $P(x)$ to the power four. In particular,

$$P_2(x) = x^3 - (\alpha_1^4 + \alpha_2^4 + \alpha_3^4)x^2 + (\alpha_1^4\alpha_2^4 + \alpha_1^4\alpha_3^4 + \alpha_2^4\alpha_3^4)x - (\alpha_1^4\alpha_2^4\alpha_3^4)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

$$\Rightarrow P_2(w) = (w - \alpha_1^4)(w - \alpha_2^4)(w - \alpha_3^4)$$

Roots of $P_2(w)$ are roots of $P(x)$ to the power four. In particular,

$$P_2(x) = x^3 - (\alpha_1^4 + \alpha_2^4 + \alpha_3^4)x^2 + (\alpha_1^4\alpha_2^4 + \alpha_1^4\alpha_3^4 + \alpha_2^4\alpha_3^4)x - (\alpha_1^4\alpha_2^4\alpha_3^4)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

For all $n \geq 0$ we have

$$P_n(w) = (w - \alpha_1^{2^n})(w - \alpha_2^{2^n})(w - \alpha_3^{2^n})$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Roots of $P_2(w)$ are roots of $P(x)$ to the power four. In particular,

$$P_2(x) = x^3 - (\alpha_1^4 + \alpha_2^4 + \alpha_3^4)x^2 + (\alpha_1^4\alpha_2^4 + \alpha_1^4\alpha_3^4 + \alpha_2^4\alpha_3^4)x - (\alpha_1^4\alpha_2^4\alpha_3^4)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

For all $n \geq 0$ we have

$$P_n(w) = (w - \alpha_1^{2^n})(w - \alpha_2^{2^n})(w - \alpha_3^{2^n})$$

Roots of $P_n(w)$ are roots of $P(x)$ to the power 2^n .

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

For all $n \geq 0$ we have

$$P_n(w) = (w - \alpha_1^{2^n})(w - \alpha_2^{2^n})(w - \alpha_3^{2^n})$$

Roots of $P_n(w)$ are roots of $P(x)$ to the power 2^n . In particular,

$$\begin{aligned} P_n(x) &= \\ x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}) \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

Roots of $P_n(w)$ are roots of $P(x)$ to the power 2^n . In particular,

$$\begin{aligned} P_n(x) &= \\ x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 &+ (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}) \end{aligned}$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then

$$\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = \lim_{n \rightarrow \infty} \sqrt[2^n]{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|} =$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then

$$\begin{aligned}\lim_{n \rightarrow \infty} \sqrt[n]{|\sigma_n|} &= \lim_{n \rightarrow \infty} \sqrt[n]{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|} = \\ &= \lim_{n \rightarrow \infty} \sqrt[n]{\left| \alpha_1^{2^n} \left(1 + \left(\frac{\alpha_2}{\alpha_1} \right)^{2^n} + \left(\frac{\alpha_3}{\alpha_1} \right)^{2^n} \right) \right|} =\end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$\begin{aligned} P_n(x) &= \\ x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}) \end{aligned}$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|} = \\ &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\left| \alpha_1^{2^n} \left(1 + \left(\frac{\alpha_2}{\alpha_1} \right)^{2^n} + \left(\frac{\alpha_3}{\alpha_1} \right)^{2^n} \right) \right|} = \\ &= \lim_{n \rightarrow \infty} |\alpha_1| \sqrt[2^n]{\left| 1 + \left(\frac{\alpha_2}{\alpha_1} \right)^{2^n} + \left(\frac{\alpha_3}{\alpha_1} \right)^{2^n} \right|} \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|} = \\ &= \lim_{n \rightarrow \infty} |\alpha_1| \sqrt[2^n]{\left|1 + \left(\frac{\alpha_2}{\alpha_1}\right)^{2^n} + \left(\frac{\alpha_3}{\alpha_1}\right)^{2^n}\right|} = |\alpha_1| \sqrt[2^n]{|1 + 0 + 0|} \end{aligned}$$

Since $|\alpha_1| > |\alpha_2| \geq |\alpha_3|$, $\lim_{n \rightarrow \infty} \left|\frac{\alpha_2}{\alpha_1}\right|^{2^n} = 0$ and $\lim_{n \rightarrow \infty} \left|\frac{\alpha_3}{\alpha_1}\right|^{2^n} = 0$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then

$$\begin{aligned}\lim_{n \rightarrow \infty} \sqrt[n]{|\sigma_n|} &= \lim_{n \rightarrow \infty} \sqrt[n]{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|} = \\ &= \lim_{n \rightarrow \infty} |\alpha_1| \sqrt[n]{\left|1 + \left(\frac{\alpha_2}{\alpha_1}\right)^{2^n} + \left(\frac{\alpha_3}{\alpha_1}\right)^{2^n}\right|} = |\alpha_1| \sqrt[n]{|1 + 0 + 0|} = |\alpha_1|\end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$P(x) = (x - \alpha_1)(x - \alpha_2)(x - \alpha_3)$$

$$= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3)$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and $\sqrt[16]{14432666} \approx 2.801942122$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$. In particular,

$$\begin{aligned} P(x) &= (x - \alpha_1)(x - \alpha_2)(x - \alpha_3) \\ &= x^3 - (\alpha_1 + \alpha_2 + \alpha_3)x^2 + (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)x - (\alpha_1\alpha_2\alpha_3) \end{aligned}$$

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$\begin{aligned} P_n(x) &= \\ x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 &+ (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}) \end{aligned}$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and $\sqrt[16]{14432666} \approx 2.801942122$. The actual root α_1 of largest absolute value is $\alpha_1 \approx 2.801937736$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then

$$\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|}} =$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\tau_n|}{|\sigma_n|}} &= \lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}{|\alpha_1^{2^n}|(|1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}{|\alpha_1^{2^n}|(|1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = \\ &= \lim_{n \rightarrow \infty} |\alpha_2| \sqrt[2^n]{\left| \frac{1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}}{1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}} \right|}} \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}{|\alpha_1^{2^n}|(|1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = \\ &= \lim_{n \rightarrow \infty} |\alpha_2| \sqrt[2^n]{\left| \frac{1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}}{1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}} \right|} = |\alpha_2| \sqrt[2^n]{\left| \frac{1+0+0}{1+0+0} \right|} \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}{|\alpha_1^{2^n}|(|1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = \\ &= \lim_{n \rightarrow \infty} |\alpha_2| \sqrt[2^n]{\left| \frac{1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}}{1 + (\frac{\alpha_2}{\alpha_1})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}} \right|} = |\alpha_2| \sqrt[2^n]{\left| \frac{1+0+0}{1+0+0} \right|} = |\alpha_2| \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and

$$\sqrt[16]{\frac{5217020805}{14432666}} \approx 1.445039605.$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and $\sqrt[16]{\frac{5217020805}{14432666}} \approx 1.445039605$. The actual root α_2 of middle absolute value is $\alpha_2 \approx 1.445041868$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

$$\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\rho_n|}{|\tau_n|}} = \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}} =$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

$$\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\rho_n|}{|\tau_n|}} = \lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}} =$$

$$\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1, α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\rho_n|}{|\tau_n|}} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}} = \\ \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n}|(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} &= \lim_{n \rightarrow \infty} |\alpha_3| \sqrt[2^n]{\frac{1}{(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\left|\frac{\tau_n}{\sigma_n}\right|} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\left|\frac{\rho_n}{\tau_n}\right|} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} |\alpha_3| \sqrt[2^n]{\frac{1}{(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = |\alpha_3| \sqrt[2^n]{\frac{1}{(1+0+0)}} \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\rho_n|}{|\tau_n|}} &= \lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n}|}{|\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n}|}} = \\ &= \lim_{n \rightarrow \infty} |\alpha_3| \sqrt[2^n]{\frac{1}{(|1 + (\frac{\alpha_3}{\alpha_2})^{2^n} + (\frac{\alpha_3}{\alpha_1})^{2^n}|)}} = |\alpha_3| \sqrt[2^n]{\frac{1}{(1+0+0)}} = |\alpha_3| \end{aligned}$$

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\rho_n|}{|\tau_n|}} = |\alpha_3|$

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and $\sqrt[16]{\frac{1}{5217020805}} \approx 0.2469796037$.

Example

Let $P(x) = x^3 - 4x^2 + 3x + 1$. Suppose α_1 , α_2 , and α_3 are the three roots of $P(x)$ and $|\alpha_1| > |\alpha_2| > |\alpha_3|$.

Letting $P_0(x) = P(x)$, for all $n \geq 1$ we let $P_{n+1}(w) = -P_n(x)P_n(-x)$ where $w = x^2$.

$$P_n(x) = x^3 - (\alpha_1^{2^n} + \alpha_2^{2^n} + \alpha_3^{2^n})x^2 + (\alpha_1^{2^n}\alpha_2^{2^n} + \alpha_1^{2^n}\alpha_3^{2^n} + \alpha_2^{2^n}\alpha_3^{2^n})x - (\alpha_1^{2^n}\alpha_2^{2^n}\alpha_3^{2^n})$$

If we let σ_n be the coefficient of x^2 in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{|\sigma_n|} = |\alpha_1|$.

If we let τ_n be the coefficient of x in $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\tau_n|}{|\sigma_n|}} = |\alpha_2|$

If we let ρ_n be the constant term of $P_n(x)$, then $\lim_{n \rightarrow \infty} \sqrt[2^n]{\frac{|\rho_n|}{|\tau_n|}} = |\alpha_3|$

In our example, $P_4(x) = x^6 - 14432666x^4 + 5217020805x^2 - 1$ and

$\sqrt[16]{\frac{1}{5217020805}} \approx 0.2469796037$. The actual root α_3 of least absolute value is $\alpha_3 \approx -0.2469796037$.

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$.

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$. Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)}x^{N-1} + \sigma_2^{(n)}x^{N-2} - \dots + (-1)^{N-1}\sigma_{N-1}^{(n)}x + (-1)^N\sigma_N^{(n)}$$

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$. Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)}x^{N-1} + \sigma_2^{(n)}x^{N-2} - \dots + (-1)^{N-1}\sigma_{N-1}^{(n)}x + (-1)^N\sigma_N^{(n)}$$

Then $\sigma_1^{(n)} = \sum_{1 \leq i \leq N} \alpha_i^{2^n}$,

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$.

Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)} x^{N-1} + \sigma_2^{(n)} x^{N-2} - \dots + (-1)^{N-1} \sigma_{N-1}^{(n)} x + (-1)^N \sigma_N^{(n)}$$

$$\text{Then } \sigma_1^{(n)} = \sum_{1 \leq i \leq N} \alpha_i^{2^n}, \sigma_2^{(n)} = \sum_{1 \leq i < j \leq N} \alpha_i^{2^n} \alpha_j^{2^n},$$

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$.

Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)}x^{N-1} + \sigma_2^{(n)}x^{N-2} - \dots + (-1)^{N-1}\sigma_{N-1}^{(n)}x + (-1)^N\sigma_N^{(n)}$$

Then $\sigma_1^{(n)} = \sum_{1 \leq i \leq N} \alpha_i^{2^n}$, $\sigma_2^{(n)} = \sum_{1 \leq i < j \leq N} \alpha_i^{2^n} \alpha_j^{2^n}, \dots$,
 $\sigma_N^{(n)} = \alpha_1^{2^n} \alpha_2^{2^n} \dots \alpha_N^{2^n}$.

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$.

Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)}x^{N-1} + \sigma_2^{(n)}x^{N-2} - \dots + (-1)^{N-1}\sigma_{N-1}^{(n)}x + (-1)^N\sigma_N^{(n)}$$

Then $\sigma_1^{(n)} = \sum_{1 \leq i \leq N} \alpha_i^{2^n}$, $\sigma_2^{(n)} = \sum_{1 \leq i < j \leq N} \alpha_i^{2^n} \alpha_j^{2^n}, \dots,$

$\sigma_N^{(n)} = \alpha_1^{2^n} \alpha_2^{2^n} \dots \alpha_N^{2^n}$. So in general, $\sigma_k^{(n)}$ is the sum of products of k factors chosen from $\alpha_1^{2^n}, \alpha_2^{2^n}, \dots, \alpha_N^{2^n}$.

Gräffe's Root Squaring Method

Suppose $P(x) = x^N + a_{N-1}x^{N-1} + \dots + a_1x + a_0$ is a polynomial of degree N with real roots $\alpha_1, \alpha_2, \dots, \alpha_N$ such that $|\alpha_1| > |\alpha_2| > \dots > |\alpha_N|$.

Letting $P_0(x) = P(x)$ and defining $P_{n+1}(w) = (-1)^N P(x)P(-x)$ where $w = x^2$ we have

$$P_n(x) = x^N - \sigma_1^{(n)}x^{N-1} + \sigma_2^{(n)}x^{N-2} - \dots + (-1)^{N-1}\sigma_{N-1}^{(n)}x + (-1)^N\sigma_N^{(n)}$$

Then $\sigma_1^{(n)} = \sum_{1 \leq i \leq N} \alpha_i^{2^n}$, $\sigma_2^{(n)} = \sum_{1 \leq i < j \leq N} \alpha_i^{2^n} \alpha_j^{2^n}, \dots,$

$\sigma_N^{(n)} = \alpha_1^{2^n} \alpha_2^{2^n} \dots \alpha_N^{2^n}$. So in general, $\sigma_k^{(n)}$ is the sum of products of k factors chosen from $\alpha_1^{2^n}, \alpha_2^{2^n}, \dots, \alpha_N^{2^n}$.

Moreover, $\lim_{n \rightarrow \infty} \sqrt[n]{|\sigma_1^{(n)}|} = |\alpha_1|$ and $\lim_{n \rightarrow \infty} \sqrt[n]{\frac{|\sigma_k^{(n)}|}{|\sigma_{k-1}^{(n)}|}} = |\alpha_k|$ for

$2 \leq k \leq N$.

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_1(x) = x^4 - 22x^3 + 125x^2 - 105x + 16$$

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$$P(3.609748279) \approx 513.9482482 \text{ and } P(-3.609748279) \approx 0.05224003$$

therefore, $\alpha_1 \approx -3.609748279$.

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$$P(3.609748279) \approx 513.9482482 \text{ and } P(-3.609748279) \approx 0.05224003$$

therefore, $\alpha_1 \approx -3.609748279$.

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_2(x) = x^4 - 234x^3 + 11037x^2 - 7025x + 256$$

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$$P(3.609748279) \approx 513.9482482 \text{ and } P(-3.609748279) \approx 0.05224003$$

therefore, $\alpha_1 \approx -3.609748279$.

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

$$P(2.829731735) \approx 232.3141638 \text{ and } P(-2.829731735) \approx 0.02550304$$

therefore, $\alpha_2 \approx -2.829731735$.

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_3(x) = x^4 - 32682x^3 + 118528181x^2 - 43699681x + 65536$$

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$$P(3.609748279) \approx 513.9482482 \text{ and } P(-3.609748279) \approx 0.05224003$$

therefore, $\alpha_1 \approx -3.609748279$.

$$|\alpha_2| \approx \sqrt[16]{\left| \frac{\sigma_2^{(4)}}{\sigma_1^{(4)}} \right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

$$P(2.829731735) \approx 232.3141638 \text{ and } P(-2.829731735) \approx 0.02550304$$

therefore, $\alpha_2 \approx -2.829731735$.

$$|\alpha_3| \approx \sqrt[16]{\left| \frac{\sigma_3^{(4)}}{\sigma_2^{(4)}} \right|} = \sqrt[16]{\frac{1894126393761729}{14046073305350949}} \approx 0.8822991819$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$P(3.609748279) \approx 513.9482482$ and $P(-3.609748279) \approx 0.05224003$
therefore, $\alpha_1 \approx -3.609748279$.

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

$P(2.829731735) \approx 232.3141638$ and $P(-2.829731735) \approx 0.02550304$
therefore, $\alpha_2 \approx -2.829731735$.

$$|\alpha_3| \approx \sqrt[16]{\left|\frac{\sigma_3^{(4)}}{\sigma_2^{(4)}}\right|} = \sqrt[16]{\frac{1894126393761729}{14046073305350949}} \approx 0.8822991819$$

$P(0.8822991819) \approx 0.000020487$ and $P(-0.8822991819) \approx 4.110279912$
therefore, $\alpha_3 \approx 0.8822991819$.

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$
$$\alpha_1 \approx -3.609748279.$$

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

$$P(2.829731735) \approx 232.3141638 \text{ and } P(-2.829731735) \approx 0.02550304$$

therefore, $\alpha_2 \approx -2.829731735$.

$$|\alpha_3| \approx \sqrt[16]{\left|\frac{\sigma_3^{(4)}}{\sigma_2^{(4)}}\right|} = \sqrt[16]{\frac{1894126393761729}{14046073305350949}} \approx 0.8822991819$$

$$P(0.8822991819) \approx 0.000020487 \text{ and } P(-0.8822991819) \approx 4.110279912$$

therefore, $\alpha_3 \approx 0.8822991819$.

$$|\alpha_4| \approx \sqrt[16]{\left|\frac{\sigma_4^{(4)}}{\sigma_3^{(4)}}\right|} = \sqrt[16]{\frac{4294967296}{1894126393761729}} \approx 0.4438353932$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$
$$\alpha_1 \approx -3.609748279.$$

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$
$$\alpha_2 \approx -2.829731735.$$

$$|\alpha_3| \approx \sqrt[16]{\left|\frac{\sigma_3^{(4)}}{\sigma_2^{(4)}}\right|} = \sqrt[16]{\frac{1894126393761729}{14046073305350949}} \approx 0.8822991819$$
$$\alpha_3 \approx 0.8822991819.$$

$$|\alpha_4| \approx \sqrt[16]{\left|\frac{\sigma_4^{(4)}}{\sigma_3^{(4)}}\right|} = \sqrt[16]{\frac{4294967296}{1894126393761729}} \approx 0.4438353932$$

$$P(0.4438353932) \approx -5.164527333 \text{ and}$$

$$P(-0.4438353932) \approx -4.672 \times 10^{-6} \text{ therefore.}$$

Example

Approximate the roots of $P(x) = x^4 + 6x^3 + 7x^2 - 7x - 4$ using four steps of the root squaring method.

$$P_4(x) = x^4 - 831056762x^3 + 14046073305350949x^2 - 1894126393761729x + 4294967296$$

$$|\alpha_1| \approx \sqrt[16]{|\sigma_1^{(4)}|} = \sqrt[16]{831056762} \approx 3.609748279$$

$$\alpha_1 \approx -3.609748279.$$

$$|\alpha_2| \approx \sqrt[16]{\left|\frac{\sigma_2^{(4)}}{\sigma_1^{(4)}}\right|} = \sqrt[16]{\frac{14046073305350949}{831056762}} \approx 2.829731735$$

$$\alpha_2 \approx -2.829731735.$$

$$|\alpha_3| \approx \sqrt[16]{\left|\frac{\sigma_3^{(4)}}{\sigma_2^{(4)}}\right|} = \sqrt[16]{\frac{1894126393761729}{14046073305350949}} \approx 0.8822991819$$

$$\alpha_3 \approx 0.8822991819.$$

$$|\alpha_4| \approx \sqrt[16]{\left|\frac{\sigma_4^{(4)}}{\sigma_3^{(4)}}\right|} = \sqrt[16]{\frac{4294967296}{1894126393761729}} \approx 0.4438353932$$

$$\alpha_4 \approx -0.4438353932.$$

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2) \dots (x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2)\dots(x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2) \dots (x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials: Every **symmetric** polynomial in variables $\alpha_1, \alpha_2, \dots, \alpha_N$ can be written in a unique way as a polynomial in $\sigma_1, \sigma_2, \dots, \sigma_N$.

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2)\dots(x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials: Every **symmetric** polynomial in variables $\alpha_1, \alpha_2, \dots, \alpha_N$ can be written in a unique way as a polynomial in $\sigma_1, \sigma_2, \dots, \sigma_N$.

A polynomial in variables x_1, x_2, \dots, x_m is symmetric if it is left unchanged after the variables x_1, x_2, \dots, x_m are permuted for every permutation of the m variables.

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2) \dots (x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials: Every **symmetric** polynomial in variables $\alpha_1, \alpha_2, \dots, \alpha_N$ can be written in a unique way as a polynomial in $\sigma_1, \sigma_2, \dots, \sigma_N$.

A polynomial in variables x_1, x_2, \dots, x_m is symmetric if it is left unchanged after the variables x_1, x_2, \dots, x_m are permuted for every permutation of the m variables.

For example, for $N = 3$, $\sigma_1 = \alpha_1 + \alpha_2 + \alpha_3$, $\sigma_2 = \alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_3\alpha_1$, and $\sigma_3 = \alpha_1\alpha_2\alpha_3$.

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2)\dots(x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials: Every **symmetric** polynomial in variables $\alpha_1, \alpha_2, \dots, \alpha_N$ can be written in a unique way as a polynomial in $\sigma_1, \sigma_2, \dots, \sigma_N$.

For example, for $N = 3$, $\sigma_1 = \alpha_1 + \alpha_2 + \alpha_3$, $\sigma_2 = \alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_3\alpha_1$, and $\sigma_3 = \alpha_1\alpha_2\alpha_3$.

Then $\alpha_1^2 + \alpha_2^2 + \alpha_3^2$ is a symmetric polynomial in the three variables α_1, α_2 , and α_3 .

Fundamental Theorem of Symmetric Polynomials

Suppose

$$P(x) = (x - \alpha_1)(x - \alpha_2) \dots (x - \alpha_N) = x^N + \sigma_1 x^{N-1} + \dots + \sigma_{N-1} x + \sigma_N$$

then σ_k is the sum of the products of k factors chosen from $\alpha_1, \alpha_2, \dots, \alpha_N$.

Fundamental Theorem of Symmetric Polynomials: Every **symmetric** polynomial in variables $\alpha_1, \alpha_2, \dots, \alpha_N$ can be written in a unique way as a polynomial in $\sigma_1, \sigma_2, \dots, \sigma_N$.

For example, for $N = 3$, $\sigma_1 = \alpha_1 + \alpha_2 + \alpha_3$, $\sigma_2 = \alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_3\alpha_1$, and $\sigma_3 = \alpha_1\alpha_2\alpha_3$.

Then $\alpha_1^2 + \alpha_2^2 + \alpha_3^2$ is a symmetric polynomial in the three variables α_1, α_2 , and α_3 .

$$\alpha_1^2 + \alpha_2^2 + \alpha_3^2 = (\alpha_1 + \alpha_2 + \alpha_3)^2 - 2(\alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_3\alpha_1) = \sigma_1^2 - 2\sigma_2.$$