

Elementary seasonal adjustment of economic data with JDemetra+: Module IV – Quality assessment

Dr Karsten Webel / Deutsche Bundesbank, DG Statistics
Virtual Seminar Series, 19-23 October 2020

Aims

Theory

- Basic understanding \leadsto Ideas, concepts
- Approach \leadsto X-11
- Pretreatment \leadsto RegARIMA models

Application


- Software \leadsto JDemetra+ (JD+)
- Specification \leadsto Options
- Results \leadsto Interpretation, quality assessment

Discussion

- Your questions \leadsto Practical problems

Road map

Structure of JDemetra+



Overview

Questions

- Data \leadsto Need for seasonal adjustment?
- SA data \leadsto Quality of seasonal adjustment?


Answers

- Dependence measures \leadsto Autocorrelations, periodogram
- Tests \leadsto Seasonality
- Heuristics \leadsto M - & Q -statistics

Dependence measures (I/III)

Autocorrelation function (ACF)

Data_{t-1} Data_t Data_{t+1} ... Data_{t+h-1} Data_{t+h} Data_{t+h+1}



Normalised time-constant linear dependence




$$\text{ACF}(h) \in [-1, 1]$$

Assumption

- $\text{Data} \rightsquigarrow$ Weakly stationary

Dependence measures (II/III)


Partial autocorrelation function (PACF)



Dependence measures (III/III)

Periodogram: AR (1) process, unit WN variance


Dominance of long cycles



Deutsche Bundesbank

SPMWSCH Out


Dominance of short cycles



Deutsche Bundesbank

SPMWSCH Out

No dominance of any cycle



Deutsche Bundesbank

SPMWSCH Out

Cycles per year	1	2	3	4	5	6
Duration in months	12	6	4	3	2.4	2
Seasonal frequency	$\frac{\pi}{6}$	$\frac{2\pi}{6}$	$\frac{3\pi}{6}$	$\frac{4\pi}{6}$	$\frac{5\pi}{6}$	π
Duration in quarters	4	2				
Seasonal frequency	$\frac{\pi}{2}$	π				

Seasonality tests in JD+ (I/VIII)

Overview

Name	Variable checked for significance	Short
Modified <i>QS</i> -test	Autocorrelations at seasonal lags	QS
Friedman-test	ANOVA with repeated measures on intra-year ranks	FT
Kruskal-Wallis-test	ANOVA without repeated measures on overall ranks	KW
Test for spectral peaks	Tukey and AR(30) spectra at seasonal frequencies	SP
Periodogram-test	Weighted sum of periodogram at seasonal frequencies	PD
<i>F</i> -test on seasonal dummies	Effects of seasonal dummies in the " <i>(pdq)</i> (000) + mean + seasonal dummies" model*	SD

* Variant 1: $(pdq) = (011)$, variant 2: (pdq) according to automatic model identification.

Seasonality tests in JD+ (II/VIII)

Ljung-Box-test

$$LB_H = \sum_{h=1}^H w(h) \text{ACF}^2(h), \quad \text{with } w(h) = \frac{T(T+2)}{T-h}$$

Hypothesis

- Data \rightsquigarrow Zero autocorrelations (i.e. independence)

Decision

- Rejection if LB_H too large

Seasonality tests in JD+ (III/VIII)

Modified QS -test

$$QS = \begin{cases} w(12) \text{ACF}^2(12) + w(24) [\max\{0, \text{ACF}(24)\}]^2, & \text{ACF}(12) > 0 \\ 0, & \text{ACF}(12) \leq 0 \end{cases}$$

Hypothesis

- Data \leadsto No positive autocorrelation at seasonal lags (i.e. no stable seasonality)

Decision

- Rejection if QS too large

Seasonality tests in JD+ (IV/VIII)

Friedman-test

$$FT = \frac{\text{Variance of month-specific mean ranks}}{\text{Variance of ranks within months}}$$

Rank assignment

- Within each year $\leadsto 1 \leq \text{rank}_{ij} \leq \tau$

Hypothesis

- Data \leadsto Same month-specific mean ranks (i.e. no stable seasonality)

Decision

- Rejection if FT -statistics too large

Seasonality tests in JD+ (V/VIII)

Kruskal-Wallis-test

$$KW = \frac{\text{Variance of month-specific mean ranks}}{\text{Variance of ranks within months}}$$

Rank assignment

- Within observation period $\leadsto 1 \leq \text{rank}_{ij} \leq T$

Hypothesis

- Data \leadsto Same month-specific mean ranks (i.e. no stable seasonality)

Decision

- Rejection if KW -statistics too large

Seasonality tests in JD+ (VI/VIII)

Test for spectral peaks

Idea

- Spectral peaks \leadsto Visual significance
- Tukey & AR(30) spectra \leadsto Combination

Hypothesis

- Data \leadsto No visually significant seasonal peaks (i.e. no seasonality)

Decision

- Rejection if too many visually significant seasonal peaks
- Consideration \leadsto Location of seasonal peaks

Seasonality tests in JD+ (VII/VIII)

Periodogram-test

$$PD = \text{Rescaled} \left[2 \sum_{j=1}^5 I(\text{SF}_j) + I(\text{SF}_6) \times \begin{cases} 1, & T \text{ even} \\ 0, & T \text{ odd} \end{cases} \right]$$

Hypothesis

- Data \rightsquigarrow No seasonal peaks (i.e. no seasonality)

Decision

- Rejection if PD -statistics too large

Seasonality tests in JD+ (VIII/VIII)

F-test on seasonal dummies

$$SD = \text{Rescaled } \left(\hat{\beta}^\top \hat{\Sigma}_{\hat{\beta}}^{-1} \hat{\beta} \right) \text{ from model}$$
$$\phi_p(B)(1-B)^d \left(\text{Data}_t - \hat{\beta}^\top \mathbf{Dummies}_t \right) = \mu + \theta_q(B) \varepsilon_t$$

Hypothesis

- Data \leadsto Zero effects for seasonal dummies (i.e. no stable seasonality)

Decision

- Rejection if *SD*-statistics too large

Additional F -tests (I/II)

Stable seasonality

$$F_S = \frac{\text{Variance of month-specific mean SI ratios}}{\text{Variance of SI ratios within months}}$$

Hypothesis

- Seasonal factors \leadsto Same month-specific averages (i.e. no stable seasonality)

Decision

- Rejection if F_S too large

Additional F -tests (II/II)

Moving seasonality

$$\left| (\hat{si})_{ij} - 100 \right| = \text{Year}_i + \text{Month}_j + \text{Residual}_{ij}$$

Hypothesis

- SI ratios \leadsto Absence of annual effects (i.e. no moving seasonality, $\text{Year}_1 = \text{Year}_2 = \dots = \text{Year}_T$)

Decision

- Rejection if F_M -statistic too large

Heuristics (I/III)

M-statistics: basic idea

Indication

- Seasonal adjustment \leadsto Quality
- Development \leadsto Statistics Canada

Construction

- Domain $\leadsto [0, 3]$
- Acceptance region $\leadsto [0, 1]$

Warning

- Relevance \leadsto Limited

Heuristics (II/III)

M-statistics: *M*7

$$M7 = \sqrt{\frac{1}{2}(T_1 + T_2)}, \quad \text{with } T_1 = \min \left\{ \frac{7}{F_S}, 9 \right\}, T_2 = \min \left\{ \frac{3F_M}{F_S}, 9 \right\}$$


Interpretation

- “Ratio” \leadsto **Stable** (F_S) & **moving** (evolutive) **seasonality** (F_M)

Heuristics (III/III)

Identifiable seasonality

Combined test for presence of identifiable seasonality



Source: Ladiray and Quenneville (2001).
Deutsche Bundesbank

S3PR0139.Chart