

Bayesian Learning (732A91) Lab1 Report

Christoforos Spyretos (chrsp415) & Marketos Damigos (marda352)

2022-04-06

Assignment 1 *Daniel Bernoulli*

Let $y_1, y_2, \dots, y_n \sim \text{Bern}(\theta)$, and the obtained sample has 13 successes out of 50 trials. The $\text{Beta}(a_0, b_0)$ prior has $a_0 = b_0 = 5$.

Task 1a

The mean value and the standard deviation of the Beta distribution for θ are calculated by the below formulas:


$$\begin{aligned} E[\theta] &= \frac{a_0 + s}{a_0 + s + b_0 + f} \\ &= \frac{18}{60} \\ &= 0.3 \end{aligned}$$

$$\begin{aligned} \text{Var}[\theta] &= \frac{(a_0 + s)(b_0 + f)}{((a_0 + s) + (b_0 + f))^2((a_0 + s) + (b_0 + f) + 1)} \\ &= \frac{18 \cdot 42}{(18 + 42)^2(18 + 42 + 1)} \\ &= 0.003442623 \end{aligned}$$


$$SD[\theta] = \sqrt{\text{Var}[\theta]} = 0.05867387$$

Where a_0 and b_0 are the arguments of the Beta prior, s is the number of successes and f is the number of failures.

Mean Values Graphs



Standard Deviation Values Graphs




From the above plots, it could be seen that both posterior's mean and standard deviation values converge to the actual mean and standard deviation values, respectively. More specifically, between 0 and approximately 250 draws in both graphs, some of the posterior's values abstain from true values. However, after the 250 draws, the posterior's values start to converge to the true ones in both graphs.

Task 1b

The posterior probability $P(\theta < 0.3|y)$ equals 0.506, and the exact probability value from the Beta posterior is 0.5150226; thus, it could be assumed that both values are pretty similar.


Task 1c



The above plot illustrates the density of the log-odds values $\phi = \log \frac{\theta}{1-\theta}$, where θ takes values from simulated draws from the Beta posterior.

Assignment 2 *Log-normal distribution and the Gini coefficient.*

Task 2a



Appendix

```
knitr::opts_chunk$set(echo = TRUE, warning = FALSE)
knitr::opts_chunk$set(tidy.opts = list(width.cutoff = 60), tidy = TRUE)
# task 1a true mean, var & sd
a0 <- 5
b0 <- 5
n <- 50
s <- 13
f <- 50 - 13
mean_true <- (a0 + s)/(a0 + s + b0 + f)
var_true <- ((a0 + s) * (b0 + f))/(((a0 + s + b0 + f)^2) * (a0 +
  s + b0 + f + 1))
sd_true <- sqrt(var_true)
set.seed(12345)

# calculate posterior's mean
mean_posterior = c()
for (i in 1:1000) {
  # rbeta generates random deviates
  mean_posterior[i] = mean(rbeta(n = i, shape1 = a0 + s, shape2 = b0 +
    f))
}

# calculate posterior's sd
sd_posterior = c()
for (i in 1:1000) {
  sd_posterior[i] = sd(rbeta(n = i, shape1 = a0 + s, shape2 = b0 +
    f))
}

# df for plots
df_plot1 <- data.frame(draws = 1:1000, mean_true = mean_true,
  sd_true = sd_true, mean_posterior = mean_posterior, sd_posterior = sd_posterior)

library(ggplot2)

# plot for mean values
ggplot(df_plot1) + geom_point(aes(x = draws, y = mean_posterior,
  color = "navy")) + geom_line(aes(x = draws, y = mean_true,
  color = "red4")) + theme(legend.position = "right") + scale_color_manual(values = c("navy",
  "red4"), name = "Samples", labels = c("Posterior's Mean Values",
  "True Mean Values")) + ggtitle("Mean Values Graphs") + xlab("Number of Draws") +
  ylab("Mean value of each sample") + theme_classic()

# plot for sd values
ggplot(df_plot1) + geom_point(aes(x = draws, y = sd_posterior,
  color = "navy")) + geom_line(aes(x = draws, y = sd_true,
  color = "red4")) + theme(legend.position = "right") + scale_color_manual(values = c("navy",
  "red4"), name = "Samples", labels = c("Posterior's Sd Values",
  "True Sd Values")) + ggtitle("Standard Deviation Values Graphs") +
  xlab("Number of Draws") + ylab("Sd value of each sample") +
  theme_classic()
set.seed(12345)
```

```

# generates 1,000 random deviates.
posterior_sample <- rbeta(n = 1000, shape1 = a0 + s, shape2 = b0 +
  f)

# posterior probability
posterior_prob <- sum(posterior_sample < 0.3)/1000

# exact posterior prob pbeta the distribution function
exact_prob <- pbeta(q = 0.3, shape1 = a0 + s, shape2 = b0 + f)
phi <- log(posterior_sample/(1 - posterior_sample))

df_plot2 <- data.frame(phi = phi)

ggplot(df_plot2, aes(x = phi)) + geom_histogram(bins = 30, color = "navy",
  fill = "steelblue2", aes(y = ..density..)) + geom_density(colour = "red4",
  size = 1) + ggtitle("Histogram of Log-Odds") + xlab("Log-Odds Values") +
  ylab("Density") + theme_classic()

# observations
obs <- c(33, 24, 48, 32, 55, 74, 23, 76, 17)

tau_2 <- sum((log(obs) - 3.5)^2)/9

# generates 10,000 random deviates.
set.seed(12345)
sigma2 <- c()
for (i in 1:10000) {
  values <- rchisq(1, 9)
  sigma2[i] <- 9 * tau_2/values
}

df_plot2 <- data.frame(sigma2 = sigma2)

ggplot(df_plot2, aes(x = sigma2)) + geom_histogram(bins = 30,
  color = "navy", fill = "steelblue2", aes(y = ..density..)) +
  geom_density(colour = "red4", size = 1) + scale_x_continuous(limits = c(0,
  1.5)) + ggtitle("Posterior Distribution") + xlab("Sigma2 Values") +
  ylab("Density") + theme_classic()

```