

Homework 1

Group 1

Contents

1	Introduction	2
2	Statement of the Problem	2
3	Data Exploration	2
3.1	Imputing Missing Values	4
3.2	Correlation Matrix	5
4	Data Preparation	6
5	Models Built	6
5.1	Model 1	6
6	Selected Model	7
7	Appendix A	8
7.1	Session Info	8
7.2	Citations	8
7.3	Data Dictionary	8
7.4	R source code	8

Prepared for:

Dr. Nathan Bastian

City University of New York - Data 621

Prepared by:

Group 1

Senthil Dhanapal

Yadu Chittampalli

Christophe Hunt

1 Introduction

The ability to analyze and predict performance of a professional baseball team using many dimensions is critical to competitive success for our organization. Therefore, we have analyzed the records of numerous professional baseball team from the years 1871 to 2006. Our hope is that the following report and the resulting predictive models will better inform the organization and assist in making data driven decisions moving forward.

"The goal of a baseball team is to win more games than any other team. Since one team has very little control over the number of games other teams win, the goal is essentially to win as many games as possible. Therefore, it is of interest to measure the player's contribution to the team's wins." Grabiner, B. D. ¹ While we do not have the variables at the player's individual contribution level, we do have the entire teams contributions as an aggregate and will analyze that information.

2 Statement of the Problem

The purpose of this report is to determine the batting, baserun, pitching, and fielding effects on a baseball team's ability to win.

3 Data Exploration

Note that each record has the performance of the team for the given year, with all of the statistics adjusted to match the performance of a 162 game season. The following Table 1 - Descriptive Statistics provides the detailed descriptive statistics regarding our variable of interest - Number of Wins and our possible explanatory variables.

We noted that several variables were missing a nontrivial amount of observations and these variables are Strikeouts by batters, Stolen Bases, Caught stealing, Batters hit by pitch (get a free base), Strikeouts by pitcher, and Double plays. So we will need to address the missing values for further analysis.

Histograms of all of the variables have been plotted below so that the distribution of the data can be visualized. In the distribution for the number of walks allowed, only two bars exist due to the excessive number of outliers.

Additionally, the skewness of each variable has been indicated in Table 1 - Descriptive Statistics. Two variables with a significant amount of skew are the number of base hits by batters and the number of walks allowed. Correspondingly, these two variables had a skew of 1.57 and 6.74 respectively.

**Table 1 : Descriptive Statistics
16 Variables 2276 Observations**

Number of wins

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	108	81	82	15.8	-0.4	54	104

lowest : 0 12 14 17 21, highest: 128 129 134 135 146

¹(Grabiner, B. D. (n.d.). The Sabermetric Manifesto. Retrieved September 10, 2016 from <http://seanlahman.com/baseball-archive/sabermetrics/sabermetric-manifesto/>)

Base Hits by batters (1B,2B,3B,HR)

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	569	1469	1454	144.6	1.6	1282	1695

lowest : 891 992 1009 1116 1122, highest: 2333 2343 2372 2496 2554



Doubles by batters (2B)

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	240	241	238	46.8	0.2	167	320


lowest : 69 112 113 118 123, highest: 382 392 393 403 458



Triples by batters (3B)

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	144	55	47	27.9	1.1	23	108

lowest : 0 8 9 11 12, highest: 166 190 197 200 223



Homeruns by batters (4B)

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	243	100	102	60.5	0.2	14	199

lowest : 0 3 4 5 6, highest: 247 249 257 260 264



Walks by batters

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	533	502	512	122.7	-1	248	670

lowest : 0 12 29 34 45, highest: 815 819 824 860 878



Strikeouts by batters

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2174	102	822	736	750	248.5	-0.3	359	1103


lowest : 0 66 67 72 74, highest: 1303 1320 1326 1335 1399



Stolen bases

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2145	131	348	125	101	87.8	2	35	302


lowest : 0 14 18 19 20, highest: 562 567 632 654 697



Caught stealing

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
1504	772	128	53	49	23	2	24	91

lowest : 0 7 11 12 14, highest: 171 186 193 200 201



Batters hit by pitch (get a free base)

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
191	2085	55	59	58	13	0.3	40	82


lowest : 29 30 35 38 39, highest: 87 88 89 90 95



Hits allowed

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	843	1779	1518	1406.8	10.3	1316	2563

lowest : 1137 1168 1184 1187 1202
highest: 16038 16871 20088 24057 30132



Homeruns allowed

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	256	106	107	61.3	0.3	18	209

lowest : 0 3 4 5 6, highest: 291 297 301 320 343



Walks allowed

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	535	553	536.5	166.4	6.7	377	757

lowest : 0 119 124 131 140, highest: 2169 2396 2840 2876 3645



Strikeouts by pitchers

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2174	102	823	818	813.5	553.1	22.2	421	1173


lowest : 0 181 205 208 252
highest: 3450 4224 5456 12758 19278



Errors

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
2276	0	549	246	159	227.8	3	100	716

lowest : 65 66 68 72 74, highest: 1567 1728 1740 1890 1898



Double Plays

n	missing	unique	Mean	Median	SD	Skew	.05 freq	.95 freq
1990	286	144	146	149	26.2	-0.4	98	186

lowest : 52 64 68 71 72, highest: 215 218 219 225 228



3.1 Imputing Missing Values

In order to address the missing values in our variables we used a nonparametric imputation method (Random Forest) to impute missing values. We chose a nonparametric method due to several variables having significant skew and greater than expected kurtosis values.

3.2 Correlation Matrix

After completing the imputation, we can implement a correlation matrix to better understand the correlation between variables in the data set. The below matrix is the results and as expected, Number of Wins appears to be most correlated to Base Hits by batters (1B,2B,3B,HR).




Figure 1: Correlation Plot of Training Data Set with imputed values

4 Data Preparation

First, we chose to eliminate two variables that had a significant number of missing data points. These variables were Batters hit by pitch (get a free base) and Caught stealing, which were missing 91.6% and 33.9% respectively.

Missing values in the remaining columns had been imputed using the random forest method as previous discussed in section 3.1.

5 Models Built

5.1 Model 1

We used the variable Base Hits by batters (1B,2B,3B,HR) which is the most correlated variable to Number of Wins as indicated in the correlation matrix. This is expected as Base Hits are necessary to win any game. Additionally, Strikeouts by batters would be negatively correlated Number of Wins because if a batter strikes out they are no able to provide runs which are critically to win.

Table 1:

	Model 1
	Number of wins
Base Hits by batters (1B,2B,3B,HR)	0.053*** (0.002)
$\sqrt{\text{Strikeouts by batters}}$	0.647*** (0.065)
Constant	-14.637*** (4.500)
Observations	2,276
R ²	0.187
Adjusted R ²	0.186
Residual Std. Error	14.210 (df = 2273)
F Statistic	261.384*** (df = 2; 2273)
Note:	*p<0.1; **p<0.05, ***p<0.01

6 Selected Model

Decide on the criteria for selecting the best multiple linear regression model. Will you select a model with slightly worse performance if it makes more sense or is more parsimonious? Discuss why you selected your model. For the multiple linear regression model, will you use a metric such as Adjusted R², RMSE, etc.? Be sure to explain how you can make inferences from the model, discuss multi-collinearity issues (if any), and discuss other relevant model output. Using the training data set, evaluate the multiple linear regression model based on (a) mean squared error, (b) R², (c) F-statistic, and (d) residual plots. Make predictions using the evaluation data set.

% Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fas.harvard.edu %
Date and time: Thu, Sep 22, 2016 - 8:16:45 PM

Table 2:	
	<i>Dependent variable:</i>
	'Number of wins'
'Base Hits by batters (1B,2B,3B,HR)'	0.053*** (0.002)
sqrt('Strikeouts by batters')	0.647*** (0.065)
Constant	−14.637*** (4.500)
Observations	2,276
R ²	0.187
Adjusted R ²	0.186
Residual Std. Error	14.210 (df = 2273)
F Statistic	261.384*** (df = 2; 2273)
Note:	*p<0.1; **p<0.05; ***p<0.01

7 Appendix A

7.1 Session Info

- R version 3.3.1 (2016-06-21), x86_64-w64-mingw32
- Locale: LC_COLLATE=English_United States.1252, LC_CTYPE=English_United States.1252, LC_MONETARY=English_United States.1252, LC_NUMERIC=C, LC_TIME=English_United States.1252
- Base packages: base, datasets, graphics, grDevices, methods, parallel, stats, utils
- Other packages: bibtex 0.4.0, corrplot 0.77, doParallel 1.0.10, dplyr 0.5.0, e1071 1.6-7, foreach 1.4.3, Formula 1.2-1, ggplot2 2.1.0, Hmisc 3.17-4, iterators 1.0.8, itertools 0.1-3, knitr 1.14, lattice 0.20-34, magrittr 1.5, missForest 1.4, pacman 0.4.1, plyr 1.8.4, randomForest 4.6-12, rJava 0.9-8, scales 0.4.0, stargazer 5.2, stringr 1.1.0, survival 2.39-5, tidyr 0.6.0, xlsx 0.5.7, xlsxjars 0.6.1
- Loaded via a namespace (and not attached): acepack 1.3-3.3, assertthat 0.1, bitops 1.0-6, chron 2.3-47, class 7.3-14, cluster 2.0.4, codetools 0.2-14, colorspace 1.2-6, data.table 1.9.6, DBI 0.5-1, digest 0.6.10, evaluate 0.9, foreign 0.8-67, formatR 1.4, grid 3.3.1, gridExtra 2.2.1, gtable 0.2.0, htmltools 0.3.5, httr 1.2.1, latticeExtra 0.6-28, lubridate 1.6.0, Matrix 1.2-7.1, munsell 0.4.3, nnet 7.3-12, R6 2.1.3, RColorBrewer 1.1-2, Rcpp 0.12.7, RCurl 1.95-4.8, RefManager 0.11.0, RJSONIO 1.3-0, rmarkdown 1.0, rpart 4.1-10, splines 3.3.1, stringi 1.1.1, tibble 1.2, tools 3.3.1, XML 3.98-1.4, yaml 2.1.13

7.2 Citations

7.3 Data Dictionary

VARIABLE.NAME..	DEFINITION	THEORETICAL.EFFECT
INDEX	Identification Variable (do not use)	None
TARGET_WINS	Number of wins	NA
TEAM_BATTING_H	Base Hits by batters (1B,2B,3B,HR)	Positive Impact on Wins
TEAM_BATTING_2B	Doubles by batters (2B)	Positive Impact on Wins
TEAM_BATTING_3B	Triples by batters (3B)	Positive Impact on Wins
TEAM_BATTING_HR	Homeruns by batters (4B)	Positive Impact on Wins
TEAM_BATTING_BB	Walks by batters	Positive Impact on Wins
TEAM_BATTING_HBP	Batters hit by pitch (get a free base)	Positive Impact on Wins
TEAM_BATTING_SO	Strikeouts by batters	Negative Impact on Wins
TEAM_BASERUN_SB	Stolen bases	Positive Impact on Wins
TEAM_BASERUN_CS	Caught stealing	Negative Impact on Wins
TEAM_FIELDING_E	Errors	Negative Impact on Wins
TEAM_FIELDING_DP	Double Plays	Positive Impact on Wins
TEAM_PITCHING_BB	Walks allowed	Negative Impact on Wins
TEAM_PITCHING_H	Hits allowed	Negative Impact on Wins
TEAM_PITCHING_HR	Homeruns allowed	Negative Impact on Wins
TEAM_PITCHING_SO	Strikeouts by pitchers	Positive Impact on Wins

7.4 R source code