

Homework 5

Christophe Hunt

February 28, 2017

Contents

1	Page 228: problem 1	2
1.1	Model to solve for long-term percentage	2
2	Page 232: problem 1	3
3	Page 240: problem 1	3
4	Page 240: problem 2	4

1 Page 228: problem 1

Consider a model for the long-term dining behavior of the students at College USA. It is found that 25% of the students who eat at the college's Grease Dining Hall return to eat there again, whereas those who eat at Sweet Dining Hall have a 93% return rate. These are the only two dining halls available on campus, and assume that all students eat at a one of these halls. Formulate a model to solve for the long-term percentage of students eating at each hall.

Table 1: Present - Next State for Dining

		NEXT STATE	
		Grease Dining Hall	Sweet Dining Hall
PRESENT STATE	Grease Dining Hall	.25	.75
	Sweet Dining Hall	.7	.93

1.1 Model to solve for long-term percentage

$$Grease_{n+1} = .25 Grease_n + .7 Sweet_N$$

$$Sweet_{n+1} = .75 Grease_n + .93 Sweet_N$$

2 Page 232: problem 1

Consider a stereo with CD player, FM-AM radio tuner, speakers (dual) and power amplifier (PA) components, as displayed with the reliability. Determine the system's reliability. what assumptions are required in your model?




Figure 1: image.

Component Reliability

$$R_{s1} = 0.95$$

$$R_{s2} = 0.98 + .97 - (.98 * .97) = 0.9994$$

$$R_{s3} = .99 + .99 - (.99 * .99) = 0.9999$$

Entire system reliability:

$$R_{s1,s2,s3} = .95 * 0.9994 * 0.9999 = 0.9493351$$

3 Page 240: problem 1

Use the basic linear model $y = ax + b$ to fit the following data sets. Provide the model, provide the values of SEE, SSR, SST, and R^2 , and provide a residual plot.

```
height <- c(60:80)
weight <- c(132, 136, 141, 145, 150, 155, 160, 165, 170,
           175, 180, 185, 190, 195, 201, 206, 212, 218,
           223, 229, 234)
```

```
slope <- function(x,y){
  return(((length(x)*sum(x*y)) - sum(x)*sum(y)) /
          ((length(x)*sum(x^2)) - sum(x)^2))
}
slope(x = height, y = weight)
```

```
## [1] 5.136364
```

```
intercept <- function(x,y){
  (sum(x^2)*sum(y) - sum(x*y)*sum(x)) /
  ((length(x)*sum(x^2)) - sum(x)^2)
}
intercept(x = height, y = weight)
```

```
## [1] -178.4978
```

The linear model $y = ax + b$ for this data set is $\$y_{\{weight\}} = 5.14x_{\{height\}} - 178.5$

We can verify the results with the `lm` function in base R.

```
library(stargazer)
```

```
##
```

```
## Please cite as:
```

```
## Hlavac, Marek (2015). stargazer: Well-Formatted Regression and Summary Statistics Tables.
```

```
## R package version 5.2. http://CRAN.R-project.org/package=stargazer
```

```
stargazer(lm(weight ~ height))
```

% Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac@fas.harvard.edu %
Date and time: Sat, Mar 04, 2017 - 2:30:11 PM

Table 2:	
	<i>Dependent variable:</i>
	weight
height	5.136*** (0.041)
Constant	-178.498*** (2.883)
Observations	21
R ²	0.999
Adjusted R ²	0.999
Residual Std. Error	1.139 (df = 19)
F Statistic	15,668.140*** (df = 1; 19)
Note:	*p<0.1; **p<0.05; ***p<0.01

4 Page 240: problem 2