

多元函数极值

二元函数Taylor公式的获得

$$\Delta z = f(x, y) - f(x_0, y_0) = ?$$

二元函数Taylor公式的获得

$$\Delta z = f(x, y) - f(x_0, y_0) = ?$$

记 $\Delta x = x - x_0, \Delta y = y - y_0$

对确定的 x, y, x_0, y_0 , 作一元辅助函数

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

二元函数Taylor公式的获得

$$\Delta z = f(x, y) - f(x_0, y_0) = ?$$

记 $\Delta x = x - x_0, \Delta y = y - y_0$

对确定的 x, y, x_0, y_0 , 作一元辅助函数

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

则显见 $\Delta z = F(1) - F(0)$

二元函数Taylor公式的获得

$$\Delta z = f(x, y) - f(x_0, y_0) = ?$$

记 $\Delta x = x - x_0, \Delta y = y - y_0$

对确定的 x, y, x_0, y_0 , 作一元辅助函数

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

则显见 $\Delta z = F(1) - F(0)$

若能写出 $F(t)$ 的表达式, 则可得 Δz 的表达式

二元函数Taylor公式的获得

$$\Delta z = f(x, y) - f(x_0, y_0) = ?$$

记 $\Delta x = x - x_0, \Delta y = y - y_0$

对确定的 x, y, x_0, y_0 , 作一元辅助函数

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

则显见 $\Delta z = F(1) - F(0)$

若能写出 $F(t)$ 的表达式, 则可得 Δz 的表达式

根据一元函数Taylor公式, 现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F'(t) = \frac{\partial f}{\partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x + \frac{\partial f}{\partial y}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta y$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F'(t) = \frac{\partial f}{\partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x + \frac{\partial f}{\partial y}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta y$$

$$F''(t) = \frac{\partial^2 f}{\partial x^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta x)^2 + 2\frac{\partial^2 f}{\partial y \partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x \Delta y \\ + \frac{\partial^2 f}{\partial y^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta y)^2 = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0 + t\Delta x, y_0 + t\Delta y); \\ \dots\dots\dots$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \dots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F'(t) = \frac{\partial f}{\partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x + \frac{\partial f}{\partial y}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta y$$

$$F''(t) = \frac{\partial^2 f}{\partial x^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta x)^2 + 2\frac{\partial^2 f}{\partial y \partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x \Delta y \\ + \frac{\partial^2 f}{\partial y^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta y)^2 = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0 + t\Delta x, y_0 + t\Delta y);$$

.....

$$F^{(n)}(t) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0 + t\Delta x, y_0 + t\Delta y)$$

$$\text{其中, } (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n = \sum_{k=0}^n C_n^k (\Delta x)^k (\Delta y)^{n-k} \frac{\partial^k}{\partial x^k} \frac{\partial^{n-k}}{\partial y^{n-k}}$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F'(t) = \frac{\partial f}{\partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x + \frac{\partial f}{\partial y}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta y$$

$$F''(t) = \frac{\partial^2 f}{\partial x^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta x)^2 + 2\frac{\partial^2 f}{\partial y \partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x \Delta y \\ + \frac{\partial^2 f}{\partial y^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta y)^2 = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0 + t\Delta x, y_0 + t\Delta y);$$

.....

$$F^{(n)}(t) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0 + t\Delta x, y_0 + t\Delta y)$$

$$\Rightarrow F^{(n)}(0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0)$$

$$F^{(n+1)}(\theta) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F'(t) = \frac{\partial f}{\partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x + \frac{\partial f}{\partial y}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta y$$

$$F''(t) = \frac{\partial^2 f}{\partial x^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta x)^2 + 2\frac{\partial^2 f}{\partial y \partial x}(x_0 + t\Delta x, y_0 + t\Delta y)\Delta x \Delta y \\ + \frac{\partial^2 f}{\partial y^2}(x_0 + t\Delta x, y_0 + t\Delta y)(\Delta y)^2 = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0 + t\Delta x, y_0 + t\Delta y);$$

.....

$$F^{(n)}(t) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0 + t\Delta x, y_0 + t\Delta y)$$

$$\Rightarrow F^{(n)}(0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0)$$

$$F^{(n+1)}(\theta) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$\Rightarrow F(1) = F(0) + F'(0) + \frac{F''(0)}{2!} + \cdots + \frac{F^{(n)}(0)}{n!} + \frac{F^{(n+1)}(\theta)}{(n+1)!}$$

根据一元函数Taylor公式,现有

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2!}t^2 + \cdots + \frac{F^{(n)}(0)}{n!}t^n + \frac{F^{(n+1)}(\theta)}{(n+1)!}t^{n+1}, \theta \in (0, t).$$

$$\Delta z = f(x, y) - f(x_0, y_0) = F(1) - F(0)$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F^{(n)}(0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0)$$

$$F^{(n+1)}(\theta) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$F(1) = F(0) + F'(0) + \frac{F''(0)}{2!} + \cdots + \frac{F^{(n)}(0)}{n!} + \frac{F^{(n+1)}(\theta)}{(n+1)!}$$

$$\Delta z = f(x, y) - f(x_0, y_0) = F(1) - F(0)$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F^{(n)}(0) = \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right)^n f(x_0, y_0)$$

$$F^{(n+1)}(\theta) = \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right)^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$F(1) = F(0) + F'(0) + \frac{F''(0)}{2!} + \cdots + \frac{F^{(n)}(0)}{n!} + \frac{F^{(n+1)}(\theta)}{(n+1)!}$$

$$f(x, y) - f(x_0, y_0) = F(1) - F(0)$$

$$= \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right) f(x_0, y_0) + \frac{1}{2!} \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right)^2 f(x_0, y_0) + \cdots$$

$$+ \frac{1}{n!} \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right)^n f(x_0, y_0) + \frac{1}{(n+1)!} \left(\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}\right)^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$\Delta z = f(x, y) - f(x_0, y_0) = F(1) - F(0)$$

$$F(t) = f(x_0 + t\Delta x, y_0 + t\Delta y), t \in [0, 1]$$

$$F^{(n)}(0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0)$$

$$F^{(n+1)}(\theta) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$F(1) = F(0) + F'(0) + \frac{F''(0)}{2!} + \cdots + \frac{F^{(n)}(0)}{n!} + \frac{F^{(n+1)}(\theta)}{(n+1)!}$$

$$f(x, y) - f(x_0, y_0) = F(1) - F(0)$$

$$= (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

$$+ \frac{1}{n!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0) + \frac{1}{(n+1)!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

$$f(x, y) = f(x_0, y_0) + (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

$$+ \frac{1}{n!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^n f(x_0, y_0) + \frac{1}{(n+1)!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^{n+1} f(x_0 + \theta\Delta x, y_0 + \theta\Delta y)$$

结论

若 $z = f(x, y)$ 于 (x_0, y_0) 点取到极值, 且 $\frac{\partial f}{\partial x}(x_0, y_0), \frac{\partial f}{\partial y}(x_0, y_0)$ 存在, 则


$$\frac{\partial f}{\partial x}(x_0, y_0) = 0, \frac{\partial f}{\partial y}(x_0, y_0) = 0.$$



结论

若 $z = f(x, y)$ 于 (x_0, y_0) 点取到极值, 且 $\frac{\partial f}{\partial x}(x_0, y_0), \frac{\partial f}{\partial y}(x_0, y_0)$ 存在, 则

$$\frac{\partial f}{\partial x}(x_0, y_0) = 0, \frac{\partial f}{\partial y}(x_0, y_0) = 0.$$



定义

使 $\frac{\partial f}{\partial x}(x_0, y_0), \frac{\partial f}{\partial y}(x_0, y_0)$ 为零的点, 称为 $z = f(x, y)$ 的驻点

极值可疑点

先考察二元函数 $z = Ax^2 + 2Bxy + Cy^2$ 的性态, 其中 A, B, C 为常数.

先考察二元函数 $z = Ax^2 + 2Bxy + Cy^2$ 的性态, 其中 A, B, C 为常数.

设 $A \neq 0$, 则 $z = A\left[\left(x + \frac{B}{A}y\right)^2 + \frac{AC - B^2}{A^2}y^2\right]$

先考察二元函数 $z = Ax^2 + 2Bxy + Cy^2$ 的性态, 其中 A, B, C 为常数.

设 $A \neq 0$, 则 $z = A[(x + \frac{B}{A}y)^2 + \frac{AC - B^2}{A^2}y^2]$

若 $AC - B^2 > 0$, 则函数值 z (除在 $(0, 0)$ 点外)

要么全为正 ($A > 0$), 为开口向上的抛物面, 全在 xoy 平面上方;

要么全为负 ($A < 0$), 为开口向下的抛物面, 全在 xoy 平面下方;

即 $AC - B^2 > 0$ 时, 函数 $z = Ax^2 + 2Bxy + Cy^2$

(除在 $(0, 0)$ 点外) 的值有确定的符号 与 A 同号

先考察二元函数 $z = Ax^2 + 2Bxy + Cy^2$ 的性态, 其中 A, B, C 为常数.

设 $A \neq 0$, 则 $z = A[(x + \frac{B}{A}y)^2 + \frac{AC - B^2}{A^2}y^2]$

若 $AC - B^2 > 0$, 则函数值 z (除在 $(0, 0)$ 点外)

要么全为正 ($A > 0$), 为开口向上的抛物面, 全在 xoy 平面上方;

要么全为负 ($A < 0$), 为开口向下的抛物面, 全在 xoy 平面下方;

即 $AC - B^2 > 0$ 时, 函数 $z = Ax^2 + 2Bxy + Cy^2$

(除在 $(0, 0)$ 点外) 的值有确定的符号 与 A 同号

若 $AC - B^2 < 0$, 则函数值 z 对不同的 x, y 有正有负, 比如 $y = 0$ 时, $z = Ax^2$ 与 A 同号,
 $x = -\frac{B}{A}y$ 时, $z = \frac{AC - B^2}{A}y^2$ 与 A 异号. 所以, $AC - B^2 < 0$ 时, 函数 $z = Ax^2 + 2Bxy + Cy^2$ 的
值(在 origin 附近)不具有确定的符号.

先考察二元函数 $z = Ax^2 + 2Bxy + Cy^2$ 的性态, 其中 A, B, C 为常数.

事实上, 如果学过线性代数, 就知道:

$A > 0, AC - B^2 > 0$ 时, 二次型 $Ax^2 + 2Bxy + Cy^2$ 正定;

$A < 0, AC - B^2 > 0$ 时, 二次型 $Ax^2 + 2Bxy + Cy^2$ 负定;

$AC - B^2 < 0$ 时, 二次型 $Ax^2 + 2Bxy + Cy^2$ 不定.

设 $z = f(x, y)$, 在 (x_0, y_0) 点, 各阶偏导数连续, 则有二元函数 Taylor 公式:

$$f(x, y) - f(x_0, y_0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

设 $z = f(x, y)$, 在 (x_0, y_0) 点, 各阶偏导数连续, 则有二元函数 Taylor 公式:

$$f(x, y) - f(x_0, y_0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

If $\frac{\partial}{\partial x} f(x_0, y_0), \frac{\partial}{\partial y} f(x_0, y_0)$ 不都为零, 则 $f(x_0, y_0)$ 非极值;

设 $z = f(x, y)$, 在 (x_0, y_0) 点, 各阶偏导数连续, 则有二元函数 Taylor 公式:

$$f(x, y) - f(x_0, y_0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

If $\frac{\partial}{\partial x} f(x_0, y_0), \frac{\partial}{\partial y} f(x_0, y_0)$ 不都为零, 则 $f(x_0, y_0)$ 非极值;

If $\frac{\partial}{\partial x} f(x_0, y_0) = \frac{\partial}{\partial y} f(x_0, y_0) = 0$

考察 $\Phi(\Delta x, \Delta y) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0)$ 在 $\forall \Delta x, \Delta y$ 变化时定号否.

定号则 $f(x_0, y_0)$ 为极值, 否则 $f(x_0, y_0)$ 非极值.

设 $z = f(x, y)$, 在 (x_0, y_0) 点, 各阶偏导数连续, 则有二元函数 Taylor 公式:

$$f(x, y) - f(x_0, y_0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

If $\frac{\partial}{\partial x} f(x_0, y_0), \frac{\partial}{\partial y} f(x_0, y_0)$ 不都为零, 则 $f(x_0, y_0)$ 非极值;

$$\text{If } \frac{\partial}{\partial x} f(x_0, y_0) = \frac{\partial}{\partial y} f(x_0, y_0) = 0$$

考察 $\Phi(\Delta x, \Delta y) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0)$ 在 $\forall \Delta x, \Delta y$ 变化时定号否.

定号则 $f(x_0, y_0)$ 为极值, 否则 $f(x_0, y_0)$ 非极值.

$$\text{即 } \Phi = (\Delta x)^2 \frac{\partial^2 f(x_0, y_0)}{\partial x^2} + 2\Delta x \Delta y \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y} + (\Delta y)^2 \frac{\partial^2 f(x_0, y_0)}{\partial y^2}$$

是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$, 在 (x_0, y_0) 点, 各阶偏导数连续, 则有二元函数 Taylor 公式:

$$f(x, y) - f(x_0, y_0) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y}) f(x_0, y_0) + \frac{1}{2!} (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0) + \cdots$$

If $\frac{\partial}{\partial x} f(x_0, y_0), \frac{\partial}{\partial y} f(x_0, y_0)$ 不都为零, 则 $f(x_0, y_0)$ 非极值;

If $\frac{\partial}{\partial x} f(x_0, y_0) = \frac{\partial}{\partial y} f(x_0, y_0) = 0$

考察 $\Phi(\Delta x, \Delta y) = (\Delta x \frac{\partial}{\partial x} + \Delta y \frac{\partial}{\partial y})^2 f(x_0, y_0)$ 在 $\forall \Delta x, \Delta y$ 变化时定号否.

定号则 $f(x_0, y_0)$ 为极值, 否则 $f(x_0, y_0)$ 非极值.

$$\text{即 } \Phi = (\Delta x)^2 \frac{\partial^2 f(x_0, y_0)}{\partial x^2} + 2\Delta x \Delta y \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y} + (\Delta y)^2 \frac{\partial^2 f(x_0, y_0)}{\partial y^2}$$

是否关于 $\forall \Delta x, \Delta y$ 定号.

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$ 在某区域内有二阶连续偏导数, 求出函数 $z = f(x, y)$ 在此区域内的驻点 (x_0, y_0) , 在驻点处求出 $A = \frac{\partial^2 f}{\partial x^2}$, $B = \frac{\partial^2 f}{\partial x \partial y}$, $C = \frac{\partial^2 f}{\partial y^2}$ 的值, 判断 $AC - B^2$ 是否大于 0.

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$ 在某区域内有二阶连续偏导数, 求出函数 $z = f(x, y)$ 在此区域内的驻点 (x_0, y_0) , 在驻点处求出 $A = \frac{\partial^2 f}{\partial x^2}$, $B = \frac{\partial^2 f}{\partial x \partial y}$, $C = \frac{\partial^2 f}{\partial y^2}$ 的值, 判断 $AC - B^2$ 是否大于 0.

若 $AC - B^2 < 0$, 则该驻点非极值点; 若 $AC - B^2 > 0$, 则该驻点为极值点, 此时

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$ 在某区域内有二阶连续偏导数, 求出函数 $z = f(x, y)$ 在此区域内的驻点 (x_0, y_0) , 在驻点处求出 $A = \frac{\partial^2 f}{\partial x^2}$, $B = \frac{\partial^2 f}{\partial x \partial y}$, $C = \frac{\partial^2 f}{\partial y^2}$ 的值, 判断 $AC - B^2$ 是否大于 0.

若 $AC - B^2 < 0$, 则该驻点非极值点; 若 $AC - B^2 > 0$, 则该驻点为极值点, 此时

$$\begin{cases} A > 0 \Rightarrow f(x_0, y_0) \text{ 是极小值} \\ A < 0 \Rightarrow f(x_0, y_0) \text{ 是极大值} \end{cases}$$

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$ 在某区域内有二阶连续偏导数, 求出函数 $z = f(x, y)$ 在此区域内的驻点 (x_0, y_0) , 在驻点处求出 $A = \frac{\partial^2 f}{\partial x^2}$, $B = \frac{\partial^2 f}{\partial x \partial y}$, $C = \frac{\partial^2 f}{\partial y^2}$ 的值, 判断 $AC - B^2$ 是否大于 0.

若 $AC - B^2 < 0$, 则该驻点非极值点; 若 $AC - B^2 > 0$, 则该驻点为极值点, 此时

$$\begin{cases} A > 0 \Rightarrow f(x_0, y_0) \text{ 是极小值} \\ A < 0 \Rightarrow f(x_0, y_0) \text{ 是极大值} \end{cases}$$

若 $AC - B^2 = 0$, 情况待定.

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

设 $z = f(x, y)$ 在某区域内有二阶连续偏导数, 求出函数 $z = f(x, y)$ 在此区域内的驻点 (x_0, y_0) , 在驻点处求出 $A = \frac{\partial^2 f}{\partial x^2}, B = \frac{\partial^2 f}{\partial x \partial y}, C = \frac{\partial^2 f}{\partial y^2}$ 的值, 判断 $AC - B^2$ 是否大于 0.

若 $AC - B^2 < 0$, 则该驻点非极值点; 若 $AC - B^2 > 0$, 则该驻点为极值点, 此时

$$\begin{cases} A > 0 \Rightarrow f(x_0, y_0) \text{ 是极小值} \\ A < 0 \Rightarrow f(x_0, y_0) \text{ 是极大值} \end{cases}$$

若 $AC - B^2 = 0$, 情况待定.

如 $z = x^4 + y^4 - x^2 - 2xy - y^2$ 有三个驻点 $(-1, -1), (0, 0), (1, 1)$

在 $(0, 0)$ 处 $A = -2, B = -2, C = -2, AC - B^2 = 0$, 此时无法去决定 $(0, 0)$ 是否是极值点.

但在 $y = x$ 上, $z = x^4 - 4x^2 = x^2(x^2 - 4) < 0 (0 < |x| \leq 1)$, 在 $y = -x$ 上, $z = 2x^4 > 0 (0 < |x| \leq 1)$, 所以 $(0, 0)$ 不是极值点.

$$\text{令 } A = \frac{\partial^2 f(x_0, y_0)}{\partial x^2}, B = \frac{\partial^2 f(x_0, y_0)}{\partial x \partial y}, C = \frac{\partial^2 f(x_0, y_0)}{\partial y^2},$$

即考察 $A(\Delta x)^2 + 2B\Delta x\Delta y + C(\Delta y)^2$ 是否关于 $\forall \Delta x, \Delta y$ 定号.

例 求 $z = 3xy - x^3 - y^3$ 的极值.

例 求 $z = 3xy - x^3 - y^3$ 的极值.

解: $\begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}$, 令 $\begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}$, 则 $\begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}$, $\begin{cases} y = x^2 \\ x = y^2 \end{cases}$,

例 求 $z = 3xy - x^3 - y^3$ 的极值.

解: $\begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}$, 令 $\begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}$, 则 $\begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}$, $\begin{cases} y = x^2 \\ x = y^2 \end{cases}$,

得驻点 $(x, y) = (0, 0), (x, y) = (1, 1)$

例 求 $z = 3xy - x^3 - y^3$ 的极值.

解: $\begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}$, 令 $\begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}$, 则 $\begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}$, $\begin{cases} y = x^2 \\ x = y^2 \end{cases}$,

得驻点 $(x, y) = (0, 0), (x, y) = (1, 1)$

$$\text{又 } \frac{\partial^2 z}{\partial x^2} = -6x, \frac{\partial^2 z}{\partial x \partial y} = 3, \frac{\partial^2 z}{\partial y^2} = -6y^2$$

例 求 $z = 3xy - x^3 - y^3$ 的极值.

$$\text{解: } \begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}, \text{ 令 } \begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}, \text{ 则 } \begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}, \begin{cases} y = x^2 \\ x = y^2 \end{cases},$$

得驻点 $(x, y) = (0, 0), (x, y) = (1, 1)$

$$\text{又 } \frac{\partial^2 z}{\partial x^2} = -6x, \frac{\partial^2 z}{\partial x \partial y} = 3, \frac{\partial^2 z}{\partial y^2} = -6y^2$$

\therefore 于 $(0, 0)$ 点, $A = 0, B = 3, C = 0, AC - B^2 = -9 < 0, \therefore (0, 0)$ 非极值点.

例 求 $z = 3xy - x^3 - y^3$ 的极值.

$$\text{解: } \begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}, \text{ 令 } \begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}, \text{ 则 } \begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}, \begin{cases} y = x^2 \\ x = y^2 \end{cases},$$

得驻点 $(x, y) = (0, 0), (x, y) = (1, 1)$

$$\text{又 } \frac{\partial^2 z}{\partial x^2} = -6x, \frac{\partial^2 z}{\partial x \partial y} = 3, \frac{\partial^2 z}{\partial y^2} = -6y^2$$

\therefore 于 $(0, 0)$ 点, $A = 0, B = 3, C = 0, AC - B^2 = -9 < 0, \therefore (0, 0)$ 非极值点.

\therefore 于 $(1, 1)$ 点, $A = -6, B = 3, C = -6, AC - B^2 = 27 > 0, \therefore (1, 1)$ 为极值点.

例 求 $z = 3xy - x^3 - y^3$ 的极值.

$$\text{解: } \begin{cases} \frac{\partial z}{\partial x} = 3y - 3x^2 \\ \frac{\partial z}{\partial y} = 3x - 3y^2 \end{cases}, \text{ 令 } \begin{cases} \frac{\partial z}{\partial x} = 0 \\ \frac{\partial z}{\partial y} = 0 \end{cases}, \text{ 则 } \begin{cases} 3y - 3x^2 = 0 \\ 3x - 3y^2 = 0 \end{cases}, \begin{cases} y = x^2 \\ x = y^2 \end{cases},$$

得驻点 $(x, y) = (0, 0), (x, y) = (1, 1)$

$$\text{又 } \frac{\partial^2 z}{\partial x^2} = -6x, \frac{\partial^2 z}{\partial x \partial y} = 3, \frac{\partial^2 z}{\partial y^2} = -6y^2$$

\therefore 于 $(0, 0)$ 点, $A = 0, B = 3, C = 0, AC - B^2 = -9 < 0, \therefore (0, 0)$ 非极值点.

\therefore 于 $(1, 1)$ 点, $A = -6, B = 3, C = -6, AC - B^2 = 27 > 0, \therefore (1, 1)$ 为极值点.

$\because A = -6 < 0 \therefore (1, 1)$ 为极大值点, 极大值为 $f(1, 1) = 3 - 1 - 1 = 1$.

