Microbial Group Evolution MCG Theory
Complete Unified Framework

Author: Michael Levin | Version: 1.0 | Date: 2026-02-14

Overview

Microbial Group Evolution (MCG) Theory is a unified framework for understanding how microbial communities evolve, organized around three interconnected dimensions: the ecological structure of communities, the genetic architecture of adaptation, and the evolutionary dynamics that shape both over time.

Scope: Applicable to bacterial, archaeal, and microbial eukaryotic communities in natural and laboratory settings.

Status: Theoretical framework with empirical validation ongoing.

Key Innovation: Integration of ecological, genetic, and evolutionary perspectives into a single mathematical framework.

---

1. Ecological Structure

Core Concepts

Community Structure: Microbial communities are characterized by species composition, abundance distributions, and spatial organization. Diversity emerges from competition, cooperation, and environmental heterogeneity.

Ecological Niches: Niches are defined by resource utilization, tolerance to environmental conditions, and interactions with other species. Niche boundaries determine which species can persist in a community.

Stability: Communities can exhibit multiple stable states. Transitions between states occur when environmental conditions cross critical thresholds.

Succession: Communities change over time in predictable patterns as species modify their environment and alter the conditions for subsequent colonizers.

1.1 Shannon Entropy and Diversity

Shannon Entropy:

$$H = -\sum_{i=1}^{S} p_i \ln(p_i)$$

Where:
- H = Shannon entropy (bits)
- S = Total number of species
- p_i = Proportion of species i

Shannon Diversity:

$$D = e^{H}$$

Interpretation:
- D = 1: Monoculture (one species dominates)
- D = 10: Moderate diversity (~10 effective species)
- D = 100: High diversity (~100 effective species)

Pielou's Evenness:

$$J = \frac{H}{\ln(S)}$$

Where:
- J = Evenness (0 to 1)
- H = Shannon entropy
- S = Total number of species

Interpretation:
- J = 1: Perfect evenness (all species equally abundant)
- J = 0.5: Moderate evenness
- J = 0: Maximum dominance (one species dominates)

Relationship: $D = e^{H}$ and $J = H / \ln(S)$

Implications:
- Diversity measures community complexity
- Evenness measures how equally resources are distributed
- Low evenness indicates competitive exclusion or resource limitation
- Shannon entropy quantifies information content of community structure

1.2 Lotka-Volterra Dynamics

Lotka-Volterra Equations:

$$\frac{dx_i}{dt} = r_i x_i \left(1 - \sum_j \alpha_{ij} x_j\right)$$

Where:
- x_i = Population of species i
- r_i = Intrinsic growth rate of species i
- α_{ij} = Interaction coefficient (effect of species j on species i)

Interpretation:
- α_{ii} = Self-limitation (intraspecific competition)
- $\alpha_{ij} > 0$ = Species j inhibits species i
- $\alpha_{ij} < 0$ = Species j facilitates species i

Applications:
- Predicting community composition
- Identifying keystone species
- Modeling invasion dynamics
- Predicting tipping points

1.3 Hill Numbers Family

Hill Numbers:

$$^{q}D = \left(\sum_{i=1}^{S} p_i^{q}\right)^{1/(1-q)}$$

Where:
- ^{q}D = Hill number of order q
- q = Sensitivity parameter
- p_i = Proportion of species i

Special Cases:
- ^{0}D = Richness (all species equally weighted)
- ^{1}D = e^{H} (Shannon diversity)
- ^{2}D = Simpson diversity (dominance-weighted)

Interpretation:
- Higher q values emphasize dominant species
- Lower q values emphasize rare species
- Hill numbers provide a family of diversity measures with different sensitivities

Implications:
- Multiple diversity measures capture different aspects of community structure
- Choice of q depends on ecological question
- Hill numbers are mathematically consistent across scales

1.4 Cross-Feeding Networks

Cross-Feeding Matrix:

$$M_{ij} = \text{Metabolite } j \text{ produced by species } i \text{ consumed by species } j$$

Network Properties:
- Connectivity: Number of cross-feeding interactions
- Modularity: Degree to which the network divides into subgroups
- Centrality: Which species are most connected
- Robustness: How many species can be lost before network collapse

Implications:
- Cross-feeding creates interdependence between species
- Metabolite exchange enables coexistence
- Network structure predicts community stability
- Disruption of cross-feeding can destabilize communities

1.5 Resource Competition

Resource Dynamics:

$$\frac{dR_k}{dt} = \text{Supply} - \text{Consumption}$$

Monod Kinetics:

$$\mu = \mu_{max} \frac{R}{K + R}$$

Where:
- μ = Growth rate
- μ_{max} = Maximum growth rate
- R = Resource concentration
- K = Half-saturation constant

Competition:
- Species with lower K values can persist at lower resource levels
- Species with higher μ_{max} values can grow faster when resources are abundant
- Trade-off between growth rate and resource affinity enables coexistence

Implications:
- Resource competition determines which species can persist
- Resource ratios determine community composition
- Resource fluctuations can destabilize communities

1.6 Priority Effects

Priority Effects:

$$\text{Outcome} = f(\text{Arrival order}, \text{Initial conditions})$$

Mechanisms:
- Niche modification: Early colonizers change environment for later arrivals
- Resource depletion: Early colonizers deplete resources needed by others
- Facilitation: Early colonizers create conditions favorable to others
- Inhibition: Early colonizers prevent others from establishing

Implications:
- Historical contingency matters
- Identical communities can diverge based on arrival order
- Predicting community composition requires knowing assembly history
- Priority effects can lock communities into alternative states

1.7 Beta Diversity Dynamics

Beta Diversity:

$$\beta = \text{Variance in community composition across sites}$$

Patterns:
- Distance-decay: Similarity decreases with distance
- Nestedness: Species-poor communities are subsets of species-rich communities
- Turnover: Species replacement without richness change

Implications:
- Spatial heterogeneity maintains regional diversity
- Dispersal limitation creates distance-decay patterns
- Habitat fragmentation disrupts nestedness patterns

1.8 Ecological Succession

Succession:

$$\text{Community}(t) = f(\text{Community}(t-1), \text{Environmental change})$$

Patterns:
- Early colonizers: Fast growth, low competitive ability
- Late colonizers: Slow growth, high competitive ability
- Climax community: Stable composition after succession completes

Implications:
- Communities change predictably over time
- Succession rates depend on environmental conditions
- Disturbance resets succession to earlier stages

---

2. Genetic Architecture

Core Concepts

Genetic Architecture: The genetic architecture of adaptation determines how many genes are involved, their effect sizes, and their interactions. Polygenic adaptation involves many genes with small effects, while monogenic adaptation involves few genes with large effects.

Adaptation Speed: Adaptation speed depends on the genetic variation available, the strength of selection, and the population size. Polygenic adaptation can be slow when selection is weak relative to genetic drift.

Trade-offs: Genetic trade-offs between traits can constrain adaptation. Pleiotropy creates correlated responses to selection.

Epistasis: Gene interactions can modify the effects of mutations and create complex fitness landscapes.

2.1 Polygenic Adaptation

Polygenic Adaptation:

$$\Delta\bar{z} = \sum_{i=1}^{n} \alpha_i \Delta p_i$$

Where:
- $\Delta\bar{z}$ = Change in mean phenotype
- n = Number of loci
- α_i = Effect size of locus i
- Δp_i = Change in allele frequency at locus i

Conditions:
- Many loci with small effects
- Weak selection relative to genetic drift
- Sufficient genetic variation
- Large population size

Implications:
- Adaptation can be slow when selection is weak
- Genetic drift can overwhelm selection at many loci
- Linkage disequilibrium can accelerate adaptation
- Clonal interference can slow adaptation in asexual populations

2.2 Fisher's Geometric Model

Fisher's Geometric Model:

$$P(\text{Adaptive mutation}) = \Phi\left(\frac{s\sqrt{n}}{2}\right)$$

Where:
- P = Probability of adaptive mutation
- Φ = Cumulative normal distribution function
- s = Selection coefficient
- n = Number of dimensions

Implications:
- Probability of adaptive mutation decreases with distance from optimum
- Probability of adaptive mutation decreases with dimensionality
- Small mutations are more likely to be adaptive than large mutations
- Adaptation slows as population approaches optimum

2.3 Clonal Interference

Clonal Interference:

$$\text{Adaptation rate} = f(\text{Mutation rate}, \text{Selection coefficient}, \text{Population size})$$

Mechanism:
- Multiple beneficial mutations arise in different lineages
- Lineages compete with each other
- Only one lineage can fix the beneficial mutation
- Other beneficial mutations are lost

Implications:
- Clonal interference slows adaptation in asexual populations
- Recombination can overcome clonal interference
- Population size determines how many lineages can compete
- Clonal interference creates selection for mutation rate modifiers

2.4 Mutational Pleiotropy

Mutational Pleiotropy:

$$\text{Correlated response} = \rho_{mu} \cdot \text{Selection response}$$

Where:
- ρ_{mu} = Mutational correlation between traits

Implications:
- Pleiotropy creates correlated responses to selection
- Positive correlation facilitates adaptation
- Negative correlation constrains adaptation
- Pleiotropy can create trade-offs between traits

2.5 Epistasis Fitness Landscapes

Epistasis:

$$\text{Fitness}(A, B) \neq \text{Fitness}(A) + \text{Fitness}(B) - \text{Fitness}(\text{Wild type})$$

Types:
- Positive epistasis: Mutations are more beneficial together than expected
- Negative epistasis: Mutations are less beneficial together than expected
- Sign epistasis: Mutation changes from beneficial to deleterious or vice versa

Implications:
- Epistasis creates rugged fitness landscapes
- Epistasis can make adaptation path-dependent
- Epistasis can create multiple adaptive peaks
- Epistasis can constrain adaptation by creating valleys between peaks

2.6 Adaptation Speed Limits

Adaptation Speed:

$$\text{Adaptation rate} = f(\text{Genetic variation}, \text{Selection strength}, \text{Population size})$$

Limiting Factors:
- Genetic variation: Mutation rate and standing variation
- Selection strength: Selection coefficient relative to genetic drift
- Population size: Number of individuals available for mutation
- Linkage disequilibrium: Recombination breaks up beneficial combinations
- Clonal interference: Competition between beneficial mutations
- Epistasis: Gene interactions modify fitness effects

Implications:
- Adaptation can be slow when any factor is limiting
- Multiple factors can interact to slow adaptation
- Overcoming one limiting factor may reveal another
- Adaptation speed varies across genes and traits

---

3. Evolutionary Dynamics

Core Concepts

Evolutionary Dynamics: Evolutionary dynamics describe how populations change over time in response to selection, mutation, drift, and migration. These forces interact to determine which genotypes persist and spread.

Selection: Selection favors genotypes with higher fitness. The strength of selection determines how quickly adaptive mutations spread.

Drift: Genetic drift is random change in allele frequencies. Drift can overwhelm selection when selection is weak or population size is small.

Mutation: Mutation creates genetic variation. Mutation rate determines how quickly new variation arises.

Migration: Migration introduces genetic variation from other populations. Migration can prevent local adaptation.

3.1 Fisher's Fundamental Theorem

Fisher's Fundamental Theorem:

$$\frac{d\bar{w}}{dt} = V_A$$

Where:
- $\bar{w}$ = Mean fitness
- V_A = Additive genetic variance

Implications:
- Mean fitness increases at rate proportional to additive genetic variance
- Adaptation slows as population approaches optimum
- Genetic variation is required for adaptation
- Selection depletes genetic variation

3.2 Price Equation

Price Equation:

$$\Delta\bar{z} = \frac{\text{Cov}(w, z)}{\bar{w}} + \frac{E(w \Delta z)}{\bar{w}}$$

Where:
- $\Delta\bar{z}$ = Change in mean phenotype
- w = Fitness
- z = Phenotype
- $\text{Cov}(w, z)$ = Selection component
- $E(w \Delta z)$ = Transmission component

Interpretation:
- Selection component: Selection favors individuals with higher phenotype
- Transmission component: Mutation changes phenotype

Implications:
- Selection component drives adaptation
- Transmission component can oppose adaptation
- Price equation partitions evolutionary change into selection and transmission components

3.3 Wright-Fisher Model

Wright-Fisher Model:

$$P(k \text{ copies}) = \binom{N}{k} p^k (1-p)^{N-k}$$

Where:
- N = Population size
- p = Allele frequency
- k = Number of copies

Implications:
- Drift is random change in allele frequencies
- Drift is strongest when population size is small
- Drift can fix alleles even when selection favors other alleles
- Drift reduces genetic variation over time

3.4 Ecological Feedback Dynamics

Ecological Feedback:

$$\text{Evolutionary change} \rightarrow \text{Ecological change} \rightarrow \text{Evolutionary change}$$

Mechanisms:
- Niche construction: Organisms modify their environment
- Resource depletion: Consumers deplete resources
- Toxin production: Organisms produce toxins that affect others
- Facilitation: Organisms create conditions favorable to others

Implications:
- Evolutionary change can create ecological change
- Ecological change can create evolutionary change
- Feedbacks can create oscillations
- Feedbacks can create multiple stable states

3.5 Frequency-Dependent Selection

Frequency-Dependent Selection:

$$\text{Fitness}(A) = f(\text{Frequency of } A)$$

Types:
- Negative frequency-dependence: Rare alleles have higher fitness
- Positive frequency-dependence: Common alleles have higher fitness

Implications:
- Negative frequency-dependence maintains polymorphism
- Positive frequency-dependence creates bistability
- Frequency-dependence can create oscillations
- Frequency-dependence can create multiple equilibria

3.6 Adaptive Dynamics

Adaptive Dynamics:

$$\frac{dx}{dt} = m \cdot s \cdot \frac{\partial w}{\partial x}$$

Where:
- x = Trait value
- m = Mutation rate
- s = Selection coefficient
- $\frac{\partial w}{\partial x}$ = Fitness gradient

Implications:
- Adaptation follows the fitness gradient
- Adaptation stops at fitness peaks
- Adaptation can be slow when fitness gradient is small
- Adaptation can be fast when fitness gradient is large

---

4. Integration

Cross-Domain Connections

Ecological → Genetic:
- Niche diversity → Genetic diversity
- Resource competition → Selection for efficiency genes
- Cross-feeding → Selection for cooperation genes
- Priority effects → Historical contingency in genetic evolution

Genetic → Ecological:
- Genetic diversity → Ecological diversity
- Adaptation → Niche expansion
- Genetic trade-offs → Ecological constraints
- Epistasis → Ecological tipping points

Evolutionary → Ecological:
- Selection → Community composition
- Drift → Random community assembly
- Mutation → Source of ecological variation
- Migration → Spatial heterogeneity

Ecological → Evolutionary:
- Ecological succession → Changing selection pressures
- Ecological feedbacks → Evolutionary feedbacks
- Ecological stability → Evolutionary stability
- Ecological tipping points → Evolutionary tipping points

Unified Principles

Diversity:
- Ecological diversity: Species richness, evenness, Hill numbers
- Genetic diversity: Polygenic adaptation, clonal interference, epistasis
- Evolutionary diversity: Selection, drift, mutation, migration

Stability:
- Ecological stability: Lotka-Volterra, cross-feeding networks, priority effects
- Genetic stability: Fisher's geometric model, epistasis, adaptation speed limits
- Evolutionary stability: Fisher's fundamental theorem, Price equation, Wright-Fisher model

Adaptation:
- Ecological adaptation: Niche construction, resource competition, succession
- Genetic adaptation: Polygenic adaptation, clonal interference, epistasis
- Evolutionary adaptation: Selection, drift, mutation, migration

Constraints:
- Ecological constraints: Resource limitation, competition, priority effects
- Genetic constraints: Genetic variation, trade-offs, epistasis
- Evolutionary constraints: Selection strength, population size, mutation rate

Feedbacks:
- Ecological feedbacks: Cross-feeding, niche construction, succession
- Genetic feedbacks: Epistasis, pleiotropy, adaptation speed
- Evolutionary feedbacks: Frequency-dependence, adaptive dynamics, ecological feedbacks

Cross-Domain Predictions

Ecological → Genetic:
- Niche diversity predicts genetic diversity
- Resource competition predicts selection for efficiency genes
- Cross-feeding predicts selection for cooperation genes
- Priority effects predict historical contingency in genetic evolution

Genetic → Ecological:
- Genetic diversity predicts ecological diversity
- Adaptation predicts niche expansion
- Genetic trade-offs predict ecological constraints
- Epistasis predicts ecological tipping points

Evolutionary → Ecological:
- Selection predicts community composition
- Drift predicts random community assembly
- Mutation predicts source of ecological variation
- Migration predicts spatial heterogeneity

Ecological → Evolutionary:
- Ecological succession predicts changing selection pressures
- Ecological feedbacks predict evolutionary feedbacks
- Ecological stability predicts evolutionary stability
- Ecological tipping points predict evolutionary tipping points

Key Mechanisms

Niche Construction:
- Organisms modify their environment
- Modified environment changes selection pressures
- Selection favors genotypes that maintain favorable conditions
- Niche construction creates feedbacks between ecology and evolution

Cross-Feeding:
- Metabolite exchange between species
- Creates interdependence between species
- Enables coexistence
- Network structure predicts community stability

Priority Effects:
- Early colonizers modify environment for later arrivals
- Resource depletion by early colonizers limits later arrivals
- Facilitation by early colonizers enables later arrivals
- Priority effects create historical contingency

Epistasis:
- Gene interactions modify fitness effects
- Creates rugged fitness landscapes
- Makes adaptation path-dependent
- Creates multiple adaptive peaks

Clonal Interference:
- Multiple beneficial mutations arise in different lineages
- Lineages compete with each other
- Only one lineage can fix the beneficial mutation
- Other beneficial mutations are lost

Frequency-Dependence:
- Fitness depends on frequency of other genotypes
- Negative frequency-dependence maintains polymorphism
- Positive frequency-dependence creates bistability
- Frequency-dependence can create oscillations

Adaptive Dynamics:
- Adaptation follows the fitness gradient
- Adaptation stops at fitness peaks
- Adaptation can be slow when fitness gradient is small
- Adaptation can be fast when fitness gradient is large

Mathematical Relationships

Diversity Measures:
- Shannon entropy: $H = -\sum p_i \ln(p_i)$
- Shannon diversity: $D = e^{H}$
- Pielou's evenness: $J = H / \ln(S)$
- Hill numbers: $^{q}D = \left(\sum p_i^{q}\right)^{1/(1-q)}$

Ecological Dynamics:
- Lotka-Volterra: $\frac{dx_i}{dt} = r_i x_i \left(1 - \sum_j \alpha_{ij} x_j\right)$
- Monod kinetics: $\mu = \mu_{max} \frac{R}{K + R}$
- Resource dynamics: $\frac{dR_k}{dt} = \text{Supply} - \text{Consumption}$

Genetic Dynamics:
- Polygenic adaptation: $\Delta\bar{z} = \sum \alpha_i \Delta p_i$
- Fisher's geometric model: $P(\text{Adaptive mutation}) = \Phi\left(\frac{s\sqrt{n}}{2}\right)$
- Epistasis: $\text{Fitness}(A, B) \neq \text{Fitness}(A) + \text{Fitness}(B) - \text{Fitness}(\text{Wild type})$

Evolutionary Dynamics:
- Fisher's fundamental theorem: $\frac{d\bar{w}}{dt} = V_A$
- Price equation: $\Delta\bar{z} = \frac{\text{Cov}(w, z)}{\bar{w}} + \frac{E(w \Delta z)}{\bar{w}}$
- Wright-Fisher model: $P(k \text{ copies}) = \binom{N}{k} p^k (1-p)^{N-k}$
- Adaptive dynamics: $\frac{dx}{dt} = m \cdot s \cdot \frac{\partial w}{\partial x}$

Implications

Ecological:
- Diversity predicts community stability
- Resource competition predicts which species persist
- Cross-feeding creates interdependence between species
- Priority effects create historical contingency in community assembly
- Succession predicts changing community composition over time

Genetic:
- Genetic architecture predicts adaptation speed
- Polygenic adaptation can be slow when selection is weak
- Epistasis makes adaptation path-dependent
- Clonal interference slows adaptation in asexual populations
- Genetic trade-offs constrain adaptation

Evolutionary:
- Selection favors genotypes with higher fitness
- Drift can overwhelm selection when selection is weak or population size is small
- Mutation creates genetic variation
- Migration introduces genetic variation from other populations
- Adaptation follows the fitness gradient

Cross-Domain:
- Ecological diversity predicts genetic diversity
- Genetic diversity predicts ecological diversity
- Ecological succession predicts changing selection pressures
- Ecological feedbacks predict evolutionary feedbacks
- Ecological stability predicts evolutionary stability
- Ecological tipping points predict evolutionary tipping points

Applications

Medicine:
- Antibiotic resistance: Clonal interference slows adaptation in asexual populations
- Antibiotic resistance: Epistasis makes adaptation path-dependent
- Antibiotic resistance: Genetic trade-offs constrain adaptation
- Antibiotic resistance: Selection favors genotypes with higher fitness
- Antibiotic resistance: Drift can overwhelm selection when selection is weak or population size is small
- Antibiotic resistance: Mutation creates genetic variation
- Antibiotic resistance: Migration introduces genetic variation from other populations
- Antibiotic resistance: Adaptation follows the fitness gradient
- Antibiotic resistance: Ecological succession predicts changing selection pressures
- Antibiotic resistance: Ecological feedbacks predict evolutionary feedbacks
- Antibiotic resistance: Ecological stability predicts evolutionary stability
- Antibiotic resistance: Ecological tipping points predict evolutionary tipping points
- Antibiotic resistance: Ecological diversity predicts genetic diversity
- Antibiotic resistance: Genetic diversity predicts ecological diversity