

1-D Simulation of Western Boundary Current

Chuning Wang

February 21, 2017

1 Model setup

The numerical model solved in this report is based on the 1-D vorticity equation

$$\frac{\partial}{\partial t}\phi_{xx} + \beta\phi_x = WC - \gamma\phi_{xx} \quad (1)$$

where the terms on the left side are the 'storage term' and ' β effect term', while terms on the right side are wind curl and friction, respectively. To solve this equation numerically, the spatial derivation terms must be discretized with finite difference method:

$$(\phi_x)_i^n = \frac{\phi_{i+1}^n - \phi_{i-1}^n}{2\Delta x} \quad (2)$$

$$(\phi_{xx})_i^n = \frac{\phi_{i+1}^n + \phi_{i-1}^n - 2\phi_i^n}{\Delta x^2} \quad (3)$$

and temporal derivation can discretized with either forward scheme

$$\frac{\partial}{\partial t}(\phi_{xx})_i = \frac{(\phi_{xx})_i^{n+1} - (\phi_{xx})_i^n}{\Delta t} \quad (4)$$

or leap frog scheme

$$\frac{\partial}{\partial t}(\phi_{xx})_i = \frac{(\phi_{xx})_i^{n+1} - (\phi_{xx})_i^{n-1}}{2\Delta t} \quad (5)$$

The superscript n and subscript i are indices for time steps and spatial grid points, respectively. Substitute Eq 4 into Eq 1 and rearrange terms, we have

$$(\phi_{xx})_i^{n+1} = (\phi_{xx})_i^n + \Delta t(-\beta(\phi_x)_i^n + WC - \gamma(\phi_{xx})_i^n) \quad (6)$$