

Moiré Superlattice as an Axion-like Optical System

Connecting Strong/Weak Coupling Alternation, Exciton-Polaritons,
and Minibands to Axion-like Emergent Physics

Honeycomb Serpentine Optics Series · Chur · 2026

Abstract

This document explores how three emergent phenomena in Moiré superlattices — spatially alternating strong/weak coupling, exciton-polariton quasi-particles, and miniband flat-band formation — collectively produce physics that is mathematically analogous to axion electrodynamics. We connect this theoretical framework to the Honeycomb Serpentine polarising-film experiment series, in which N20 DC micro-motors drive a rotating Moiré fringe system and optical phase modulation is directly observable via cluster-formation and satellite-to-cluster transition dynamics.

1. Background: What Is an Axion?

In particle physics, the axion is a hypothetical elementary particle proposed by Peccei and Quinn (1977) to resolve the strong CP problem. Its defining interaction with the electromagnetic field is governed by the term:

$$\mathcal{L}_{\text{axion}} = (g_{a\gamma\gamma} / 4) \cdot \theta \cdot F_{\mu\nu} \cdot \tilde{F}^{\mu\nu} = g_{a\gamma\gamma} \cdot \theta \cdot \mathbf{E} \cdot \mathbf{B}$$

where θ is an extremely small vacuum angle, $F_{\mu\nu}$ is the electromagnetic field tensor, and $g_{a\gamma\gamma}$ is the axion-photon coupling constant. Three features make axions distinctive:

- Extremely light mass (10^{-10} – 10^{-3} eV), far below any known Standard Model particle.
- Extremely weak interactions — passes through ordinary matter almost unimpeded.
- Couples indirectly to the strong force through gluon anomaly, resolving CP asymmetry.

In condensed matter, an **effective axion field** $\theta(\mathbf{r},t)$ can emerge from topological band structure without requiring a new fundamental particle. This is the sense in which Moiré systems are 'axion-like'.

2. Moiré Superlattice — Mechanisms Relevant to This Experiment

2.1 Strong / Weak Coupling Alternation and $\theta(\mathbf{r})$

When two periodic lattices (or polarising films) are stacked with a relative twist angle ϕ , regions of strong AA-stacking (constructive overlap) alternate with AB/BA regions (weak coupling). The spatial period of this alternation defines the Moiré wavelength:

$$\lambda_M = \lambda / (2 \cdot \sin(\phi/2))$$

The coupling strength $J(\mathbf{r})$ modulates sinusoidally across the supercell. Crucially, a spatially varying coupling constant maps directly onto a **spatially varying effective gauge potential** $A_{\text{eff}}(\mathbf{r})$, and the curl of this potential

yields an effective magnetic flux — the Berry curvature. Integrating the Berry curvature over the Moiré Brillouin zone gives the topological invariant θ :

$$\theta = (1/2\pi) \int \Omega(k) d^2k$$

When $\theta = \pi$ the system is in the axion insulator phase; when $\theta \neq 0, \pi$ it breaks time-reversal symmetry and supports a Hall-like magneto-electric response identical in form to the axion term L_{axion} above.

2.2 Exciton-Polaritons as the Photon-Axion Mixing Analogue

A polariton is a quantum superposition of a photon and an exciton (bound electron-hole pair). In transition-metal dichalcogenide (TMD) Moiré systems such as MoSe₂/WSe₂, the Moiré potential traps excitons at lattice sites, and strong light-matter coupling produces:

$$H_{\text{coupling}} = g \cdot (a^\dagger + a)(b^\dagger + b)$$

where a is the photon annihilation operator and b the exciton operator. In the real axion Lagrangian the analogous coupling reads $g_{\text{axion}} \cdot \theta \cdot E \cdot B$. The mathematical structure is identical: a bosonic matter field linearly coupled to the electromagnetic field with a coupling constant that encodes the topological angle. When this coupling is topologically protected (i.e., θ is locked by band topology), the polariton becomes a faithful laboratory model of axion-photon mixing, accessible via standard optical spectroscopy.

2.3 Minibands, Flat Bands, and the Axion Insulator Phase

Zone-folding of the original band structure into the smaller Moiré Brillouin zone creates **minibands**. As the twist angle decreases, miniband widths W narrow according to:

$$W \propto t \cdot (a/\lambda_M)^2$$

where t is the hopping integral and a the lattice constant. At the 'magic angle' ($\phi \approx 1.1^\circ$ for graphene) $W \rightarrow 0$, producing a **flat band** in which kinetic energy $t \ll U$ (Coulomb interaction). In this regime strong correlations dominate, and topological band invariants stabilise the axion insulator phase ($\theta = \pi$). Materials such as MnBi₂Te₃ realise exactly this phase experimentally.

3. Axion vs Moiré Effective Analogy — Side-by-Side

Property	True Axion (particle)	Moiré Effective Analogy
Origin	Peccei-Quinn symmetry breaking	Geometric Berry phase
θ value	Cosmological, $\sim 10^{-16}$	Tunable $0 - 2\pi$ via twist angle
E·B coupling	In vacuum	In effective medium (material)
New particle?	Yes (hypothetical)	No — emergent quasi-particle
Experimental access	Extremely difficult	Optical / electrical measurements
Mass scale	$10^{-5} - 10^{-3}$ eV	Quasi-particle gap (meV range)
Time-reversal	Breaks T if $\theta \neq 0, \pi$	Magnetic Moiré breaks T

4. Connection to the Honeycomb Serpentine Experiment Series

The Honeycomb Serpentine experiment uses stacked polarising films driven by N20 DC micro-motors to create a dynamic, rotating Moiré fringe pattern. The following correspondences map the theoretical framework directly

onto the experimental observables.

4.1 Polarising-film twist $\rightarrow \theta(r)$ phase field

The relative angle ϕ between the input and output polarising films sets the Moiré wavelength $\lambda_M = \lambda / (2 \sin(\phi/2))$. Spatially, the transmission function of two crossed polarisers follows Malus's law:

$$T(r) = \cos^2(\Delta\phi(r))$$

The local phase $\Delta\phi(r)$ varies across the fringe pattern exactly as the effective axion field $\theta(r)$ varies across a topological Moiré supercell. Removing the input-side privacy film (as in the most recent experimental run) is equivalent to removing a boundary condition on θ , freeing the phase field and producing the observed satellite-to-cluster transition.

4.2 Cluster formation/detachment $\rightarrow$ domain-wall dynamics

In axion insulators, boundaries between regions with $\theta = 0$ and $\theta = \pi$ host topologically protected surface states. In the polarising-film system, cluster formation marks the nucleation of a high-contrast fringe domain, while detachment is the propagation of its boundary — directly analogous to axion domain-wall motion. The measured quasi-periodic detachment frequency:

$$f_{\text{detach}} \approx 0.108 \text{ Hz (period} \approx 9.3 \text{ s)}$$

provides a direct experimental handle on the 'domain-wall velocity' in this optical analogue system.

4.3 Rocket-trajectory fringe pattern $\rightarrow$ polariton group velocity

The rocket-trajectory-like fringe propagation observed in the satellite-to-cluster transition corresponds physically to a polariton wavepacket propagating with a group velocity set by the light-matter coupling strength g . In an ideal exciton-polariton system:

$$v_{\text{group}} = (\partial E / \partial k) \approx c \cdot (\Delta E_{\text{Rabi}} / E_{\text{photon}})^{(1/2)}$$

The fringe trajectory slope in the space-time brightness map thus encodes the effective coupling constant of the optical Moiré system.

4.4 Planned experiment: input-side film re-insertion $\rightarrow \theta$ boundary condition

Re-inserting the input-side privacy protection film will re-impose a hard boundary condition on $\theta(r)$, analogous to pinning the axion field at the surface of a topological insulator. Predicted observables:

- Suppression of the satellite-to-cluster transition (θ is pinned, domain walls cannot nucleate freely).
- Recovery of the quasi-periodic fringe spectrum in the range 0.091–0.389 Hz.

• Possible appearance of a new frequency corresponding to the cavity round-trip of the effective polariton (set by film thickness and motor speed).

Varying the input-film angle independently of the output film will allow systematic mapping of θ as a continuously tunable experimental parameter.

5. Key Equations at a Glance

Symbol / Equation	Physical meaning
$\lambda_M = \lambda / (2 \sin(\phi/2))$	Moiré superlattice period
$\theta = (1/2\pi) \int \Omega(k) d^2k$	Topological axion angle (Berry curvature integral)
$L = g_{a\gamma\gamma} \cdot \theta \cdot E \cdot B$	Axion-photon coupling Lagrangian
$H = g(a^\dagger+a)(b^\dagger+b)$	Polariton Hamiltonian (photon-exciton coupling)
$W \propto t \cdot (a/\lambda_M)^2$	Miniband width — narrows toward flat band
$T(r) = \cos^2(\Delta\phi(r))$	Malus's law as optical $\theta(r)$ field
$f_{\text{detach}} \approx 0.108 \text{ Hz}$	Measured cluster detachment frequency (domain-wall rate)

6. Conclusions

Three independent mechanisms in Moiré superlattices — strong/weak coupling alternation, exciton-polariton formation, and miniband flat-band physics — converge on the same effective Lagrangian as axion electrodynamics. The mathematical correspondence is not merely analogical: the Berry-curvature integral that defines θ is structurally identical in both cases, and the polariton coupling Hamiltonian maps onto the axion-photon interaction term term-by-term.

The Honeycomb Serpentine polarising-film experiment provides a macroscopic, fully accessible realisation of this physics. The twist angle ϕ controls θ continuously; the cluster-detachment dynamics probe domain-wall propagation; the fringe trajectory encodes the effective polariton group velocity. The planned follow-up experiment — re-inserting the input-side privacy film — will directly test the θ boundary-condition hypothesis and may reveal a new frequency mode associated with polariton cavity round-trips.

Key takeaway

Moiré fringe systems are not merely visual curiosities. When the coupling geometry is controlled (twist angle, film identity, motor speed), they become tuneable table-top laboratories for topological and axion-like electromagnetic phenomena — accessible with polarising film, an N20 motor, and a camera.

Selected References

[1] Peccei, R.D. & Quinn, H.R. (1977). CP conservation in the presence of pseudoparticles. *Phys. Rev. Lett.* 38, 1440.

[2] Li, L. et al. (2021). Quantum anomalous Hall effect from intertwined Moiré bands. *Nature* 600, 641–646.

- [3] Qi, X.-L., Hughes, T.L. & Zhang, S.-C. (2008). Topological field theory of time-reversal invariant insulators. *Phys. Rev. B* 78, 195424.
- [4] Shimazaki, Y. et al. (2020). Strongly correlated electrons and hybrid excitons in a Moiré heterostructure. *Nature* 580, 472–477.
- [5] Chur (2025–2026). Honeycomb Serpentine Moiré Optics Experiment Series. GitHub repository.