1 Exercise 11.9

验证 point separation, $||T|| = 0 \Rightarrow \forall x, ||Tx|| \leq M||x|| = 0 \Rightarrow T = 0$ 验证 absolute homogeneity, $\forall a \in \mathbb{F}, v \in \mathcal{V}(\mathbb{F}^n, \mathbb{F}^m)$ 有 $||av|| = \inf\{M \geq 0 : \forall x \in \mathbb{F}^n, |avx| = |a||vx| \leq M|x|\}$ 若 |a| = 0, ||av|| = 0 = |a|||v||, 否则,有 $||av|| = \inf M \geq 0 : \forall x \in \mathbb{F}^n, |vx| \leq \frac{M}{|a|}|x|$, 也有 ||av|| = |a|||v||验证 triangle inequality, $\forall u, v \in \mathcal{V}(\mathbb{F}^n, \mathbb{F}^m)$ $||u+v|| = \inf\{M \geq 0 : \forall x \in \mathbb{F}^n, |(u+v)x| = |ux+vx| \leq M|x|\}$ 设 $||u|| = M_1, ||v|| = M_2$,则有 $\forall x \in \mathbb{F}^n, |ux| \leq M_1|x|, |vx| \leq M_2|x| \Rightarrow |ux| + |vx| \leq (M_1 + M_2)|x|$ 有 $||u+v|| \leq M_1 + M_2 = ||u|| + ||v||$ 所以有 $||\cdot||$ 是一个范数

2 Exercise 11.13

验证 non-negativity, 由范数的非负性, $d(S,T) = ||S-T|| \ge 0$ 验证 identity of indiscernibles, $d(S,T) = 0 \Leftrightarrow ||S-T|| = 0 \Leftrightarrow \forall x |(S-T)x| = 0 = |Sx-Tx| \Leftrightarrow \forall x Sx = Tx \Leftrightarrow S = T$

验证 real positivity,由 $||T|| \ge \inf\{M \ge 0\} = 0 \Rightarrow ||T|| \ge 0$

验证 symmetry, d(S,T) = ||S-T|| = ||T-S|| = d(T,S)验证 triangle inequality, $d(A,B) = ||A-B|| \le ||A-C|| + ||C-B|| = d(A,C) + d(C,B)$

3 Exercise 11.16

real positivity, point separation, absolute homogeneity 显然, 下证 triangle inequality:

$$||u+v||^2 = \sum_{j=1}^n ||ue_j + ve_j||^2$$
(1)

$$= \sum_{j=1}^{n} (||ue_j||^2 + ||ve_j||^2 + 2\langle ue_j, ve_j \rangle)$$
(2)

$$= ||u||^2 + ||v||^2 + 2\sum_{j=1}^n \langle ue_j, ve_j \rangle (\text{由内积的性质})$$
(3)

$$\leq ||u||^2 + ||v||^2 + 2\sqrt{\sum_{j=1}^n ||ue_j||^2 ||ve_j||^2}$$
(由柯西不等式) (4)

$$\leq ||u||^2 + ||v||^2 + 2\sqrt{\sum_{j=1}^n ||ue_j||^2 \sum_{j=1}^n ||ve_j||^2}$$
(5)

$$\leq ||u||^2 + ||v||^2 + 2\sqrt{||u||^2||v||^2} \tag{6}$$

$$\leq ||u||^2 + ||v||^2 + 2||u||||v|| \tag{7}$$

$$\leq (||u|| + ||v||)^2 \tag{8}$$

(9)

两边同时开平方即可得 $||u+v|| \leq ||u|| + ||v||$

4 Exercise 11.20

由 11.18 有 $|TSx| \le |T||Sx|$ 则 $|TS|^2 = \sum_{j=1}^k ||TSe_j||^2 \le \sum_{j=1}^k (|T||Se_j||)^2 = |T|^2 \sum_{j=1}^k (|Se_j||)^2 = |T|^2 |S|^2$ 两边同时开平方即可得 $|TS| \le |T||S|$