

The Fourier Transform (Part 2)

Dr Chris Jobling (c.p.jobling@swansea.ac.uk)

Digital Technium 123

Office Hours: 12:00-13:00 Mondays

You can view the notes for this presentation in [HTML](#) and [PDF](#).

The source code of this presentation is available in Markdown format from GitHub: [ft1.md](#).

The GitHub repository [EG-247 Resources](#) also contains the source code for all the Matlab/Simulink examples and the Laboratory Exercises.

Note on Notation

If you have been reading both Karris and Boulet you may have noticed a difference in the notation used in the definition of Fourier Transform:

- Karris uses $F(\omega)$
- Boulet uses $F(j\omega)$

I checked other sources and [Hsu \(Schaum's Signals and Systems\)](#) and [Morrell \(The Fourier Analysis Video Series on YouTube\)](#) both use the $F(\omega)$ notation.

According to Wikipedia [Fourier Transform: Other Notations](#) both are used only by electronic engineers anyway and either would be acceptable.

There is some advantage in using Boulet's notation $F(j\omega)$ in that it helps to reinforce the idea that Fourier Transform is a special case of the Laplace Transform and it was the notation that I used in the last session.

In these notes, I've used the other convention on the basis that its the more likely to be seen in your support materials.

However, I am happy to change back if you find the addition of j useful.

You should be aware that Fourier Transforms are in general complex so whatever the notation used to *represent* the transform, we are still dealing with real and imaginary parts or magnitudes and phases when we use the *actual transforms* in analysis.

Scope and Background Reading

This session continues our introduction to the Fourier Transform with several examples extracted from tables of transform tables.

The material in this presentation and notes is based on Chapter 8 (Starting at Section 8.4) of [Steven T. Karris, Signals and Systems: with Matlab Computation and Simulink Modelling, 5th Edition](#). from the Required Reading List. I also used Chapter 5 of [Benoit Boulet, Fundamentals of Signals and Systems](#) from the Recommended Reading List.

Agenda

Last time

- Fourier Transform as the Limit of a Fourier Series
- Doing the Maths
- The Fourier Transform
- Properties of the Fourier Transform
- Some Examples
- Computing Fourier Transforms in Matlab

This Time

- Tables of Transform Pairs
- Examples of Selected Transforms
- Relationship between Laplace and Fourier
- Fourier Transforms of Common Signals

Reminder of the Definitions

Last time we derived the Fourier Transform by evaluating what would happen when a periodic signal was made periodic. Let us restate the definitions.

The Fourier Transform

In the signals and systems context, the Fourier Transform is used to convert a function of time $f(t)$ to a function of radian frequency $F(\omega)$:

$$\mathcal{F}\{f(t)\} = \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt = F(\omega).$$

The Inverse Fourier Transform

In the signals and systems context, the *Inverse Fourier Transform* is used to convert a function of frequency $F(\omega)$ to a function of time $f(t)$:

$$\mathcal{F}^{-1}\{F(\omega)\} = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{j\omega t} d\omega = f(t).$$

Note, the factor 2π is introduced because we are changing units from radians/second to seconds.

Duality of the transform

Note the similarity of the Fourier and its Inverse. This has important consequences in filter design and later when we consider sampled data systems.

Table of Common Fourier Transform Pairs

This table is adapted from Table 8.9 of Karris. See also: [Wikibooks: Engineering Tables/Fourier Transform Table](#) and [Fourier Transform—WolframMathworld](#) for more complete references.

Examples of Selected Transforms

Relationship between $f(t)$ and $F(\omega)$

In most of the work we will do in this course, and in practice, the *signals* that we use with the Fourier transform will be a real continuous aperiodic functions of time that are zero when $t = 0$.

The Fourier transforms of such a signals will be complex continuous function of frequency which have real and imaginary parts and will exist at both positive and negative values of ω .

It is often most convenient to deal with the transformed “spectrum” by considering the magnitude and phase and we will therefore often plot $F(\omega)$ on two separate graphs as *magnitude* $|F(\omega)|$ and *phase* $\angle F(\omega)$ (where phase is measured in radians) plotted against frequency $\omega \in [-\infty, \infty]$ (in radians/second).

We most often represent the *system* by its so-called *frequency response* and we will be interested in what effect the system has on the signal $f(t)$.

As for the Laplace transform, this is more conveniently determined by exploiting the *time convolution property*. That is by performing a Fourier transform of

the signal, multiplying it by the system's frequency response and then inverse Fourier transforming the result.

Have these ideas in mind as we go through the examples in the rest of this session.

The Dirac Delta

$$\delta(t) \Leftrightarrow 1$$




Figure 1: Fourier transform of $\delta(t)$

Proof: uses sampling and sifting properties of $\delta(t)$.

Matlab:

```
syms t;
fourier(dirac(t))
```

Related:

$$\delta(t - t_0) \Leftrightarrow e^{-j\omega t_0}$$

DC

$$1 \Leftrightarrow 2\pi\delta(\omega)$$

Matlab:

```
syms A t;
fourier(1)
```




Figure 2: Fourier transform of DC

Related by frequency shifting property:

$$e^{j\omega_0 t} \Leftrightarrow 2\pi\delta(\omega - \omega_0)$$

Cosine wave (Sinewave with even symmetry)

$$\cos(t) = \frac{1}{2} (e^{j\omega_0 t} + e^{-j\omega_0 t}) \Leftrightarrow \pi\delta(\omega - \omega_0) + \pi\delta(\omega + \omega_0)$$




Figure 3: Fourier transform of cosine wave

Note: $f(t)$ is real and even. $F(\omega)$ is also real and even.

Sine wave

$$\sin(t) = \frac{1}{j2} (e^{j\omega_0 t} - e^{-j\omega_0 t}) \Leftrightarrow j\pi\delta(\omega - \omega_0) - j\pi\delta(\omega + \omega_0)$$

Note: $f(t)$ is real and odd. $F(\omega)$ is imaginary and odd.




Figure 4: Fourier transform of sine wave

Signum (Sign)

The signum function is a function whose value is equal to

$$\operatorname{sgn} x = \begin{cases} -1 & x < 0 \\ 0 & x = 0 \\ +1 & x > 0 \end{cases}$$

This function is often used to model a *voltage comparitor* in circuits.

The transform is:

$$\operatorname{sgn} x = u_0(t) - u_0(-t) = \frac{2}{j\omega}$$

Example 1: Unit Step

Use the signum function to show that

$$\mathcal{F}\{u_0(t)\} = \pi\delta(\omega) + \frac{1}{j\omega}$$

Clue

Define




Figure 5: Signum function

$$u_0(t) = 2 \operatorname{sgn} x - 1$$

Does that help?




Figure 6: Unit step defined using signum

Proof

$$\operatorname{sgn} x = 2u_0(t) - 1$$

so

$$u_0(t) = \frac{1}{2} [1 + \operatorname{sgn} x]$$

From previous results $1 \Leftrightarrow 2\pi\delta(\omega)$ and $\operatorname{sgn} x = 2/(j\omega)$ so by linearity

$$u_0(t) \Leftrightarrow \pi\delta(\omega) + \frac{1}{j\omega}$$

QED

Graph of unit step

$$u_0(t) \Leftrightarrow \pi\delta(\omega) + \frac{1}{j\omega}$$




Figure 7: FT of unit step

Unit step is neither even nor odd so the Fourier transform is complex with real part $F_{\text{Re}}(\omega) = \pi\delta(\omega)$ and imaginary part $F_{\text{Im}}(\omega) = 1/(j\omega)$. The real part is even, and the imaginary part is odd.

Example 2

Use the results derived so far to show that

$$e^{j\omega_0 t} u_0(t) \Leftrightarrow \pi \delta(\omega - \omega_0) + \frac{1}{j(\omega - \omega_0)}$$

Hint: linearity plus frequency shift property.

Example 3

Use the results derived so far to show that

$$\sin \omega_0 t u_0(t) \Leftrightarrow \frac{\pi}{j2} [\delta(\omega - \omega_0) - \delta(\omega + \omega_0)] + \frac{\omega_0}{\omega_0^2 - \omega^2}$$

Hint: Euler's formula plus solution to example 2.

Important note: the equivalent example in Karris (Section 8.4.9 Eq. 8.75 pp 8-23—8-24) is wrong!

See worked solution on Blackboard for the corrected proof.

Example 4

Use the result of Example 3 to determine the Fourier transform of $\cos \omega_0 t u_0(t)$.

Answer

$$\cos \omega_0 t u_0(t) \Leftrightarrow \frac{\pi}{2} [\delta(\omega - \omega_0) + \delta(\omega + \omega_0)] + \frac{j\omega}{\omega_0^2 - \omega^2}$$

Derivation of the Fourier Transform from the Laplace Transform

If a signal is a function of time $f(t)$ which is zero for $t \leq 0$, we can obtain the Fourier transform from the Laplace transform by substituting s by $j\omega$.

Example 5: Single Pole Filter

Given that

$$\mathcal{L}\{e^{-at}u_0(t)\} = \frac{1}{s+a}$$

Compute

$$\mathcal{F}\{e^{-at}u_0(t)\}$$

Boulet gives the graph of this function.

Example 6: Complex Pole Pair (cos term)

Given that

$$\mathcal{L}\{e^{-at} \cos \omega_0 t u_0(t)\} = \frac{s+a}{(s+a)^2 + \omega_0^2}$$

Compute

$$\mathcal{F}\{e^{-at} \cos \omega_0 t u_0(t)\}$$

Boulet gives the graph of this function.

Fourier Transforms of Common Signals

We shall conclude this session by computing as many of the the Fourier transform of some common signals as we have time for.

- rectangular pulse
- triangular pulse
- periodic time function
- unit impulse train (model of regular sampling)

I will not provide notes for these, but you will find more details in Chapter 8 of Karris and Chapter 5 of Boulet and I have created some worked examples to help with revision which you'll find on Blackboard.

Suggestions for Further Reading

Boulet has several interesting amplifications of the material presented by Karris. You would be well advised to read these. Particular highlights which we will not have time to cover:

- Time multiplication and its relation to amplitude modulation (pp 182—183).
- Fourier transform of the complex exponential signal $e^{(\alpha+j\beta)t}$ with graphs (pp 184—187).
- Use of inverse Fourier series to determine $f(t)$ from a given $F(j\omega)$ and the “ideal” low-pass filter (pp 188—191).
- The Duality of the Fourier transform (pp 191—192).

End of Part 2

Summary

- Tables of Transform Pairs
- Examples of Selected Transforms
- Relationship between Laplace and Fourier
- Fourier Transforms of Common Signals

Next Time

- The Fourier Transform for Systems and Circuit Analysis

Homework

Attempt Questions 1—6 of the End of Chapter Problems (Section 8.10) in Karris.