

# Crosstalk of TGF- $\beta$ and Estrogen Receptor Signaling in Breast Cancer

Arja M. Band · Marikki Laiho

Received: 23 February 2011 / Accepted: 28 February 2011 / Published online: 11 March 2011  
© Springer Science+Business Media, LLC 2011

**Abstract** Estrogen receptor- $\alpha$  (ER $\alpha$ ) and transforming growth factor (TGF)- $\beta$  signaling pathways are major regulators during mammary gland development, function and tumorigenesis. Predominantly, they have opposing roles in proliferation and apoptosis. While ER $\alpha$  signaling supports growth and differentiation and is antiapoptotic, mammary gland epithelia cells are very sensitive to TGF- $\beta$ —induced cell cycle arrest and apoptosis. Their regulatory pathways intersect, and ER $\alpha$  blocks TGF- $\beta$  pathway by multiple means, including direct interactions of its signaling components, Smads. However, relatively little is known of the dysfunction of their interactions in cancer. A better understanding would help to develop new strategies for breast cancer treatment.

**Keywords** Estrogen receptor · TGF- $\beta$  signaling · Breast cancer · Smad · SnoN

## Abbreviations

AF	Activation function
EMT	Epithelial-to-mesenchymal transition
ER $\alpha$	Estrogen receptor- $\alpha$
ER $\beta$	Estrogen receptor- $\beta$
HAT	Histone acetyltransferase
HDAC	Histone deacetylase
MAPK	Mitogen activated protein kinase
PI3K	Phosphatidylinositol 3-OH kinase
R-Smad	Receptor regulated Smad
SnoN	Ski-related novel gene
TAM	Tamoxifen
TGF- $\beta$	Transforming growth factor- $\beta$

## Introduction

Estrogen receptors (ER $\alpha$  and ER $\beta$ ) are ligand-inducible transcription factors and members of the nuclear hormone receptors. ER $\alpha$  signaling is essential for the development of the adult mammary gland. ER $\alpha$  knock-out mouse lacks the development that occurs during pre- and postpubertal stages [1, 2]. Due to its ability to stimulate proliferation ER $\alpha$  is also a driving force during mammary gland tumorigenesis. Approximately 70% of breast cancers are ER $\alpha$  positive and estrogen-dependent. Therefore anti-estrogens like tamoxifen metabolites (TAM) acting as competitive inhibitors of the receptor, or suppression of the synthesis of the ligand by aromatase inhibitors, have been the mainstay of endocrine therapy for breast cancer. ER $\beta$  has been identified relatively recently [3]. It may act as a repressor of ER $\alpha$  transcriptional activity [4] and as a tumor suppressor in other organs, like the prostate [5]. Its expression pattern only partly overlaps with that of ER $\alpha$

---

Financial support: Academy of Finland (grant no. 129699).

---

A. M. Band · M. Laiho  
Molecular Cancer Biology Program, Biomedicum Helsinki and  
Haartman Institute, University of Helsinki,  
Helsinki, Finland

M. Laiho  
Sidney Kimmel Comprehensive Cancer Center, The Johns  
Hopkins University School of Medicine,  
Baltimore, MD, USA

M. Laiho (✉)  
Department of Radiation Oncology and Molecular Radiation  
Sciences, The Johns Hopkins University School of Medicine,  
1550 Orleans Street, Room 444,  
Baltimore, MD 21231, USA  
e-mail: mlaiho1@jhmi.edu

but is expressed in stromal cells in the mammary gland [6]. ER $\beta$  knock-out mouse models suggest that it has a role in the development of ovaries, bladder, and prostate but not the mammary gland [2, 7, 8]. Much less is known about the role of ER $\beta$  tumorigenesis than that of ER $\alpha$ .


TGF- $\beta$  belongs to the TGF- $\beta$  superfamily of cytokines. It consists of more than 30 factors including nodal, activins, inhibins, and bone morphogenic proteins. TGF- $\beta$  is a major regulator in many essential cellular processes including proliferation, differentiation, migration, immune response, and apoptosis. The role of TGF- $\beta$  in cancer is rather complex. It has a dual role in tumor progression. Due to its anti-proliferative effect on most cells it acts as a tumor suppressor, but due to its ability to promote epithelial-to-mesenchymal transition (EMT) and to suppress immune surveillance, it appears to convert to a tumor promoter in later stages of cancer [9].

ER $\alpha$  and TGF- $\beta$  have opposing roles in cell proliferation and apoptosis of epithelial cells. Normal ER $\alpha$ —expressing luminal cells rarely proliferate [10]. However, prolonged exposure to estrogen promotes breast tumorigenesis and ER $\alpha$  has a strong mitogenic activity in breast cancer cells. ER $\alpha$  signaling enhances the transcriptional activation of cyclin D1 and c-Myc [11–13]. ER $\alpha$  prevents apoptosis by controlling both the extrinsic and intrinsic apoptotic pathways and by promoting cell survival [14], whereas TGF- $\beta$  causes cell cycle arrest by inhibiting cyclin dependent kinase activities and by reducing the expression levels of c-Myc in epithelial cells [15]. TGF- $\beta$  can also promote apoptosis or cell survival in a cell-type and context-dependent manner [16].

Considering the intricate roles of these major signaling pathways, ER $\alpha$  and TGF- $\beta$ , in mammary epithelial cells biology and tumorigenesis, and their extensive interactions, a deeper understanding how these pathways crosstalk and are dysregulated would be highly relevant for a better clinical translation. In this article we review the current knowledge and raise aspects for future consideration.

### Regulation of TGF- $\beta$ Signaling Pathway

TGF- $\beta$  family members signal through cell membrane receptors through a cascade of positive and negative



**Figure 1** Intersections between TGF- $\beta$  and ER $\alpha$  pathways

regulation steps that culminate in activation of transcriptional activator complexes. Active TGF- $\beta$  binds to serine/threonine kinase type II receptor, which leads to the recruitment of the type I serine/threonine kinase receptor to the complex and its subsequent phosphorylation. The signaling cascade is mediated by the phosphorylation of receptor regulated Smads (R-Smads), which, for the TGF- $\beta$  family, are represented by Smad2 and Smad3. Phosphorylated R-Smads form a complex with co-Smad, Smad4. Once formed, the complex is translocated to the nucleus where it recruits transcriptional co-activators like p300 and CBP to induce acetylation and activation of the expression of TGF- $\beta$  target genes [17, 18].

Well-controlled TGF- $\beta$  signaling cascade is essential for cell and tissue homeostasis. Therefore the duration and timing are governed by negative regulators, Smad7, SnoN (Ski-related novel gene) and Ski [18, 19]. Upon TGF- $\beta$  stimulation Smad7 is transported from the nucleus to the plasma membrane where it binds the TGF- $\beta$  type I receptor [20]. At the plasma membrane it recruits phosphatases or ubiquitin ligases leading to degradation or dephosphorylation of the type I receptor [21–23]. Smad7 also prevents the formation of the functional transcription complex at the Smad binding element [24]. The negative feedback loop is reinstated by TGF- $\beta$ -induced production of Smad7 to a steady state level.

Similarly to Smad7, SnoN and Ski provide negative feedback loops. TGF- $\beta$  induces rapid proteasomal degradation of SnoN to allow induction of its target genes [25–30]. SnoN and Ski can block TGF- $\beta$  induced transcription through various mechanisms. They recruit N-CoR-histone deacetylase (HDAC) complex to the promoter [31–33]. Alternatively, they may bind to the phosphorylated Smads and prevent the translocation of the Smad complex to the nucleus, or prevent the formation of the active Smad complex at the promoter [34]. Consequent to the initial decrease of SnoN, TGF- $\beta$  induces the production of SnoN to the steady state level (Fig. 1). In some cell types, TGF- $\beta$  induces degradation of also Ski [35, 36]. In addition, SnoN and Ski modify the TGF- $\beta$  signaling cascade by regulating Smad7 production. Both block the Smad-dependent transcriptional induction of Smad7 [37, 38]. Conversely, degradation of SnoN or Ski increases production of Smad7 and causes downregulation of TGF- $\beta$  receptors [37–39]. Thus, upregulation of SnoN and Ski shuts down the production of Smad7 and thus closes down the feedback loop of TGF- $\beta$  signaling.

Although Smad signaling is the main mediator of TGF- $\beta$  responses, TGF- $\beta$  activates also other pathways, like mitogen activated protein kinase (MAPK), phosphatidylinositol 3-kinase (PI3K)/Akt and Rho-like GTPase signaling pathways [40].

## Regulation of ER $\alpha$ Signaling Pathway

ER $\alpha$  nuclear hormone receptor consists of several functionally relevant domains. Two acidic activation domains are located in the amino- and carboxy-termini (activation function 1 and 2, AF-1 and AF-2), which mediate the ligand-dependent transcriptional activation. AF-1 contributes to the constitutive transcriptional activity of the ER $\alpha$  while AF-2 is ligand-dependent. DNA-binding domain is located in a central zinc finger region whereas the ligand-binding domain overlaps with AF-2 [41].

Upon ligand binding the conformation of ER $\alpha$  changes and promotes receptor dimerization and high-affinity binding to estrogen response element (ERE) [42]. ER $\alpha$  regulates gene transcription by formation of a multiprotein complex containing general transcription factors, co-activators, co-repressors, co-integrators, histone acetyltransferases (HATs) and HDACs that provide tissue- and context-specific activation of ER $\alpha$  signaling [43]. Transcriptional activation requires relaxation of the chromatin to loosen the interaction between histones and DNA. This is facilitated by acetylation of the histones. Co-activators enhance transcription by recruiting HATs to the chromatin or contain themselves HAT activity. Co-repressors, on the other hand, may recruit HDACs to remove acetylation and turn off the transcription. Histone methyltransferase also modulates the promoter for transcription. Chromatin immunoprecipitation assays have revealed that up to 46 transcription factors cycle on and off the *pS2* gene promoter in a cyclic manner [44]. Thus, acetylation and methylation status regulate the permissiveness of the promoter and are dependent on the specific association of multiprotein complexes.

In addition to the chromatin remodeling proteins, ER $\alpha$  interacts with and is regulated by several co-activators that enhance its transcriptional activity. Many of these contain a LXXLL motif, a so called nuclear receptor binding domain that ER $\alpha$  binds to [45, 46]. The best characterized co-activators are the p160 family that consists of steroid receptor co-activators (SRC-1, SRC-2), GRIP1 and AIB1 (SRC-3).

In addition to the classical model where ER $\alpha$  bind to EREs in the promoter region, ER $\alpha$  can also regulate gene expression by binding to other transcription factors that are already bound to DNA. These transcription factors include Sp1, AP1, Runx1 and FOXA1. The whole genome contains tens of thousands of putative ER $\alpha$  binding sites but the number of sites at which ER $\alpha$  binds to the chromatin is much less, perhaps thousands [47, 48]. Genome-wide maps for ER $\alpha$  binding sites has indicated that most of the ER $\alpha$  binding to the promoters requires DNA binding capacity of ER $\alpha$  and are dependent on accessibility factors such as Runx1 or FOXA1 [49, 50].

Apart from co-activators and co-repressors, ER $\alpha$  activity can be also modulated by phosphorylation, acetylation and methylation. Several kinases have been implicated in the phosphorylation and transcriptional activation of ER $\alpha$  reporters. These kinases include protein kinase A, MAPK and cyclin A-CDK2 [51–53], opening up the possibility of different signaling pathways to modulate ER $\alpha$  signaling. Acetylation of ER $\alpha$  may either enhance or attenuate its transcriptional activity [54–56]. Methylation of ER $\alpha$  leads to its stabilization [57].

Rapid response to estrogen by several protein kinases has lead to the identification of a membrane bound pool of ERs which may consist of 5–10% of the total amount of ER in cell [58]. Membrane-localized ER dimerizes upon ligand-binding and transactivates several protein kinase cascades. These include MAPK, PKA, protein kinase C, PI3K and Src kinase [58]. Collectively, these are referred to as non-genomic mechanism of ER $\alpha$  because they do not require DNA binding ability or transcriptional activation by ER $\alpha$ . Both nuclear and membrane bound pools of ER contribute the overall cellular response to estrogen. In cancer, intervention of both pathways may be relevant.

## Crosstalk Between TGF- $\beta$ and ER $\alpha$ Pathways in the Breast

The finding that ER $\alpha$ —expressing luminal cells rarely express proliferation markers is intriguing and indicative that ER $\alpha$  mainly functions to conserve and support the differentiated state. However, the breast is an organ that is highly responsive to hormonal stimulation. In order to explore the relationship between proliferation, ER $\alpha$  and TGF- $\beta$  signaling, Ewan and colleagues detected their expression in mouse mammary glands at estrus, and showed co-localization of phosphorylated Smads and nuclear ER $\alpha$ . This indicated co-regulation of the pathways and suggested that TGF- $\beta$  could act to restrict ER $\alpha$ -mediated proliferation [59]. Furthermore, the proliferation rate of mammary gland cells was significantly increased in a mouse model of heterozygous TGF- $\beta$ 1 expression, and proliferation marker Ki67 expressing cells frequently co-expressed ER $\alpha$ .

Several other observations have been made over the years for the crosstalk between ER activity and TGF- $\beta$  signaling. Estrogen receptor activation has been reported to inhibit transcriptional activity of TGF- $\beta$  reporter assays by up to 60% [60–63]. It inhibits TGF- $\beta$ —induced cell migration [61]. Furthermore, microarray analysis in MCF-7 breast cancer cells revealed that TGF- $\beta$  treatment increased more than two fold the expression of 956 genes whereas estrogen treatment reduced the expression of 683 genes of these [63], indicating that ER $\alpha$  is a major modifier of TGF- $\beta$

signaling cascade. Similarly, activin and ER signaling reciprocally suppress each other [64].

Overexpression of Snail transcription factor, which drives EMT, in ER $\alpha$ -expressing MCF-7 cells led to down-regulation of ER $\alpha$  and increased expression of TGF- $\beta$  signaling components [65]. Additionally, it has been reported that inhibition of breast cancer cell growth by antiestrogens TAM and ICI 182,780, is mediated by TGF- $\beta$  [66]. Treatment of MCF-7 cells with antiestrogen leads to increased secretion of active forms of TGF- $\beta$ 1 and TGF- $\beta$ 2 as well as increased expression of TGF- $\beta$  type II receptor [67, 68]. These antiestrogen activities are mediated through p38 MAPK [68]. Furthermore, expression levels of the type II receptor in ER-negative tumors correlate with more aggressive tumors [68].

TGF- $\beta$  is also a potent immunosurveillance factor. In mouse mammary tumor models generated by deletion of the type II receptor, accrual of immune cells in the tumor, and notably a subset of tumor-promoting immune suppressive myeloid cells, was observed [69]. Therefore an important feature of TGF- $\beta$  is to regulate immune responses. Bieri and colleagues conducted molecular profiling of breast cancer expression signatures and compared to those of TGF- $\beta$  response signatures [70]. Intriguingly, the TGF- $\beta$  response signature correlated with ER-negative tumors and poor prognosis. However, complete absence of TGF- $\beta$  signaling correlated with ER-positive and lymph node-positive tumors and poor prognosis. The latter profile appeared to result from increased expression of chemokines driving the recruitment of the myeloid suppressor cells [69]. On the other hand, also antiestrogens may induce immunosuppression in the tumor microenvironment in a manner that depends on TGF- $\beta$  signaling [71]. While the explicit connections of these ER regulatory events to TGF- $\beta$ —regulated pathways remain to be established, they indicate that both pathways are interlinked in their ability to maintain epithelial tissue homeostasis and govern also the metastatic ability of breast cancer cells.

#### Smad-ER $\alpha$ Interactions

Estrogen inhibits TGF- $\beta$  signaling (Fig. 1). Estrogen treatment has been reported to reduce the phosphorylation of Smad2 and Smad3 [61, 63]. ER $\alpha$  physically interacts with Smad2 and Smad3 but only when ER $\alpha$  is in a ligand activated form [60, 63]. This interaction leads to recruitment of ubiquitin ligase, Smurf1, to the complex and ubiquitylation of Smad2/3 and subsequently Smad degradation through ubiquitin–proteasome pathway [63]. However, also contrasting reports on the ability of ER $\alpha$  to reduce Smad2/3 levels exist [62, Band and Laiho, unpublished]. The ability of ER $\alpha$  to enhance the degradation of Smads and to reduce TGF- $\beta$  signaling does not require ER $\alpha$  DNA binding or transcrip-

tional activity. Therefore ER $\alpha$  appears to regulate TGF- $\beta$  signaling also through a non-genomic mechanism. Interestingly, also ER $\beta$  may inhibit TGF- $\beta$  signaling by enhancing Smad2/3 degradation in an estrogen-dependent manner [63]. Consequently, activation of ER $\alpha$  signaling leads to the reduction of both TGF- $\beta$  induced migration and invasion of breast cancer cells [61, 63].

Smad4 acts as an inhibitor of ER $\alpha$ . Smad4 interaction with ER $\alpha$  is activated by antiestrogens, occurs at the ERE binding sites, and negatively regulates ER $\alpha$  transcriptional activity [72, 73]. In contrast to Smad2/3, the interaction with ER $\alpha$  does not lead to its degradation [72]. It is believed that overexpression of Smad3 or inhibition of Smad4 leads to a change of role of TGF- $\beta$  from a repressor to an activator of the ER $\alpha$  signaling cascade [73]. On the other hand, Smad4 can induce apoptosis in an ER $\alpha$ -dependent manner in ER $\alpha$  positive but not in ER $\alpha$  negative breast cancer cells. Smad4 promotes expression of the pro-apoptotic proteins, Bim and Bax, as well as release of cytochrome c [74].

#### SnoN in Breast Cancer

The negative regulator of TGF- $\beta$  signaling, SnoN, is widely expressed in adult and embryonic cells. SnoN is structurally and functionally homologous to Ski [75, 76]. Four isoforms of Sno exist in human cells: SnoA, SnoN, SnoN2 and SnoI. The level of SnoN is directly linked to its ability to repress TGF- $\beta$  signaling. Overexpression of SnoN or stabilization of SnoN, by mutation of its three lysine residues that are ubiquitylated, leads to resistance to TGF- $\beta$ —induced growth arrest [25, 27]. Several ubiquitin ligases are recruited to SnoN by the Smad2/3 and responsible for the proteasomal degradation [30]. The inability of TGF- $\beta$  to cause SnoN degradation results in resistance to TGF- $\beta$ —induced growth arrest in esophageal cancer cells [77]. A high level of SnoN expression has been found in many cancers and cancer cell lines including breast cancer [75, 78, 79]. Like TGF- $\beta$ , SnoN has a dual role in cancer. It has been reported to act as both a tumor promoter and a tumor suppressor. When the levels of SnoN are reduced by RNA interference, the ability of the cells to respond to TGF- $\beta$  induced growth arrest is restored [79]. A reduction of SnoN expression also reduces anchorage-independent growth of MDA-MB-231 breast cancer cells in soft agar and tumor growth in nude mice [79]. Furthermore, elevated SnoN levels co-operate with polyoma middle T-antigen in the formation of aggressive multifocal adenocarcinomas in mouse mammary glands [80]. On the other hand, SnoN $^{+/-}$ heterozygous mice are hypersensitive to chemical carcinogens and develop spontaneous lymphomas [81]. This suggests that SnoN may also act as a tumor suppressor. Furthermore, downregulation of SnoN leads to



increased EMT and metastasis during breast tumorigenesis [79].

SnoN is ubiquitously expressed in adult and embryonic tissues albeit at low levels. In mammary gland, its expression is detected in luminal epithelial cells of the ducts and in lobular cells [78]. In mouse, SnoN expression in the mammary gland peaks in late pregnancy and early lactation and is then rapidly downregulated [80]. These associations indicate that its expression, and likely function, is coupled to mammary epithelial cell differentiation. We have previously investigated SnoN levels in human breast cancer [78]. We analyzed SnoN expression by immunohistochemistry in a tissue microarray of 1122 breast carcinomas. Ninety percent of tumors contained a moderate to high level of SnoN. Low levels of SnoN expression correlated with longer distant disease free survival than in patients whose tumors contained high levels of SnoN. Interestingly, this phenomenon was only observed in ER $\alpha$  positive tumors. Consequently, we have recently investigated the possible mechanism of crosstalk between TGF- $\beta$  and ER $\alpha$  signaling pathways. We find that SnoN contains two highly conserved nuclear receptor binding LxxLL-like motifs. SnoN interacts with ER $\alpha$  at pS2 target gene promoter and is a potent transcriptional activator of ER $\alpha$  (Band and Laiho, unpublished). These findings indicate that SnoN acts as ER $\alpha$ -coactivator. Therefore TGF- $\beta$  and ER $\alpha$  pathways intersect also through SnoN, which further underscores its importance in breast cancer.

## Conclusions

Crosstalk between ER $\alpha$ , insulin-like growth factor and epidermal growth factor/HER2 pathways have been intensively studied. Relatively little is known of the interaction between ER and TGF- $\beta$  signaling cascade although they are two major regulators during the mammary gland development and tumorigenesis.

Anti-estrogen therapy such as TAM has been a successful endocrine treatment for breast cancer for decades. However, its long-term use is limited by TAM resistance which develops for most of the treated women within three years. The relevance of the plasma membrane bound ER, and its effects on calcium signaling, cAMP and activation of several kinases open up the possibility of ER signaling to influence several signaling pathways. New strategies have been developed using combination treatments blocking other signaling pathways with estrogen. Dysregulation of these crosstalk mechanisms may affect the outcome for the breast cancer treatment. Therefore, further understanding of crosstalk of ER and other pathways is needed to be able to develop these treatments.

While evidence is accruing that estrogen signaling inhibits TGF- $\beta$  pathway by multiple means through the Smads (direct interaction, decreased phosphorylation and activity, increased degradation), these responses are challenging to convert to a translation of their implications in breast cancer. The more so considering the dual aspects of TGF- $\beta$  acting both as a tumor suppressor and tumor promoter during breast carcinogenesis and the profound effect of the tumor microenvironment on these activities. Therefore, more molecular models, combined with analyses of the interaction at different stages of breast tumorigenesis, and addressing of their relevance in the clinical setting will take forward our understanding of these signaling cascades and their clinical translation.

## References

1. Bocchinfuso WP, Korach KS. Mammary gland development and tumorigenesis in estrogen receptor knockout mice. *J Mammary Gland Biol Neoplasia*. 1997;2:323–34.
2. Couse JF, Korach KS. Estrogen receptor null mice: what have we learned and where will they lead us? *Endocr Rev*. 1999;20:358–417.
3. Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci U S A*. 1996;93:5925–30.
4. Hall JM, McDonnell DP. The estrogen receptor beta-isoform (ERbeta) of the human estrogen receptor modulates ERalpha transcriptional activity and is a key regulator of the cellular response to estrogens and antiestrogens. *Endocrinology*. 1999;140:5566–78.
5. Mak P, Leav I, Pursell B, Bae D, Yang X, Taglienti CA, et al. ERbeta impedes prostate cancer EMT by destabilizing HIF-1alpha and inhibiting VEGF-mediated snail nuclear localization: implications for Gleason grading. *Cancer Cell*. 2010;17:319–32.
6. Palmieri C, Saji S, Sakaguchi H, Cheng G, Sunter A, O'Hare MJ, et al. The expression of oestrogen receptor (ER)-beta and its variants, but not ERalpha, in adult human mammary fibroblasts. *J Mol Endocrinol*. 2004;33:35–50.
7. Kregel JH, Hodgins JB, Couse JF, Enmark E, Warner M, Mahler JF, et al. Generation and reproductive phenotypes of mice lacking estrogen receptor beta. *Proc Natl Acad Sci U S A*. 1998;95:15677–82.
8. Ogawa S, Chan J, Chester AE, Gustafsson JA, Korach KS, Pfaff DW. Survival of reproductive behaviors in estrogen receptor beta gene-deficient (betaERKO) male and female mice. *Proc Natl Acad Sci U S A*. 1999;96:12887–92.
9. Massagué J. TGF $\beta$  in cancer. *Cell*. 2008;134:215–30.
10. Clarke RB, Howell A, Potten CS, Anderson E. Dissociation between steroid receptor expression and cell proliferation in the human breast. *Cancer Res*. 1997;57:4987–91.
11. Dubik D, Dembinski TC, Shiu RP. Stimulation of c-myc oncogene expression associated with estrogen-induced proliferation of human breast cancer cells. *Cancer Res*. 1987;47:6517–21.
12. Musgrove EA, Lee CS, Buckley MF, Sutherland RL. Cyclin D1 induction in breast cancer cells shortens G1 and is sufficient for cells arrested in G1 to complete the cell cycle. *Proc Natl Acad Sci U S A*. 1994;91:8022–6.
13. Butt AJ, McNeil CM, Musgrove EA, Sutherland RL. Downstream targets of growth factor and oestrogen signalling and endocrine

- resistance: the potential roles of c-Myc, cyclin D1 and cyclin E. *Endocr Relat Cancer*. 2005;12:S47–59.
14. Lewis-Wambi JS, Jordan VC. Estrogen regulation of apoptosis: how can one hormone stimulate and inhibit? *Breast Cancer Res*. 2009;11:206.
  15. Donovan J, Slingerland J. Transforming growth factor-beta and breast cancer: cell cycle arrest by transforming growth factor-beta and its disruption in cancer. *Breast Cancer Res*. 2000;2:116–24.
  16. Heldin CH, Landström M, Moustakas A. Mechanism of TGF-beta signaling to growth arrest, apoptosis, and epithelial-mesenchymal transition. *Curr Opin Cell Biol*. 2009;21:166–76.
  17. Schmierer B, Hill CS. TGFbeta-SMAD signal transduction: molecular specificity and functional flexibility. *Nat Rev Mol Cell Biol*. 2007;8:970–82.
  18. Lönn P, Morén A, Raja E, Dahl M, Moustakas A. Regulating the stability of TGFbeta receptors and Smads. *Cell Res*. 2009;19:21–35.
  19. Luo K. Ski and SnoN: negative regulators of TGF-beta signaling. *Curr Opin Genet Dev*. 2004;14:65–70.
  20. Hayashi H, Abdollah S, Qiu Y, Cai J, Xu YY, Grinnell BW, et al. The MAD-related protein Smad7 associates with the TGFbeta receptor and functions as an antagonist of TGFbeta signaling. *Cell*. 1997;89:1165–73.
  21. Kavsak P, Rasmussen RK, Causing CG, Bonni S, Zhu H, Thomsen GH, et al. Smad7 binds to Smurf2 to form an E3 ubiquitin ligase that targets the TGF beta receptor for degradation. *Mol Cell*. 2000;6:1365–75.
  22. Ebisawa T, Fukuchi M, Murakami G, Chiba T, Tanaka K, Imamura T, et al. Smurf1 interacts with transforming growth factor-beta type I receptor through Smad7 and induces receptor degradation. *J Biol Chem*. 2001;276:12477–80.
  23. Shi W, Sun C, He B, Xiong W, Shi X, Yao D, et al. GADD34-PP1c recruited by Smad7 dephosphorylates TGFbeta type I receptor. *J Cell Biol*. 2004;164:291–300.
  24. Zhang S, Fei T, Zhang L, Zhang R, Chen F, Ning Y, et al. Smad7 antagonizes transforming growth factor beta signaling in the nucleus by interfering with functional Smad-DNA complex formation. *Mol Cell Biol*. 2007;27:4488–99.
  25. Sun Y, Liu X, Ng-Eaton E, Lodish HF, Weinberg RA. SnoN and Ski protooncoproteins are rapidly degraded in response to transforming growth factor beta signaling. *Proc Natl Acad Sci U S A*. 1999;96:12442–7.
  26. Bonni S, Wang HR, Causing CG, Kavsak P, Stroschein SL, Luo K, et al. TGF-beta induces assembly of a Smad2-Smurf2 ubiquitin ligase complex that targets SnoN for degradation. *Nat Cell Biol*. 2001;3:587–95.
  27. Stroschein SL, Bonni S, Wrana JL, Luo K. Smad3 recruits the anaphase-promoting complex for ubiquitination and degradation of SnoN. *Genes Dev*. 2001;15:2822–36.
  28. Wan Y, Liu X, Kirschner MW. The anaphase-promoting complex mediates TGF-beta signaling by targeting SnoN for destruction. *Mol Cell*. 2001;8:1027–39.
  29. Levy L, Howell M, Das D, Harkin S, Episkopou V, Hill CS. Arkadia activates Smad3/Smad4-dependent transcription by triggering signal-induced SnoN degradation. *Mol Cell Biol*. 2007;27:6068–83.
  30. Band A, Laiho M. Proteasome in the control of TGF- $\beta$  signaling during development and cancer. In Jakowlew S, editor. *Transforming growth factor-beta in cancer therapy*. Totowa, New Jersey; Humana Press; 2008. pp. 243–57.
  31. Stroschein SL, Wang W, Zhou S, Zhou Q, Luo K. Negative feedback regulation of TGF-beta signaling by the SnoN oncoprotein. *Science*. 1999;286:771–4.
  32. Luo K, Stroschein SL, Wang W, Chen D, Martens E, Zhou S, et al. The Ski oncoprotein interacts with the Smad proteins to repress TGFbeta signaling. *Genes Dev*. 1999;13:2196–206.
  33. Nomura T, Khan MM, Kaul SC, Dong HD, Wadhwa R, Colmenares C, et al. Ski is a component of the histone deacetylase complex required for transcriptional repression by Mad and thyroid hormone receptor. *Genes Dev*. 1999;13:412–23.
  34. Suzuki H, Yagi K, Kondo M, Kato M, Miyazono K, Miyazawa K. c-Ski inhibits the TGF-beta signaling pathway through stabilization of inactive Smad complexes on Smad-binding elements. *Oncogene*. 2004;23:5068–76.
  35. Nagano Y, Mavrikis KJ, Lee KL, Fujii T, Koinuma D, Sase H, et al. Arkadia induces degradation of SnoN and c-Ski to enhance transforming growth factor-beta signaling. *J Biol Chem*. 2007;282:20492–501.
  36. Le Scolan E, Zhu Q, Wang L, Bandyopadhyay A, Javelaud D, Mauviel A, et al. Transforming growth factor-beta suppresses the ability of Ski to inhibit tumor metastasis by inducing its degradation. *Cancer Res*. 2008;68:3277–85.
  37. Denissova NG, Liu F. Repression of endogenous Smad7 by Ski. *J Biol Chem*. 2004;279:28143–8.
  38. Tabata T, Kokura K, Ten Dijke P, Ishii S. Ski co-repressor complexes maintain the basal repressed state of the TGF-beta target gene, SMAD7, via HDAC3 and PRMT5. *Genes Cells*. 2009;14:17–28.
  39. Band AM, Björklund M, Laiho M. The phosphatidylinositol 3-kinase/Akt pathway regulates transforming growth factor-beta signaling by destabilizing ski and inducing Smad7. *J Biol Chem*. 2009;284:35441–9.
  40. Zhang YE. Non-Smad pathways in TGF-beta signaling. *Cell Res*. 2009;19:128–39.
  41. Hall JM, McDonnell DP. Coregulators in nuclear estrogen receptor action: from concept to therapeutic targeting. *Mol Interv*. 2005;5:343–57.
  42. Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engström O, et al. Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature*. 1997;389:753–8.
  43. Cheskis BJ, Greger JG, Nagpal S, Freedman LP. Signaling by estrogens. *J Cell Physiol*. 2007;213:610–7.
  44. Métiévier R, Penot G, Hübner MR, Reid G, Brand H, Kos M, et al. Estrogen receptor-alpha directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target promoter. *Cell*. 2003;115:751–63.
  45. Heery DM, Kalkhoven E, Hoare S, Parker MG, Heery DM, Kalkhoven E, et al. A signature motif in transcriptional co-activators mediates binding to nuclear receptors. *Nature*. 1997;387:733–6.
  46. Torchia J, Rose DW, Inostroza J, Kamei Y, Westin S, Glass CK, et al. The transcriptional co-activator p/CIP binds CBP and mediates nuclear-receptor function. *Nature*. 1997;387:677–84.
  47. Vega VB, Lin CY, Lai KS, Kong SL, Xie M, Su X, et al. Multiplatform genome-wide identification and modeling of functional human estrogen receptor binding sites. *Genome Biol*. 2006;7:R82.
  48. Lin CY, Vega VB, Thomsen JS, Zhang T, Kong SL, Xie M, et al. Whole-genome cartography of estrogen receptor alpha binding sites. *PLoS Genet*. 2007;3:e87.
  49. Stender JD, Kim K, Charn TH, Komm B, Chang KC, Kraus WL, et al. Genome-wide analysis of estrogen receptor alpha DNA binding and tethering mechanisms identifies Runx1 as a novel tethering factor in receptor-mediated transcriptional activation. *Mol Cell Biol*. 2010;30:3943–55.
  50. Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key determinant of estrogen receptor function and endocrine response. *Nat Genet*. 2011;43:27–33.
  51. Le Goff P, Montano MM, Schodin DJ, Katzenellenbogen BS. Phosphorylation of the human estrogen receptor. Identification of hormone-regulated sites and examination of their influence on transcriptional activity. *J Biol Chem*. 1994;269:4458–66.

52. Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, et al. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science*. 1995;270:1491–4.
53. Trowbridge JM, Rogatsky I, Garabedian MJ. Regulation of estrogen receptor transcriptional enhancement by the cyclin A/Cdk2 complex. *Proc Natl Acad Sci U S A*. 1997;94:10132–7.
54. Wang C, Fu M, Angeletti RH, Siconolfi-Baez L, Reutens AT, Albanese C, et al. Direct acetylation of the estrogen receptor alpha hinge region by p300 regulates transactivation and hormone sensitivity. *J Biol Chem*. 2001;276:18375–83.
55. Cui Y, Zhang M, Pestell R, Curran EM, Welshons WV, Fuqua SA. Phosphorylation of estrogen receptor alpha blocks its acetylation and regulates estrogen sensitivity. *Cancer Res*. 2004;64:9199–208.
56. Kim MY, Woo EM, Chong YT, Homenko DR, Kraus WL. Acetylation of estrogen receptor alpha by p300 at lysines 266 and 268 enhances the deoxyribonucleic acid binding and transactivation activities of the receptor. *Mol Endocrinol*. 2006;20:1479–93.
57. Subramanian K, Jia D, Kapoor-Vazirani P, Powell DR, Collins RE, Sharma D, et al. Regulation of estrogen receptor alpha by the SET7 lysine methyltransferase. *Mol Cell*. 2008;30:336–47.
58. Levin ER. Plasma membrane estrogen receptors. *Trends Endocrinol Metab*. 2009;20:477–82.
59. Ewan KB, Oketch-Rabah HA, Ravani SA, Shyamala G, Moses HL, Barcellos-Hoff MH, et al. Proliferation of estrogen receptor-alpha-positive mammary epithelial cells is restrained by transforming growth factor-beta1 in adult mice. *Am J Pathol*. 2005;167:409–17.
60. Matsuda T, Yamamoto T, Muraguchi A, Saatcioglu F. Cross-talk between transforming growth factor-beta and estrogen receptor signaling through Smad3. *J Biol Chem*. 2001;276:42908–14.
61. Malek D, Gust R, Kleuser B. 17-Beta-estradiol inhibits transforming-growth-factor-beta-induced MCF-7 cell migration by Smad3-repression. *Eur J Pharmacol*. 2006;534:39–47.
62. Cherlet T, Murphy LC. Estrogen receptors inhibit Smad3 transcriptional activity through Ap-1 transcription factors. *Mol Cell Biochem*. 2007;306:33–42.
63. Ito I, Hanyu A, Wayama M, Goto N, Katsuno Y, Kawasaki S, et al. Estrogen inhibits transforming growth factor beta signaling by promoting Smad2/3 degradation. *J Biol Chem*. 2010;285:14747–55.
64. Burdette JE, Woodruff TK. Activin and estrogen crosstalk regulates transcription in human breast cancer cells. *Endocr Relat Cancer*. 2007;14:679–89.
65. Dhasarathy A, Kajita M, Wade PA. The transcription factor snail mediates epithelial to mesenchymal transitions by repression of estrogen receptor-alpha. *Mol Endocrinol*. 2007;21:2907–18.
66. Buck MB, Pfizenmaier K, Knabbe C. Antiestrogens induce growth inhibition by sequential activation of p38 mitogen-activated protein kinase and transforming growth factor-beta pathways in human breast cancer cells. *Mol Endocrinol*. 2004;18:1643–57.
67. Chen H, Tritton TR, Kenny N, Absher M, Chiu JF. Tamoxifen induces TGF-beta 1 activity and apoptosis of human MCF-7 breast cancer cells in vitro. *J Cell Biochem*. 1996;61:9–17.
68. Buck MB, Knabbe C. TGF-beta signaling in breast cancer. *Ann N Y Acad Sci*. 2006;1089:119–26.
69. Bieri B, Moses HL. Gain or loss of TGFbeta signaling in mammary carcinoma cells can promote metastasis. *Cell Cycle*. 2009;8:3319–27.
70. Bieri B, Chung CH, Parker JS, Stover DG, Cheng N, Chytil A, et al. Abrogation of TGF-beta signaling enhances chemokine production and correlates with prognosis in human breast cancer. *J Clin Invest*. 2009;119:1571–82.
71. Joffroy CM, Buck MB, Stope MB, Popp SL, Pfizenmaier K, Knabbe C. Antiestrogens induce transforming growth factor beta-mediated immunosuppression in breast cancer. *Cancer Res*. 2010;70:1314–22.
72. Wu L, Wu Y, Gathings B, Wan M, Li X, Grizzle W, et al. Smad4 as a transcription corepressor for estrogen receptor alpha. *J Biol Chem*. 2003;278:15192–200.
73. Ren Y, Wu L, Frost AR, Grizzle W, Cao X, Wan M. Dual effects of TGF-beta on ERalpha-mediated estrogenic transcriptional activity in breast cancer. *Mol Cancer*. 2009;8:111.
74. Li Q, Wu L, Oelschlager DK, Wan M, Stockard CR, Grizzle WE, et al. Smad4 inhibits tumor growth by inducing apoptosis in estrogen receptor-alpha-positive breast cancer cells. *J Biol Chem*. 2005;280:27022–8.
75. Deheuninck J, Luo K. Ski and SnoN, potent negative regulators of TGF-beta signaling. *Cell Res*. 2009;19:47–57.
76. Jahchan NS, Luo K. SnoN in mammalian development, function and diseases. *Curr Opin Pharmacol*. 2010;10:670–5.
77. Edmiston JS, Yeudall WA, Chung TD, Lebman DA. Inability of transforming growth factor-beta to cause SnoN degradation leads to resistance to transforming growth factor-beta-induced growth arrest in esophageal cancer cells. *Cancer Res*. 2005;65:4782–8.
78. Zhang F, Lundin M, Ristimäki A, Heikkilä P, Lundin J, Isola J, et al. Ski-related novel protein N (SnoN), a negative controller of transforming growth factor-beta signaling, is a prognostic marker in estrogen receptor-positive breast carcinomas. *Cancer Res*. 2003;63:5005–10.
79. Zhu Q, Krakowski AR, Dunham EE, Wang L, Bandyopadhyay A, Berdeaux R, et al. Dual role of SnoN in mammalian tumorigenesis. *Mol Cell Biol*. 2007;27:324–39.
80. Jahchan NS, You YH, Muller WJ, Luo K. Transforming growth factor-beta regulator SnoN modulates mammary gland branching morphogenesis, postlactational involution, and mammary tumorigenesis. *Cancer Res*. 2010;70:4204–13.
81. Shinagawa T, Dong HD, Xu M, Maekawa T, Ishii S. The sno gene, which encodes a component of the histone deacetylase complex, acts as a tumor suppressor in mice. *EMBO J*. 2000;19:2280–91.