

Phosphatidylinositol 3-Kinase and Antiestrogen Resistance in Breast Cancer

Todd W. Miller, Justin M. Balko, and Carlos L. Arteaga

All authors: Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN.

Submitted April 22, 2011; accepted August 4, 2011; published online ahead of print at www.jco.org on October 17, 2011.

Supported by National Institutes of Health Grant No. K99CA142899 (T.W.M.); Breast Cancer Specialized Program of Research Excellence Grant No. P50CA98131; Vanderbilt-Ingram Cancer Center Support Grant No. P30CA68485; a grant from the Breast Cancer Research Foundation (C.L.A.); American Cancer Society Clinical Research Professorship Grant No. CRP-07-234 (C.L.A.); the Lee Jeans Translational Breast Cancer Research Program (C.L.A.); and Stand Up to Cancer/American Association for Cancer Research Dream Team Translational Cancer Research Grant No. SU2C-AACR-DT0209 (C.L.A.).

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Clinical Trials repository link available on JCO.org.

Corresponding author: Carlos L. Arteaga, MD, Division of Hematology-Oncology, VUMC, 2200 Pierce Ave, 777 PRB, Nashville, TN 37232-6307; e-mail: carlos.arteaga@vanderbilt.edu.

© 2011 by American Society of Clinical Oncology

0732-183X/11/2933-4452/\$20.00

DOI: 10.1200/JCO.2010.34.4879

ABSTRACT

Although antiestrogen therapies targeting estrogen receptor (ER) α signaling prevent disease recurrence in the majority of patients with hormone-dependent breast cancer, a significant fraction of patients exhibit de novo or acquired resistance. Currently, the only accepted mechanism linked with endocrine resistance is amplification or overexpression of the *ERBB2* (human epidermal growth factor receptor 2 [*HER2*]) proto-oncogene. Experimental and clinical evidence suggests that hyperactivation of the phosphatidylinositol 3-kinase (PI3K) pathway, the most frequently mutated pathway in breast cancer, promotes antiestrogen resistance. PI3K is a major signaling hub downstream of *HER2* and other receptor tyrosine kinases. PI3K activates several molecules involved in cell-cycle progression and survival, and in ER-positive breast cancer cells, it promotes estrogen-dependent and -independent ER transcriptional activity. Preclinical tumor models of antiestrogen-resistant breast cancer often remain sensitive to estrogens and PI3K inhibition, suggesting that simultaneous targeting of the PI3K and ER pathways may be most effective. Herein, we review alterations in the PI3K pathway associated with resistance to endocrine therapy, the state of clinical development of PI3K inhibitors, and strategies for the clinical investigation of such drugs in hormone receptor-positive breast cancer.

J Clin Oncol 29:4452-4461. © 2011 by American Society of Clinical Oncology

INTRODUCTION

Approximately 75% of breast cancers express estrogen receptor (ER) α and/or progesterone receptor (PR). Hormone receptor expression typically indicates a degree of estrogen dependence for cancer cell growth. Treatment for these patients inhibits ER function either by antagonizing ligand binding to ER (tamoxifen and other selective ER modulators [SERMs]), downregulating ER (fulvestrant), or blocking estrogen biosynthesis (aromatase inhibitors [AIs]). Although endocrine therapies have changed the natural history of hormone-dependent breast cancer, many tumors exhibit de novo or acquired resistance. For example, more than 30% of patients with early ER-positive breast cancer relapse within 15 years after adjuvant therapy with tamoxifen, and 17% of patients treated with an AI relapse within 9 years.^{1,2} An accepted mechanism of resistance to endocrine therapy involves overexpression of the *ERBB2* (*HER2*) proto-oncogene.³⁻⁵ However, less than 10% of ER-positive breast cancers overexpress *HER2*, suggesting that for most ER-positive breast cancers, mechanisms of escape from endocrine therapy remain to be elucidated.

Tamoxifen has been a standard treatment for ER-positive breast cancer for more than 30 years.

This SERM has dual agonistic/antagonistic effects on ER transcription and breast cancer cell growth.^{6,7} In contrast, AIs induce estrogen deprivation without agonistic effects on ER. AIs are clinically equivalent if not modestly superior to tamoxifen.^{2,8} Data on the clinical activity of fulvestrant as first-line therapy in the metastatic setting are limited but suggest it may be superior to AIs despite its inability to completely downregulate ER in patients' tumors.⁹⁻¹¹ It is unclear whether mechanisms of antiestrogen resistance are common to SERMs, AIs, and ER downregulators. Studies in human cell lines and xenografts have shown that growth factor receptor signaling pathways, particularly those that converge on phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase (MAPK/ERK), can mediate resistance to all forms of endocrine therapy. PI3K is the most frequently altered pathway in breast cancer, with mutation and/or amplification of the genes encoding the PI3K catalytic subunits p110 α (*PIK3CA*) and p110 β (*PIK3CB*); the PI3K regulatory subunit p85 α (*PIK3R1*); receptor tyrosine kinases (RTKs) such as *HER2* (*ERBB2*) and fibroblast growth factor receptor 1 (*FGFR1*); K-Ras; PI3K effectors AKT1, AKT2, and PDK1; and loss of the lipid phosphatases phosphatase and tensin homolog (PTEN) and INPP4B (Table 1).

Table 1. PI3K Pathway Alterations Associated With Response to Endocrine Therapy

Gene	Protein	Aberration	Effect on Signaling	Frequency	Patient Prognosis	Reference
<i>ERBB2</i>	HER2	Gene amplification or overexpression	Hyperactivation of ErbB2 signaling (PI3K, MEK)	Approximately 10% of ER-positive tumors	Correlates with worse outcome	Ellis et al, ³ Arpino et al, ⁴ De Laurentiis et al ⁵
<i>PTEN</i>		Loss-of-function mutation or reduced expression	Hyperactivation of PI3K signaling	37% to 44% of ER-positive tumors	No consistent correlation	Pérez-Tenorio et al, ¹² Saal et al, ¹³ Shoman et al ¹⁴
<i>PIK3CA</i>	p110 α , PI3K	Activating mutation	Hyperactivation of PI3K signaling	28% to 47% of ER-positive tumors	No correlation or correlation with better outcome	Pérez-Tenorio et al, ¹² Baselga et al, ¹⁵ Stemke-Hale et al, ¹⁶ Ellis et al, ¹⁷ Campbell et al ¹⁸
<i>PIK3CB</i>	p110 β , PI3K	Amplification	Unknown	5% of all cases	Unknown	Crowder et al ¹⁹
<i>IGF1R</i> , <i>INSR</i>	IGF-1R, InsR	Receptor activation	Activates IGF-1R/InsR signaling (PI3K, MEK)	48% of ER-positive tumors	Unknown	Law et al ²⁰
<i>FGFR1</i>		Amplification	Hyperactivation of FGFR signaling (PI3K, MEK)	11.6% of ER-positive tumors	Correlates with shorter RFS	Turner et al ²¹
<i>RPS6K1</i>	p70S6K	Amplification	Activates mTORC1, protein translation	8.8% to 12.5% of all tumors	Unknown	Monni et al ²²
<i>INPP4B</i>		Reduced expression or genomic loss	Hyperactivation of PI3K signaling	8.4% to 37.7% of ER-positive tumors	Correlates with worse outcome	Gewinner et al, ²³ Fedele et al ²⁴
<i>PIK3R1</i>	p85 α , PI3K	Inactivating mutation	Derepression of catalytic activity of p110 α	2% of all tumors	Unknown	Jaiswal et al ²⁵
<i>AKT1</i>		Activating mutation	Hyperactivation of AKT signaling	2.6% to 3.8% of ER-positive tumors	Unknown	Stemke-Hale et al, ¹⁶ Loi et al, ²⁶ Carpten et al ²⁷
<i>AKT2</i>		Amplification	Hyperactivation of AKT signaling	2.8% of all tumors	Unknown	Bellacosa et al ²⁸
<i>EGFR</i>		Amplification	Hyperactivation of EGFR signaling (PI3K, MEK)	0.5% of ER-positive tumors	Unknown	Al-Kuraya et al ²⁹
<i>PDK1</i>		Amplification or overexpression	Hyperactivation of PDK1 signaling (AKT, mTORC1)	21% of all tumors	Unknown	Maurer et al ³⁰
<i>KRAS</i>		Activating mutation	Hyperactivation of Ras signaling	4% to 6% of all tumors	Unknown	Rochlitz et al, ³¹ Di Nicolantonio et al, ³²

Abbreviations: EGFR, epidermal growth factor receptor; ER, estrogen receptor; FGFR, fibroblast growth factor receptor 1; HER2, human epidermal growth factor receptor 2; IGF-1R, insulin-like growth factor-1 receptor; InsR, insulin receptor; mTORC1, mammalian target of rapamycin complex 1; PI3K, phosphatidylinositol 3-kinase; RFS, relapse-free survival.

CROSSTALK BETWEEN ER AND PI3K PATHWAYS

PI3K is activated by growth factor RTKs and G-protein–coupled receptors (GPCRs). PI3K phosphorylates phosphatidylinositol 4,5-bisphosphate (PIP₂) to produce phosphatidylinositol 3,4,5-trisphosphate (PIP₃).³³ In turn, PIP₃ recruits several pleckstrin homology (PH) domain–containing proteins to the plasma membrane such as PDK1, serine/threonine-protein kinase, and AKT, which on activation drive cell-cycle progression and survival.^{34,35} Negative regulation of this pathway is conferred by PTEN and INPP4B, which dephosphorylate PIP₃ and PIP₂, respectively.^{23,36} AKT activates the mammalian target of rapamycin (mTOR) –containing complex 1 (TORC1),³⁷ which regulates protein synthesis. mTOR is also part of another complex (ie, TORC2), which lies upstream of AKT.³⁸

In addition to its pro-survival and growth-promoting roles, the PI3K pathway interacts with ER directly and indirectly (Fig 1). ER

phosphorylation at Ser₁₆₇ by AKT or p70S6K increases estrogen-induced, tamoxifen-induced, and ligand-independent ER transcriptional activity.^{39,40} Additionally, PI3K and Ras promote c-Jun phosphorylation.^{41–45} c-Jun complexes with c-Fos to form the AP-1 complex, which cooperates with ER transcription.^{46–48} Other oncogenic kinase pathways (eg, MAPK, protein kinase C) also contribute to the modulation of ER and transcription cofactors.⁴⁹

The activation of ER by growth factor RTK signaling is reciprocated in a feed-forward fashion, whereby ER promotes the transcription of genes encoding ligands, RTKs, and signaling adaptors (Fig 1). Clinical evidence also suggests that ER function may sustain PI3K pathway activation in breast cancer cells. Neoadjuvant treatment of patients with ER-positive breast cancer with the AI letrozole reduces P-AKT_{S473} and P-mTOR_{S2448} tumor levels (markers of PI3K activity), which correlates with clinical response and improved disease outcome.⁵⁰ Another study detected a reduction in P-S6 levels (marker of




Fig 1. Reciprocal crosstalk between estrogen receptor (ER) α and growth factor receptor signaling pathways. Receptor tyrosine kinases (RTKs) and G-protein-coupled receptors activate phosphatidylinositol 3-kinase (PI3K; blue) and MEK signaling pathways. These signal transducers can then phosphorylate ER (green) and/or coactivators and corepressors to modulate ER transcriptional activity not necessarily dependent on ER ligands. In turn, ER transcribes genes encoding components of growth factor signaling pathways, thus completing signaling cycle of RTKs to ER to RTKs. ER also complexes with RTKs and Src to rapidly induce nongenomic signaling. ER-interacting proteins shown in color. EGFR, epidermal growth factor receptor; IGF-1R, insulin-like growth factor-1 receptor; IGFBP, insulin-like growth factor binding protein; IRS-1, insulin receptor substrate 1; JNK, c-Jun N-terminal kinase; PTEN, phosphatase and tensin homolog; SGK3, serum/glucocorticoid-regulated kinase 3; TGF α , transforming growth factor alpha; TORC1, target of rapamycin complex 1.

TORC1 activity) after neoadjuvant letrozole.¹⁵ Hence, estrogen deprivation may suppress ER-positive breast cancer cell growth in part by decreasing PI3K/AKT/mTOR signaling.

Experimentally, PI3K pathway activation has been causally associated with de novo and acquired resistance to endocrine therapy. RNAi-mediated knockdown of *PTEN* and overexpression of oncogenes that activate PI3K/AKT signaling (eg, *HER2*, type 1 insulin-like growth factor receptor (*IGF1R*), activated mutant *AKT1*) confer resistance to tamoxifen, fulvestrant, and estrogen deprivation in ER-positive breast cancer cells. In most such models, inhibition of PI3K has reversed antiestrogen resistance.^{7,39,51} In another example, tamoxifen-resistant ER-positive MCF-7 breast cancer xenografts exhibited increased expression of IGF-1R, HER2, and epidermal growth factor receptor (EGFR).⁵² One model of resistance to estrogen deprivation employs overexpression of aromatase in MCF-7 cells followed by selection with an AI. AI-resistant MCF-7/aromatase cells and xenografts have exhibited higher levels of HER2 and the EGFR ligand amphiregulin along with lower levels of ER.^{53,54} We reported that four of four long-term estrogen-deprived (LTED) ER-positive breast cancer cell lines exhibited amplification of *PI3K/AKT/mTOR* signaling, and three of four lines showed hyperactivation of IGF-1R and/or InsR.⁵⁵ Chronic exposure of MCF-7 cells and xenografts to fulvestrant has been shown to elicit similar changes.⁵⁶⁻⁵⁸

Emerging evidence also implicates nongenomic, estrogen-induced signaling in the activation of PI3K. Estrogen stimulation rapidly initiates intracellular kinase signaling, including activation of IGF-1R/InsR, EGFR, Src, PI3K, and MEK (Fig 1).^{51,59-61} This implies that a non-nuclear ER binds estrogen and transmits signals to membrane and cytosolic kinases. Several membrane ERs have been identified, including ER α , splice variants of ER α (ER-36, ER-46), ER β , and GPCR 30.⁶²⁻⁶⁴ Although ER α localizes to the plasma membrane in cultured cells, this has not been demonstrated in human tumors.⁶⁵ Membrane ER might activate oncogenic kinases to promote endocrine resistance; however, these mechanisms remain to be shown clinically.

PI3K pathway activation is required for growth of breast cancer cells resistant to endocrine therapy. Growth of four of four LTED cell lines in the absence of estrogen is inhibited by treatment with the PI3K/mTOR inhibitor BEZ235 or the TORC1 inhibitor everolimus.⁵⁵ Treatment of mice bearing estrogen-independent MCF-7 xenografts with the pan-PI3K inhibitor BKM120, or bearing letrozole-resistant MCF-7/aromatase xenografts with wortmannin, has been shown to slow tumor growth (unpublished data).^{66,67} Also, treatment of MCF-7 and MCF-7/LTED cells with the Ras inhibitor farnesylthiosalicylic acid decreases mTOR signaling and hormone-independent growth.⁶⁸ Interestingly, a recent study showed that exogenous estrogen prevents

the apoptotic effects of BEZ235 and RNAi-mediated silencing of PI3K in ER-positive cells.¹⁹ Most breast cancers that adapt to antiestrogen therapy retain ER and, presumably, estrogen sensitivity. These data imply that treatment of patients harboring ER-positive breast cancers with a PI3K-targeted therapy alone (without endocrine agent) might be insufficient to maximally inhibit tumor growth.

ASSOCIATION OF PI3K HYPERACTIVATION WITH RESISTANCE TO ENDOCRINE THERAPY

Gain-of-function oncogenes and/or loss of tumor suppressors in breast cancer cells may confer antiestrogen resistance via activation of PI3K. For example, HER2 overexpression predicts weaker response to neoadjuvant AIs or tamoxifen and worse outcome after adjuvant endocrine therapy compared with ER-positive/HER2-negative breast cancers.³⁻⁵ Patients with FGFR1-overexpressing ER-positive tumors exhibit a shorter relapse-free survival (RFS) after adjuvant tamoxifen.²¹ Patients with ER-positive/INPP4B-deficient tumors show worse survival compared with patients with ER-positive/INPP4B-positive tumors.²³ Although loss-of-function mutations in *PTEN* are rare in ER-positive breast cancer, immunohistochemical (IHC) studies have reported a wide range of PTEN loss but found no association of PTEN level with outcome after tamoxifen therapy. Whether other mutations in the PI3K pathway correlate with antiestrogen resistance remains to be determined.

Point mutations in *PIK3CA*, the gene encoding the p110 α catalytic subunit of PI3K, are the most common genetic alterations of this pathway in breast cancer. Up to 80% of *PIK3CA* mutations occur in hotspots within the helical (E542K and E545K) and kinase (H1047R) domains of p110 α . They increase PI3K activity, induce cellular transformation in vitro and tumorigenicity in vivo when overexpressed in human mammary epithelial cells, and induce mammary tumor formation in transgenic mice.⁶⁹⁻⁷² Although such mutations occur in 28% to 47% of ER-positive breast cancers, their clinical significance remains unclear. In retrospective studies, *PIK3CA* mutations in primary ER-positive tumors have correlated with good long-term outcome.^{12,15-18,26} In one report, *PIK3CA* mutations were linked with lower activation of PI3K (assessed by P-AKT) compared with PTEN deficiency in breast tumors.¹⁶ In another study, a *PIK3CA*-mutant gene expression signature was associated with low P-AKT, low TORC1 signaling, and good outcome after adjuvant tamoxifen.²⁶ This association of *PIK3CA*-mutant status and good prognosis in ER-positive disease does not negate the possibility that combinations of antiestrogens and PI3K pathway inhibitors would be clinically more effective than antiestrogens alone in such tumors. Indeed, a recent analysis of a cohort of 217 patients with multiple solid tumor types enrolled onto phase I trials with PI3K/AKT/mTOR inhibitors showed a higher response rate among patients with *PIK3CA*-mutant versus *PIK3CA*-wild-type cancers.⁷³

Understanding the relationship between *PIK3CA* mutations and endocrine resistance may be confounded by evidence suggesting that these genetic alterations may arise late in tumor development. For example, *PIK3CA* mutation status is discordant between invasive carcinoma and ductal carcinoma in situ in 33% of patient cases, between primary breast tumors and synchronous lymph node metastases in 13% of patient cases, between primary tumors and asynchronous metastases in 18% to 33% of patient cases, and even within microdissected regions of the same tumor.⁷⁴⁻⁷⁶

In addition to mutational analyses, tumor protein and gene expression profiling of PI3K pathway activation may provide a biomarker to identify patients with antiestrogen-resistant tumors. For example, a gene expression signature of PTEN loss, derived from a comparison of PTEN-positive and -negative tumors by IHC, was predictive of poor RFS after tamoxifen, whereas PTEN IHC status alone was not.¹³ A gene expression signature of PI3K activation, based on levels of phosphoprotein markers (eg, P-AKT, P-p70S6K) in ER-positive tumors, was enriched in luminal B breast cancers.⁷⁷ This suggests that luminal B tumors have higher PI3K activity, which may contribute to their lower response to antiestrogens compared with luminal A tumors.⁷⁸ Similarly, we identified a tumor protein signature of PI3K pathway activation that predicts poor outcome after adjuvant endocrine therapy.⁵⁵ Therefore, signatures of PI3K activation may complement mutational analyses for the identification of PI3K-driven tumors.

RATIONALE FOR COMBINED USE OF ER AND PI3K INHIBITORS

One of the first PI3K pathway-targeted drugs to be tested in combination with endocrine therapy was the TORC1 inhibitor everolimus. Patients with ER-positive tumors were randomly assigned to neoadjuvant letrozole with or without everolimus for 4 months before surgery. The combination therapy induced a higher clinical response and greater suppression of tumor cell proliferation compared with letrozole alone.¹⁵ In another study, patients with advanced disease who had progressed while receiving an AI were randomly assigned to tamoxifen with or without everolimus. Patients receiving tamoxifen plus everolimus showed a significantly improved rate of clinical benefit, time to progression, and disease-free survival compared with women receiving tamoxifen alone.⁷⁹

Although combinations containing TORC1 inhibitors have shown clinical activity in breast cancer, inhibition of TORC1 relieves negative feedback on activators of PI3K (eg, IGF-1R, IRS-1, HER3), which in turn promote cell survival.⁸⁰⁻⁸² These results suggest that direct inhibitors of PI3K may be more effective. However, inhibition of PI3K or AKT also results in feedback upregulation/activation of several RTKs, which, by providing an upstream input to PI3K, may counteract drug action.⁸³⁻⁸⁶ These data suggest that PI3K/AKT/TORC1 inhibitors should be combined with RTK inhibitors to induce an optimal antitumor effect. Consistent with this notion, studies with human xenografts have shown that combinations of inhibitors targeting HER2 and PI3K, HER2 and AKT, HER2 and TORC1, or EGFR and AKT are each superior to single-agent treatments.^{80,83-85}

Additional rationales for combined inhibition of PI3K and ER come from studies in HER2-positive breast cancer. HER2 overexpression, a potent activator of PI3K, confers endocrine resistance.⁸⁷⁻⁹⁰ Preclinical and clinical data suggest that ER-positive breast cancer cells initially inhibited by tamoxifen or estrogen deprivation can upregulate HER2 to bypass ER blockade.^{88,91} The interdependence of these pathways is highlighted by examples in which HER2 inhibition with trastuzumab or the tyrosine kinase inhibitor lapatinib restores or upregulates ER levels or transcriptional activity in breast cancer cells and patient tumors.^{92,93} Furthermore, treatment with AIs or downregulation of ER with fulvestrant or RNAi has inhibited the growth of HER2-positive tumors or cells that had progressed with trastuzumab or lapatinib.^{92,93} These data suggest that combined inhibition of ER

and HER2 signaling may provide more effective control of ER-positive/HER2-positive tumors. This inference is supported by results from two clinical trials. In the TAnDEM (Trastuzumab in Dual Human Epidermal Growth Factor Receptor Type 2 [HER2] ER + Metastatic Breast Cancer) study, 207 patients with ER-positive/HER2-positive metastatic breast cancer were randomly assigned to the AI anastrozole with or without trastuzumab. A second study randomly assigned 219 patients with ER-positive/HER2-positive metastatic breast cancer to letrozole with or without lapatinib. In both trials, progression-free survival and clinical benefit were superior in the combination arm compared with the AI-alone arm, suggesting that both HER2 and ER should be simultaneously targeted for maximal therapeutic efficacy.^{94,95} Whether combined inhibition of PI3K and ER is equally effective against ER-positive/HER2-positive tumors remains to be determined.

PI3K PATHWAY INHIBITORS IN CLINICAL DEVELOPMENT

Several drugs targeting multiple levels of the PI3K network (ie, PI3K, AKT, mTOR) are in clinical development. Class IA PI3K isoforms are heterodimeric lipid kinases that contain a p110 catalytic subunit and p85 regulatory subunit. Three genes (ie, *PIK3CA*, *PIK3CB*, and *PIK3CD*) encode the homologous p110 α , p110 β , and p110 δ isozymes, respectively.^{96,97} p110 δ expression is largely restricted to immune and hematopoietic cells, whereas p110 α and p110 β are ubiquitously expressed.⁹⁸ p110 α is essential for signaling and growth of tumors driven by *PIK3CA* mutations, RTKs, and/or mutant Ras, whereas p110 β lies downstream of GPCRs and has been shown to mediate tumorigenesis in PTEN-deficient cells.⁹⁹ A number of ATP-mimetics that bind competitively and reversibly to the ATP-binding pocket of p110 are in early clinical development.^{100,101} These include the pan-PI3K inhibitors BKM120, XL-147, PX-866, PKI-587, and GDC-0941; p110 α -specific inhibitors BYL719, GDC-0032, and INK-1117; p110 δ -specific inhibitor CAL-101; and dual PI3K/mTOR inhibitors BEZ235, BGT226, PF-4691502, GDC-0980, and XL-765.

The pan-PI3K and p110 α -specific inhibitors are equally potent against oncogenic p110 α mutants.¹⁰²⁻¹⁰⁴ The rationale for the development of isozyme-specific antagonists is to allow higher doses of anti-p110 α and anti-p110 β drugs to be delivered without incurring adverse effects caused by pan-PI3K inhibitors. Interim results from a phase I trial with the p110 δ -specific inhibitor CAL-101 in patients with hematologic malignancies showed that treatment reduced P-AKT levels by more than 90% in peripheral blood lymphocytes and induced objective clinical responses.¹⁰⁵ Recently completed phase I trials with BKM120, BEZ235, and XL-147 showed that treatment partially inhibited PI3K as measured by levels of P-S6 and P-AKT in patients' skin or tumors and [¹⁸F]fluorodeoxyglucose uptake measured by PET. Main toxicities were rash, hyperglycemia, diarrhea, fatigue, and mood alterations. Few clinical responses were observed in patients with and without detectable PI3K pathway mutations, although screening for genetic lesions was not comprehensive.¹⁰⁶⁻¹⁰⁸

Both allosteric and ATP-competitive pan-inhibitors of the three isoforms of AKT are being developed. AZD5363, GDC-0068, GSK2141795, and GSK690693 are ATP-competitive compounds that have shown antitumor activity in preclinical models and recently entered phase I trials.^{109,110} Allosteric inhibitors such as MK-2206 bind to the AKT PH domain and/or hinge region to promote a con-

formation incapable of membrane localization.¹¹¹ MK-2206 inhibits AKT signaling in vivo and suppresses growth of breast cancer xenografts harboring *PIK3CA* mutations or *ERBB2* amplification.¹¹² Phase I data have shown that treatment with MK-2206 decreases levels of P-AKT, P-proline-rich AKT substrate 40, and P-GSK3 β in tumor cells, peripheral blood mononuclear cells, and hair follicles.^{113,114}

The mTOR kinase is a component of PI3K-driven oncogenesis that functions within two signaling complexes: TORC1 and TORC2.^{37,38} The macrolide rapamycin and its analogs complex with FK506-binding protein (FKBP12), which then binds to mTOR and inhibits the kinase activity of TORC1 but not TORC2.³⁸ The formulation problems of rapamycin prompted the development of analogs such as CCI-779 (temsirolimus), RAD001 (everolimus), AP-23573 (deferolimus), and MK-8669 (ridaferolimus). These rapalogs have shown cytostatic activity in preclinical models and clinical trials, particularly in patients with renal cell cancer and in those with mutations in the tuberous sclerosis complex (upstream of TORC1) who harbor renal angioliopomas. Compounds that target the ATP-binding cleft of mTOR (ie, OSI-027, AZD8055, INK-128), and are thus active against both TORC1 and TORC2, are also in phase I trials.¹¹⁵

CLINICAL INVESTIGATION OF PI3K PATHWAY INHIBITORS IN ER-POSITIVE BREAST CANCER

The somatic alterations described (ie, *PIK3CA* and *AKT1* mutations, PTEN and INPP4B loss, *PIK3CB* and *AKT2* amplification, and so on) identify cancers with aberrant activation of and potential dependence on the PI3K pathway. This is an important consideration for the selection of patients into trials with PI3K inhibitors. Like mutant PI3K, other somatic mutations in tumors have revealed molecules critical for cancer survival and progression. Pharmacologic targeting of these mutants has resulted in remarkable clinical responses in patients bearing tumors with such mutant oncogenes. Examples include imatinib and dasatinib in chronic myelogenous leukemia harboring the *BCR-ABL* oncogene, EGFR TKIs gefitinib and erlotinib in lung cancers with *EGFR*-activating mutations, HER2 antagonists trastuzumab and lapatinib in breast cancers with *HER2* gene amplification, and Raf inhibitors against metastatic melanomas containing B-Raf-activating mutations.¹¹⁶

As with other targeted therapies, only a fraction of patients with tumors containing PI3K pathway mutations will likely benefit from single-agent PI3K inhibitors. There is increasing agreement that initial phase II efficacy studies with PI3K inhibitors in patients with advanced disease should be enriched with, if not limited to, patients harboring mutations in this pathway. However, testing these drugs in single-arm phase II trials in patients with metastatic cancer with PI3K pathway alterations is intrinsically problematic because: first, the difficulty of obtaining biopsies from metastatic sites, and second, the limitations of assessing tumor response as a meaningful clinical end point in the absence of a placebo control arm. We will discuss alternative approaches that may address these issues.

PRESURGICAL AND NEOADJUVANT CLINICAL TRIALS

There are examples of short-term, pharmacodynamic trials providing information that can be later used for patient selection into trials with

novel targeted therapies such as PI3K antagonists. In ER-positive breast cancer, data from presurgical studies suggest that such trials can be used to predict longer-term outcome after adjuvant endocrine therapy. Dowsett et al¹¹⁷ reported that patients with ER-positive tumors with high Ki67 scores after 2 weeks of neoadjuvant antiestrogen therapy had shorter RFS compared with those with low Ki67 scores. Ellis et al¹¹⁸ found that Ki67 scores and ER status after 3 to 4 months of neoadjuvant endocrine therapy were predictive of RFS after adjuvant tamoxifen. Furthermore, ER-positive/HER2-negative tumors showed a reduction in Ki67 in response to neoadjuvant letrozole, whereas ER-positive/HER2-positive tumors did not.³ Therefore, presurgical evaluation of molecular markers after short-term treatment may be valuable to predict long-term patient benefit. However, no single biomarker has been shown to accurately predict disease outcome in individual patients. Ki67 score was found to vary between biopsies from the same breast tumor such that a change greater than 36% to 50% in the post-treatment compared with pretreatment biopsy would be required to detect a potentially informative difference.^{119,120} Furthermore, identifying patients who would benefit from neoadjuvant therapy using Ki67 score is challenging in tumors with rather low baseline cell proliferation.¹²⁰

Although some PI3K pathway mutations (eg, *HER2*, *FGFR1* amplification) have been linked with antiestrogen resistance, the role of *PIK3CA* mutations is less clear.^{3-5,21} Neoadjuvant trials provide a setting to investigate this role. For example, in a neoadjuvant study of letrozole with or without everolimus, *PIK3CA*^{exon9} mutations were associated with a statistically lower reduction in cell proliferation in response to letrozole, as measured by a change in the Ki67 IHC score on day 15, compared with *PIK3CA*-wild-type tumors.¹⁵ We have observed similar preliminary data in a presurgical trial of short-term

letrozole (10 to 21 days) in patients with ER-positive/HER2-negative breast cancer (NCT00651976). In this study, analysis of the first 21 patients showed that *PIK3CA*-mutant tumors exhibited a statistically lower reduction in Ki67 score compared with tumors with wild-type *PIK3CA* (data not shown). However, another study in which tumor cell proliferation was assessed at baseline and after 4 months of antiestrogen therapy found no association between *PIK3CA* status and drug-induced change in Ki67 score.¹⁷ This discrepancy may be the result of the different timing (2 to 3 weeks v 4 months) of the biopsies from which cell proliferation was assessed.

RANDOMIZED TRIALS OF ANTIESTROGENS WITH OR WITHOUT PI3K PATHWAY INHIBITORS

Although PI3K pathway activation may confer antiestrogen resistance, breast tumors that acquire hormone independence may still be stimulated by (host) estradiol. Thus, we speculate that PI3K inhibitors may have limited efficacy against ER-positive breast cancers as single agents. Because antiestrogen therapies are effective against a large fraction of ER-positive tumors as single agents, there is a need for randomized clinical trials of standard ER-targeted therapies versus combinations targeting both the ER and PI3K pathways.

In Table 2, we summarize the randomized trials in which inhibitors of TORC1, HER2, EGFR, IGF-1R, protein kinase C- β /PDK1/p70S6K, and farnesyltransferase have been combined with endocrine therapy. The molecular targets of these drugs rely on PI3K and have been linked to endocrine resistance. Neoadjuvant treatment with letrozole and the TORC1 inhibitor everolimus more effectively suppressed tumor cell proliferation and increased clinical response compared with letrozole alone in patients with early-stage ER-positive

Table 2. Prospective Randomized Trials Testing Combinations of PI3K Pathway Inhibitors and Endocrine Therapies

Kinase Target	Trial Design	Phase	Reference
mTOR	Letrozole \pm everolimus in patients with early-stage ER-positive breast cancer	II	Baselga et al ¹⁵
	Exemestane \pm everolimus in ER-positive metastatic breast cancer after progression while receiving another AI	III	NCT00863655 (BOLERO-2; ongoing)
	Tamoxifen \pm everolimus in ER-positive metastatic breast cancer after progression while receiving AI	II	TAMRAD; Bachelot et al ⁷⁹
HER2	Anastrozole \pm trastuzumab in patients with ER-positive/HER2-positive metastatic breast cancer	III	Kaufman et al ⁹⁵
HER2/EGFR	Letrozole \pm lapatinib in ER-positive metastatic breast cancer	III	Johnston et al ⁹⁴
	AI \pm lapatinib or AI + fulvestrant \pm lapatinib in ER-positive metastatic breast cancer after progression while receiving AI	III	NCT00688194 (ongoing)
	AI \pm trastuzumab or lapatinib or both in patients with ER-positive/HER2-positive metastatic breast cancer	III	NCT01160211 (not yet open)
EGFR	Anastrozole \pm gefitinib in early-stage ER-positive breast cancer	II	Smith et al ¹²¹
	Anastrozole \pm gefitinib in metastatic ER-positive patients	II	Cristofanilli et al ¹²²
PKC β , PDK1, p70S6K	Fulvestrant \pm enzastaurin in ER-positive metastatic breast cancer after progression while receiving AI	II	NCT00451555 (ongoing)
Farnesyl transferase	Letrozole \pm tipifarnib in ER-positive metastatic breast cancer after progression on tamoxifen	II	Johnston et al ¹²³
IGF-1R/InsR	BMS-754807 \pm letrozole in ER-positive metastatic breast cancer after progression while receiving AI	II	NCT01225172 (ongoing)
IGF-1R	IMCA12 \pm same antiestrogen (AI, fulvestrant, or tamoxifen) after progression in patients with ER-positive metastatic breast cancer	II	NCT00728949 (ongoing)
	AI or fulvestrant \pm AMG479 in ER-positive advanced or metastatic breast cancer after progression while receiving endocrine Tx	II	Kaufman et al ¹²⁴

Abbreviations: AI, aromatase inhibitor; BOLERO, Breast cancer trials of OraL Everolimus; EGFR, epidermal growth factor receptor; ER, estrogen receptor; HER2, human epidermal growth factor receptor 2; IGF-1R, insulin-like growth factor-1 receptor; InsR, insulin receptor; mTOR, mammalian target of rapamycin; PI3K, phosphatidylinositol 3-kinase; PKC, protein kinase C; TAMRAD, Tamoxifen and RAD001.




Fig 2. Diagram of neoadjuvant clinical trial with phosphatidylinositol 3-kinase (PI3K) pathway inhibitor. Patients with estrogen receptor (ER) –positive breast cancer eligible for neoadjuvant therapy randomly assigned to standard treatment (eg, aromatase inhibitor [AI]) with or without PI3K pathway inhibitor. Core biopsies obtained before and after 2 weeks of therapy to document effects on tumor cell proliferation, apoptosis, and pathway inactivation (ie, downregulation of P-AKT, P-S6, P-proline-rich AKT substrate 40 by immunohistochemistry). Incorporation of noninvasive [^{18}F]fluorodeoxyglucose (FDG) –positron emission tomography (PET) at 2 weeks could identify early metabolic changes; however, PI3K pathway inhibitors may induce hyperglycemia,¹⁰⁶⁻¹⁰⁸ which could limit utility of FDG-PET. Primary end point of clinical response could be evaluated after approximately 4 to 6 months of therapy by measuring tumor with calipers, ultrasound (U/S), and/or mammography. Absence of tumor in surgical specimen would be scored as pathologic complete response (CR). Rate of breast-conserving surgery could be compared between both arms. Molecular mining of baseline, 2-week, and surgical biopsies could identify biomarkers that could be used for selection of patients enrolled onto subsequent studies of combination. HER2, human epidermal growth factor receptor 2; TUNEL, terminal deoxynucleotidyl nick-end labeling.

breast cancer.¹⁵ In the TAMRAD (Tamoxifen and RAD001) trial, 111 patients bearing AI-resistant metastatic breast cancer (resistance defined as primary [relapse during adjuvant AI or < 6 months after starting adjuvant AI in metastatic setting] or secondary [relapse \geq 6 months on adjuvant AI or prior response to AI and subsequent metastatic progression]) were randomly assigned to tamoxifen with or without everolimus. Patients in the combination arm showed an improved clinical benefit rate (61% *v* 42%), time to progression (8.6 *v* 4.5 months), and overall survival compared with patients receiving tamoxifen alone. Patients with secondary but not primary AI resistance who received both drugs showed an increased time to progression compared with patients receiving tamoxifen alone.⁷⁹ In contrast, results from other studies have not favored the combination arm. For example, addition of the IGF-1R antibody AMG479 to fulvestrant or exemestane did not alter progression-free survival in patients with ER-positive breast cancer compared with endocrine therapy alone.¹²⁴ The addition of the farnesyltransferase inhibitor tipifarnib to letrozole did not improve response in patients with advanced ER-positive disease compared with letrozole alone.¹²³

The neoadjuvant trial design depicted in Figure 2 illustrates an approach that can be used to determine whether to pursue combinations of PI3K inhibitors and antiestrogens in patients with ER-positive breast cancer. Such trials would have to be performed after safety of the combinations has been documented in phase I studies. Patients would be randomly assigned to standard endocrine therapy with or without a PI3K pathway inhibitor. A research biopsy could be obtained after 2 weeks to document effects on tumor cell proliferation, apoptosis, and ER/PI3K signaling and for wider exploratory mutational analysis. Incorporation of noninvasive imaging with [^{18}F]fluorodeoxyglucose–positron emission tomography at this time point could identify metabolic changes indicative of a pharmacodynamic effect. Study end points would be clinical and/or pathologic complete response (CR) after 4 to 6 months of therapy. Historically, pathologic CR has had limitations as an end point of neoadjuvant trials in ER-positive breast cancer. However, we cannot rule out that as we optimize targeted approaches such as antiestrogens in combination with PI3K inhibitors in ER-positive/PI3K-mutant

tumors, a pathologic CR rate could be used as a primary end point for a trial of this design. This approach potentially addresses the following questions: First, is there an early (at 2 weeks) difference in cellular and molecular response between treatment arms? Second, is clinical and/or pathologic response superior in the arm containing the PI3K pathway inhibitor? A difference in favor of the combination would support further development of PI3K inhibitors and endocrine therapy in patients with advanced disease. Third, is there a tissue and/or imaging pharmacodynamic biomarker in the baseline, 2-week, and/or surgical specimen that correlates with response/resistance to the combination? If so, such a biomarker could be used to select patients with advanced disease who would likely benefit from the combination in phase II trials.

DISCUSSION

Alterations in the PI3K pathway are the most common somatic mutations in ER-positive breast cancer. Experimental and clinical evidence implies that such mutations are associated with antiestrogen resistance. Many PI3K pathway inhibitors are in clinical development. Early clinical data suggest that this strategy is feasible and that as single agents, these drugs are well tolerated. Although there is crosstalk between the ER and PI3K/AKT pathways, these networks also signal independently. Because most ER-positive breast cancers that acquire resistance to antiestrogens retain ER and responsiveness to estrogens, we speculate PI3K inhibitors should be used in combination with antiestrogens in patients who progress while receiving the latter. To determine if combined inhibition of PI3K and ER is more active than antiestrogen therapy alone against ER-positive tumors with mutations in the PI3K pathway, randomized clinical trials are required.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject

matter under consideration in this article. Certain relationships marked with a “U” are those for which no compensation was received; those relationships marked with a “C” were compensated. For a detailed description of the disclosure categories, or for more information about ASCO’s conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: None **Consultant or Advisory**

Role: Carlos L. Arteaga, Roche (C), Monogram Biosciences (C), AstraZeneca (C), GlaxoSmithKline (C) **Stock Ownership:** None

Honoraria: None **Research Funding:** Carlos L. Arteaga, U3-Amgen, AstraZeneca, Exelixis **Expert Testimony:** None **Other Remuneration:** None

AUTHOR CONTRIBUTIONS

Manuscript writing: All authors

Final approval of manuscript: All authors

REFERENCES

1. Early Breast Cancer Trialists’ Collaborative Group: Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: An overview of the randomised trials. *Lancet* 365:1687-1717, 2005
2. Forbes JF, Cuzick J, Buzdar A, et al: Effect of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: 100-month analysis of the ATAC trial. *Lancet Oncol* 9:45-53, 2008
3. Ellis MJ, Tao Y, Young O, et al: Estrogen-independent proliferation is present in estrogen-receptor HER2-positive primary breast cancer after neoadjuvant letrozole. *J Clin Oncol* 24:3019-3025, 2006
4. Arpino G, Green SJ, Allred DC, et al: HER-2 amplification, HER-1 expression, and tamoxifen response in estrogen receptor-positive metastatic breast cancer: A southwest oncology group study. *Clin Cancer Res* 10:5670-5676, 2004
5. De Laurentiis M, Arpino G, Massarelli E, et al: A meta-analysis on the interaction between HER-2 expression and response to endocrine treatment in advanced breast cancer. *Clin Cancer Res* 11:4741-4748, 2005
6. Gottardis MM, Jordan VC: Development of tamoxifen-stimulated growth of MCF-7 tumors in athymic mice after long-term antiestrogen administration. *Cancer Res* 48:5183-5187, 1988
7. Shou J, Massarweh S, Osborne CK, et al: Mechanisms of tamoxifen resistance: Increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer. *J Natl Cancer Inst* 96:926-935, 2004
8. Bartlett JMS, Brookes CL, van de Velde CJH, et al: Final results of a prospectively planned biomarker analysis: HER1-3 as predictive markers of benefit from early treatment with aromatase inhibitors versus tamoxifen in the TEAM pathology sub-study. *Cancer Res* 70, 2010 (abstr S2-4)
9. Robertson JF, Llombart-Cussac A, Rolski J, et al: Activity of fulvestrant 500 mg versus anastrozole 1 mg as first-line treatment for advanced breast cancer: Results from the FIRST study. *J Clin Oncol* 27:4530-4535, 2009
10. Robertson JFR, Dixon JM, Sibbering DM, et al: Tumor biomarker changes following pre-surgical treatment with 500 mg fulvestrant plus anastrozole versus 500 mg fulvestrant alone and 1 mg anastrozole alone. *Cancer Res* 69, 2009 (abstr 24)
11. Robertson JFR, Lindemann JPO, Llombart-Cussac A, et al: A comparison of fulvestrant 500 mg with anastrozole as first-line treatment for advanced breast cancer: Follow-up analysis from the “FIRST” study. *Cancer Res* 70, 2010 (abstr S1-3)
12. Pérez-Tenorio G, Alkhori L, Olsson B, et al: PIK3CA mutations and PTEN loss correlate with similar prognostic factors and are not mutually exclusive in breast cancer. *Clin Cancer Res* 13:3577-3584, 2007
13. Saal LH, Johansson P, Holm K, et al: Poor prognosis in carcinoma is associated with a gene expression signature of aberrant PTEN tumor suppressor pathway activity. *Proc Natl Acad Sci U S A* 104:7564-7569, 2007
14. Shoman N, Klassen S, McFadden A, et al: Reduced PTEN expression predicts relapse in patients with breast carcinoma treated by tamoxifen. *Mod Pathol* 18:250-259, 2005
15. Baselga J, Semiglazov V, van Dam P, et al: Phase II randomized study of neoadjuvant everolimus plus letrozole compared with placebo plus letrozole in patients with estrogen receptor-positive breast cancer. *J Clin Oncol* 27:2630-2637, 2009
16. Stemke-Hale K, Gonzalez-Angulo AM, Lluch A, et al: An integrative genomic and proteomic analysis of PIK3CA, PTEN, and AKT mutations in breast cancer. *Cancer Res* 68:6084-6091, 2008
17. Ellis MJ, Lin L, Crowder R, et al: Phosphatidylinositol-3-kinase alpha catalytic subunit mutation and response to neoadjuvant endocrine therapy for estrogen receptor positive breast cancer. *Breast Cancer Res Treat* 119:379-390, 2010
18. Campbell IG, Russell SE, Choong DY, et al: Mutation of the PIK3CA gene in ovarian and breast cancer. *Cancer Res* 64:7678-7681, 2004
19. Crowder RJ, Phommaly C, Tao Y, et al: PIK3CA and PIK3CB inhibition produce synthetic lethality when combined with estrogen deprivation in estrogen receptor-positive breast cancer. *Cancer Res* 69:3955-3962, 2009
20. Law JH, Habibi G, Hu K, et al: Phosphorylated insulin-like growth factor-I/insulin receptor is present in all breast cancer subtypes and is related to poor survival. *Cancer Res* 68:10238-10246, 2008
21. Turner N, Pearson A, Sharpe R, et al: FGFR1 amplification drives endocrine therapy resistance and is a therapeutic target in breast cancer. *Cancer Res* 70:2085-2094, 2010
22. Monni O, Barlund M, Mousses S, et al: Comprehensive copy number and gene expression profiling of the 17q23 amplicon in human breast cancer. *Proc Natl Acad Sci U S A* 98:5711-5716, 2001
23. Gewinner C, Wang ZC, Richardson A, et al: Evidence that inositol polyphosphate 4-phosphatase type II is a tumor suppressor that inhibits PI3K signaling. *Cancer Cell* 16:115-125, 2009
24. Fedele CG, Ooms LM, Ho M, et al: Inositol polyphosphate 4-phosphatase II regulates PI3K/AKT signaling and is lost in human basal-like breast cancers. *Proc Natl Acad Sci U S A* 107:22231-22236, 2010
25. Jaiswal BS, Janakiraman V, Kijavini NM, et al: Somatic mutations in p85alpha promote tumorigenesis through class IA PI3K activation. *Cancer Cell* 16:463-474, 2009
26. Loi S, Haibe-Kains B, Majaj S, et al: PIK3CA mutations associated with gene signature of low mTORC1 signaling and better outcomes in estrogen receptor-positive breast cancer. *Proc Natl Acad Sci U S A* 107:10208-10213, 2010
27. Carpten JD, Faber AL, Horn C, et al: A transforming mutation in the pleckstrin homology domain of AKT1 in cancer. *Nature* 448:439-444, 2007
28. Bellacosa A, de Feo D, Godwin AK, et al: Molecular alterations of the AKT2 oncogene in ovarian and breast carcinomas. *Int J Cancer* 64:280-285, 1995
29. Al-Kuraya K, Schraml P, Torhorst J, et al: Prognostic relevance of gene amplifications and coamplifications in breast cancer. *Cancer Res* 64:8534-8540, 2004
30. Maurer M, Su T, Saal LH, et al: 3-phosphoinositide-dependent kinase 1 potentiates upstream lesions on the phosphatidylinositol 3-kinase pathway in breast carcinoma. *Cancer Res* 69:6299-6306, 2009
31. Rochlitz CF, Scott GK, Dodson JM, et al: Incidence of activating ras oncogene mutations associated with primary and metastatic human breast cancer. *Cancer Res* 49:357-360, 1989
32. Di Nicolantonio F, Arena S, Tabernero J, et al: Deregulation of the PI3K and KRAS signaling pathways in human cancer cells determines their response to everolimus. *J Clin Invest* 120:2858-2866, 2010
33. Whitman M, Downes CP, Keeler M, et al: Type I phosphatidylinositol kinase makes a novel inositol phospholipid, phosphatidylinositol-3-phosphate. *Nature* 332:644-646, 1988
34. Anderson KE, Coadwell J, Stephens LR, et al: Translocation of PDK-1 to the plasma membrane is important in allowing PDK-1 to activate protein kinase B. *Curr Biol* 8:684-691, 1998
35. Franke TF, Yang SI, Chan TO, et al: The protein kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase. *Cell* 81:727-736, 1995
36. Maehama T, Dixon JE: The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem* 273:13375-13378, 1998
37. Gingras AC, Kennedy SG, O’Leary MA, et al: 4E-BP1, a repressor of mRNA translation, is phosphorylated and inactivated by the Akt(PKB) signaling pathway. *Genes Dev* 12:502-513, 1998
38. Sarbassov DD, Ali SM, Kim DH, et al: Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr Biol* 14:1296-1302, 2004
39. Campbell RA, Bhat-Nakshatri P, Patel NM, et al: Phosphatidylinositol 3-kinase/AKT-mediated activation of estrogen receptor alpha: A new model for anti-estrogen resistance. *J Biol Chem* 276:9817-9824, 2001
40. Yamnik RL, Digilova A, Davis DC, et al: S6 kinase 1 regulates estrogen receptor alpha in control

of breast cancer cell proliferation. *J Biol Chem* 284:6361-6369, 2009

41. Coso OA, Chiariello M, Yu JC, et al: The small GTP-binding proteins Rac1 and Cdc42 regulate the activity of the JNK/SAPK signaling pathway. *Cell* 81:1137-1146, 1995
42. Minden A, Lin A, Claret FX, et al: Selective activation of the JNK signaling cascade and c-Jun transcriptional activity by the small GTPases Rac and Cdc42Hs. *Cell* 81:1147-1157, 1995
43. Logan SK, Falasca M, Hu P, et al: Phosphatidylinositol 3-kinase mediates epidermal growth factor-induced activation of the c-Jun N-terminal kinase signaling pathway. *Mol Cell Biol* 17:5784-5790, 1997
44. Hibi M, Lin A, Smeal T, et al: Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain. *Genes Dev* 7:2135-2148, 1993
45. Derijard B, Hibi M, Wu IH, et al: JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 76:1025-1037, 1994
46. Rauscher FJ 3rd, Cohen DR, Curran T, et al: Fos-associated protein p39 is the product of the jun proto-oncogene. *Science* 240:1010-1016, 1988
47. DeNardo DG, Cuba VL, Kim H, et al: Estrogen receptor DNA binding is not required for estrogen-induced breast cell growth. *Mol Cell Endocrinol* 277:13-25, 2007
48. Petz LN, Ziegler YS, Loven MA, et al: Estrogen receptor alpha and activating protein-1 mediate estrogen responsiveness of the progesterone receptor gene in MCF-7 breast cancer cells. *Endocrinology* 143:4583-4591, 2002
49. Massarweh S, Schiff R: Unraveling the mechanisms of endocrine resistance in breast cancer: New therapeutic opportunities. *Clin Cancer Res* 13:1950-1954, 2007
50. Generali D, Fox SB, Brizzi MP, et al: Down-regulation of phosphatidylinositol 3'-kinase/AKT/molecular target of rapamycin metabolic pathway by primary letrozole-based therapy in human breast cancer. *Clin Cancer Res* 14:2673-2680, 2008
51. Miller TW, Perez-Torres M, Narasanna A, et al: Loss of Phosphatase and Tensin homologue deleted on chromosome 10 engages ErbB3 and insulin-like growth factor-I receptor signaling to promote antiestrogen resistance in breast cancer. *Cancer Res* 69:4192-4201, 2009
52. Massarweh S, Osborne CK, Creighton CJ, et al: Tamoxifen resistance in breast tumors is driven by growth factor receptor signaling with repression of classic estrogen receptor genomic function. *Cancer Res* 68:826-833, 2008
53. Sabnis G, Schayowitz A, Golubeva O, et al: Trastuzumab reverses letrozole resistance and amplifies the sensitivity of breast cancer cells to estrogen. *Cancer Res* 69:1416-1428, 2009
54. Wang X, Masri S, Phung S, et al: The role of amphiregulin in exemestane-resistant breast cancer cells: Evidence of an autocrine loop. *Cancer Res* 68:2259-2265, 2008
55. Miller TW, Hennessy BT, Gonzalez-Angulo AM, et al: Hyperactivation of phosphatidylinositol-3 kinase promotes escape from hormone dependence in estrogen receptor-positive human breast cancer. *J Clin Invest* 120:2406-2413, 2010
56. Fan M, Yan PS, Hartman-Frey C, et al: Diverse gene expression and DNA methylation profiles correlate with differential adaptation of breast cancer cells to the antiestrogens tamoxifen and fulvestrant. *Cancer Res* 66:11954-11966, 2006
57. Frogne T, Benjaminsen RV, Sonne-Hansen K, et al: Activation of ErbB3, EGFR and Erk is essential for growth of human breast cancer cell lines with acquired resistance to fulvestrant. *Breast Cancer Res Treat* 114:263-275, 2009
58. Massarweh S, Osborne CK, Jiang S, et al: Mechanisms of tumor regression and resistance to estrogen deprivation and fulvestrant in a model of estrogen receptor-positive, HER-2/neu-positive breast cancer. *Cancer Res* 66:8266-8273, 2006
59. Simoncini T, Hafezi-Moghadam A, Brazil DP, et al: Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* 407:538-541, 2000
60. Song RX, McPherson RA, Adam L, et al: Linkage of rapid estrogen action to MAPK activation by ERalpha-Shc association and Shc pathway activation. *Mol Endocrinol* 16:116-127, 2002
61. Song RX, Fan P, Yue W, et al: Role of receptor complexes in the extranuclear actions of estrogen receptor alpha in breast cancer. *Endocr Relat Cancer* 13:S3-S13, 2006 (suppl 1)
62. Wang Z, Zhang X, Shen P, et al: A variant of estrogen receptor-(alpha), hER-(alpha)36: transduction of estrogen- and antiestrogen-dependent membrane-initiated mitogenic signaling. *Proc Natl Acad Sci U S A* 103:9063-9068, 2006
63. Li L, Haynes MP, Bender JR: Plasma membrane localization and function of the estrogen receptor alpha variant (ER46) in human endothelial cells. *Proc Natl Acad Sci U S A* 100:4807-4812, 2003
64. Filardo EJ, Quinn JA, Bland KL, et al: Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30, and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol* 14:1649-1660, 2000
65. Pedram A, Razandi M, Levin ER: Nature of functional estrogen receptors at the plasma membrane. *Mol Endocrinol* 20:1996-2009, 2006
66. Voliva CF, Pecchi S, Burger M, et al: Biological characterization of NVP-BKM120, a novel inhibitor of phosphoinositide 3-kinase in phase I/II clinical trials. *Cancer Res* 70, 2010 (abstr 4498)
67. Sabnis G, Golubeva O, Jelovac D, et al: Inhibition of the phosphatidylinositol 3-kinase/Akt pathway improves response of long-term estrogen-deprived breast cancer xenografts to antiestrogens. *Clin Cancer Res* 13:2751-2757, 2007
68. Yue W, Wang J, Li Y, et al: Farnesylthiosalicylic acid blocks mammalian target of rapamycin signaling in breast cancer cells. *Int J Cancer* 117:746-754, 2005
69. Meyer DS, Brinkhaus H, Muller U, et al: Luminal expression of PIK3CA mutant H1047R in the mammary gland induces heterogeneous tumors. *Cancer Res* 71:4344-4351, 2011
70. Zhao L, Vogt PK: Helical domain and kinase domain mutations in p110alpha of phosphatidylinositol 3-kinase induce gain of function by different mechanisms. *Proc Natl Acad Sci U S A* 105:2652-2657, 2008
71. Isakoff SJ, Engelman JA, Irie HY, et al: Breast cancer-associated PIK3CA mutations are oncogenic in mammary epithelial cells. *Cancer Res* 65:10992-11000, 2005
72. Samuels Y, Wang Z, Bardelli A, et al: High frequency of mutations of the PIK3CA gene in human cancers. *Science* 304:554, 2004
73. Janku F, Tsimberidou AM, Garrido-Laguna I, et al: PIK3CA Mutations in Patients with Advanced Cancers Treated with PI3K/AKT/mTOR Axis Inhibitors. *Mol Cancer Ther* 10:558-565, 2011
74. Miron A, Varadi M, Carrasco D, et al: PIK3CA mutations in situ and invasive breast carcinomas. *Cancer Res* 70:5674-5678, 2011
75. Dupont Jensen J, Laenkholm AV, Knoop A, et al: PIK3CA mutations may be discordant between primary and corresponding metastatic disease in breast cancer. *Clin Cancer Res* 17:667-677, 2011
76. Gonzalez-Angulo AM, Ferrer-Lozano J, Stemke-Hale K, et al: PI3K pathway mutations and PTEN levels in primary and metastatic breast cancer. *Mol Cancer Ther* 10:1093-1101, 2011
77. Creighton CJ, Fu X, Hennessy BT, et al: Proteomic and transcriptomic profiling reveals a link between the PI3K pathway and lower estrogen receptor (ER) levels and activity in ER+ breast cancer. *Breast Cancer Res* 12:R40, 2010
78. Sorlie T, Perou CM, Tibshirani R, et al: Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 98:10869-10874, 2001
79. Bachelot T, Bourcier C, Cropet C, et al: TAMRAD: A GINECO randomized phase II trial of everolimus in combination with tamoxifen versus tamoxifen alone in patients (pts) with hormone-receptor positive, HER2 negative metastatic breast cancer (MBC) with prior exposure to aromatase inhibitors (AI). *Cancer Res* 70, 2010 (abstr S1-6)
80. Miller TW, Forbes JT, Shah C, et al: Inhibition of mammalian target of rapamycin is required for optimal antitumor effect of HER2 inhibitors against HER2-overexpressing cancer cells. *Clin Cancer Res* 15:7266-7276, 2009
81. O'Reilly KE, Rojo F, She QB, et al: mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates Akt. *Cancer Res* 66:1500-1508, 2006
82. Carracedo A, Ma L, Teruya-Feldstein J, et al: Inhibition of mTORC1 leads to MAPK pathway activation through a PI3K-dependent feedback loop in human cancer. *J Clin Invest* 118:3065-3074, 2008
83. Chakrabarty A, Sanchez V, Kuba MG, et al: Breast cancer special feature: Feedback upregulation of HER3 (ErbB3) expression and activity attenuates antitumor effect of PI3K inhibitors. *Proc Natl Acad Sci U S A* [epub ahead of print on February 28, 2011]
84. Chandralapaty S, Sawai A, Scaltriti M, et al: AKT inhibition relieves feedback suppression of receptor tyrosine kinase expression and activity. *Cancer Cell* 19:58-71, 2011
85. Garrett JT, Olivares MG, Rinehart C, et al: Transcriptional and posttranslational up-regulation of HER3 (ErbB3) compensates for inhibition of the HER2 tyrosine kinase. *Proc Natl Acad Sci U S A* 108:5021-5026, 2011
86. Amin DN, Sergina N, Ahuja D, et al: Resiliency and vulnerability in the HER2-HER3 tumorigenic driver. *Sci Transl Med* 2:16ra7, 2011
87. Lipton A, Ali SM, Leitzel K, et al: Serum HER-2/neu and response to the aromatase inhibitor letrozole versus tamoxifen. *J Clin Oncol* 21:1967-1972, 2003
88. Arpino G, Wiechmann L, Osborne CK, et al: Crosstalk between the estrogen receptor and the HER tyrosine kinase receptor family: Molecular mechanism and clinical implications for endocrine therapy resistance. *Endocr Rev* 29:217-233, 2008
89. Benz CC, Scott GK, Sarup JC, et al: Estrogen-dependent, tamoxifen-resistant tumorigenic growth of MCF-7 cells transfected with HER2/neu. *Breast Cancer Res Treat* 24:85-95, 1992
90. Pietras RJ, Arboleda J, Reese DM, et al: HER-2 tyrosine kinase pathway targets estrogen receptor and promotes hormone-independent growth in

human breast cancer cells. *Oncogene* 10:2435-2446, 1995

91. Gutierrez MC, Detre S, Johnston S, et al: Molecular changes in tamoxifen-resistant breast cancer: Relationship between estrogen receptor, HER-2, and p38 mitogen-activated protein kinase. *J Clin Oncol* 23:2469-2476, 2005

92. Xia W, Bacus S, Hegde P, et al: A model of acquired autoresistance to a potent ErbB2 tyrosine kinase inhibitor and a therapeutic strategy to prevent its onset in breast cancer. *Proc Natl Acad Sci U S A* 103:7795-7800, 2006

93. Munzone E, Curigliano G, Rocca A, et al: Reverting estrogen-receptor-negative phenotype in HER-2-overexpressing advanced breast cancer patients exposed to trastuzumab plus chemotherapy. *Breast Cancer Res* 8:R4, 2006

94. Johnston S, Pippen J Jr, Pivot X, et al: Lapatinib combined with letrozole versus letrozole and placebo as first-line therapy for postmenopausal hormone receptor-positive metastatic breast cancer. *J Clin Oncol* 27:5538-5546, 2009

95. Kaufman B, Mackey JR, Clemens MR, et al: Trastuzumab plus anastrozole versus anastrozole alone for the treatment of postmenopausal women with human epidermal growth factor receptor 2-positive, hormone receptor-positive metastatic breast cancer: Results from the randomized phase III TAnDEM study. *J Clin Oncol* 27:5529-5537, 2009

96. Engelman JA, Luo J, Cantley LC: The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet* 7:606-619, 2006

97. Cantley LC: The phosphoinositide 3-kinase pathway. *Science* 296:1655-1657, 2002

98. Vanhaesebroeck B, Leivers SJ, Panayotou G, et al: Phosphoinositide 3-kinases: A conserved family of signal transducers. *Trends Biochem Sci* 22:267-272, 1997

99. Jia S, Roberts TM, Zhao JJ: Should individual PI3 kinase isoforms be targeted in cancer? *Curr Opin Cell Biol* 21:199-208, 2009

100. Garcia-Echeverria C, Sellers WR: Drug discovery approaches targeting the PI3K/Akt pathway in cancer. *Oncogene* 27:5511-5526, 2008

101. Maira SM, Voliva C, Garcia-Echeverria C: Class IA phosphatidylinositol 3-kinase: From their biologic implication in human cancers to drug discovery. *Expert Opin Ther Targets* 12:223-238, 2008

102. Maira SM, Stauffer F, Brueggen J, et al: Identification and characterization of NVP-BEZ235, a new orally available dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor with potent in vivo antitumor activity. *Mol Cancer Ther* 7:1851-1863, 2008

103. Garlich JR, De P, Dey N, et al: A vascular targeted pan phosphoinositide 3-kinase inhibitor pro-drug, SF1126, with antitumor and antiangiogenic activity *Cancer Res* 68:206-215, 2008

104. Folkes AJ, Ahmadi K, Alderton WK, et al: The identification of 2-(1H-indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-d]pyrimidine (GDC-0941) as a potent, selective, orally bio-available inhibitor of class I PI3 kinase for the treatment of cancer. *J Med Chem* 51:5522-5532, 2008

105. Furman RR, Byrd JC, Flinn IW, et al: Interim results from a phase I study of CAL-101, a selective oral inhibitor of phosphatidylinositol 3-kinase p110δ isoform, in patients with relapsed or refractory hematologic malignancies. *J Clin Oncol* 28:241s, 2010 (suppl; abstr 3032)

106. Baselga J, De Jonge MJ, Rodon J, et al: A first-in-human phase I study of BKM120, an oral pan-class I PI3K inhibitor, in patients (pts) with advanced solid tumors. *J Clin Oncol* 28:233s, 2010 (suppl; abstr 3003)

107. Burris H, Rodon J, Sharma S, et al: First-in-human phase I study of the oral PI3K inhibitor BEZ235 in patients (pts) with advanced solid tumors. *J Clin Oncol* 28:234s, 2010 (suppl; abstr 3005)

108. Edelman G, Bedell C, Shapiro G, et al: A phase I dose-escalation study of XL147 (SAR245408), a PI3K inhibitor administered orally to patients (pts) with advanced malignancies. *J Clin Oncol* 28:234s, 2010 (suppl; abstr 3004)

109. Rhodes N, Heerding DA, Duckett DR, et al: Characterization of an Akt kinase inhibitor with potent pharmacodynamic and antitumor activity. *Cancer Res* 68:2366-2374, 2008

110. Davies TG, Verdonk ML, Graham B, et al: A structural comparison of inhibitor binding to PKB, PKA and PKA-PKB chimera. *J Mol Biol* 367:882-894, 2007

111. Toral-Barza L, Zhang WG, Huang X, et al: Discovery of lactoquinomycin and related pyranonaphthoquinones as potent and allosteric inhibitors of AKT/PKB: Mechanistic involvement of AKT catalytic activation loop cysteines. *Mol Cancer Ther* 6:3028-3038, 2007

112. She QB, Chandralapaty S, Ye Q, et al: Breast tumor cells with PI3K mutation or HER2 amplification are selectively addicted to Akt signaling. *PLoS ONE* 3:e3065, 2008

113. Yap TA, Yan L, Patnaik A, et al: Final results of a translational phase I study assessing a QOD schedule of the potent AKT inhibitor MK-2206 incorporating predictive, pharmacodynamic (PD), and functional imaging biomarkers. *J Clin Oncol* 29:194s, 2011 (suppl; abstr 3001)

114. Biondo A, Yap TA, Yan L, et al: Phase I clinical trial of an allosteric AKT inhibitor, MK-2206, using a once weekly (QW) dose regimen in patients with advanced solid tumors. *J Clin Oncol* 29:203s, 2011 (suppl; abstr 3037)

115. Fasolo A, Sessa C: MTOR inhibitors in the treatment of cancer. *Expert Opin Investig Drugs* 17:1717-1734, 2008

116. Stuart D, Sellers WR: Linking somatic genetic alterations in cancer to therapeutics. *Curr Opin Cell Biol* 21:304-310, 2009

117. Dowsett M, Smith IE, Ebbs SR, et al: Prognostic value of Ki67 expression after short-term presurgical endocrine therapy for primary breast cancer. *J Natl Cancer Inst* 99:167-170, 2007

118. Ellis MJ, Tao Y, Luo J, et al: Outcome prediction for estrogen receptor-positive breast cancer based on postneoadjuvant endocrine therapy tumor characteristics. *J Natl Cancer Inst* 100:1380-1388, 2008

119. Ellis PA, Smith IE, Detre S, et al: Reduced apoptosis and proliferation and increased Bcl-2 in residual breast cancer following preoperative chemotherapy. *Breast Cancer Res Treat* 48:107-116, 1998

120. Assersohn L, Salter J, Powles TJ, et al: Studies of the potential utility of Ki67 as a predictive molecular marker of clinical response in primary breast cancer. *Breast Cancer Res Treat* 82:113-123, 2003

121. Smith IE, Walsh G, Skene A, et al: A phase II placebo-controlled trial of neoadjuvant anastrozole alone or with gefitinib in early breast cancer. *J Clin Oncol* 25:3816-3822, 2007

122. Cristofanilli M, Valero V, Mangalik A, et al: Phase II, randomized trial to compare anastrozole combined with gefitinib or placebo in postmenopausal women with hormone receptor-positive metastatic breast cancer. *Clin Cancer Res* 16:1904-1914, 2010

123. Johnston SR, Semiglazov VF, Manikhas GM, et al: A phase II, randomized, blinded study of the farnesyltransferase inhibitor tipifarnib combined with letrozole in the treatment of advanced breast cancer after antiestrogen therapy. *Breast Cancer Res Treat* 110:327-335, 2008

124. Kaufman PA, Ferrero JM, Bourgeois H, et al: A randomized, double-blind, placebo-controlled, phase 2 study of AMG 479 with exemestane (E) or fulvestrant (F) in postmenopausal women with hormone-receptor positive (HR+) metastatic (M) or locally advanced (LA) breast cancer (BC). *Cancer Res* 70, 2010 (abstr S1-4)