

PI3K-PKB/Akt Pathway

Brian A. Hemmings and David F. Restuccia

Friedrich Miescher Institute for Biomedical Research, Basel 4058, Switzerland

Correspondence: david.restuccia@fmi.ch

Identification of the phosphoinositide-3-kinase–protein kinase B/Akt (PI3K-PKB/Akt) pathway and activating receptor tyrosine kinases (RTKs) began in earnest in the early 1980s through vigorous attempts to characterize insulin receptor signaling (for review, see Alessi 2001; Brazil and Hemmings 2001). These humble beginnings led to the identification of the components and mechanism of insulin receptor signaling via insulin receptor substrate (IRS) proteins to PI3K and consequent PKB/Akt-mediated activation by 3-phosphoinositide-dependent protein kinase 1 (PDK1). With the discovery of the potent contribution of PI3K and PKB/Akt activation to tumorigenesis, intense research into the regulation of this pathway led to the discovery of the negative regulators, the protein phosphatase 2 (PP2A), phosphatase and tensin homolog (PTEN), and the PH-domain leucine-rich-repeat-containing protein phosphatases (PHLPP1/2). More recently, the elusive PKB/

Akt hydrophobic motif kinases—i.e., the mammalian target of rapamycin (mTOR), when associated with the mTOR complex 2 (mTORC2), and the DNA-dependent protein kinase (DNA-PK)—were identified, as was the ability of Ras to affect the PI3K-PKB/Akt pathway via PI3K, completing our current model of the PI3K-PKB/Akt pathway.

The PI3K-PKB/Akt pathway is highly conserved, and its activation is tightly controlled via a multistep process (as shown in Fig. 1). Activated receptors directly stimulate class 1A PI3Ks bound via their regulatory subunit or adapter molecules such as the insulin receptor substrate (IRS) proteins. This triggers activation of PI3K and conversion by its catalytic domain of phosphatidylinositol (3,4)-bisphosphate (PIP₂) lipids to phosphatidylinositol (3,4,5)-trisphosphate (PIP₃). PKB/Akt binds to PIP₃ at the plasma membrane, allowing PDK1 to access and phosphorylate




Figure 1. PKB/Akt activation downstream of RTKs via the P13K pathway.

Editors: Lewis Cantley, Tony Hunter, Richard Sever, and Jeremy Thorner
Additional Perspectives on Signal Transduction available at www.cshperspectives.org

Copyright © 2012 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a011189
Cite this article as *Cold Spring Harb Perspect Biol* 2012;4:a011189




Figure 2. Signalling events activating PKB/Akt and cellular functions regulated by PKB/Akt.

T308 in the “activation loop,” leading to partial PKB/Akt activation (Alessi et al. 1997). This PKB/Akt modification is sufficient to activate mTORC1 by directly phosphorylating and inactivating proline-rich Akt substrate of 40 kDa (PRAS40) and tuberous sclerosis protein 2 (TSC2) (Vander Haar et al. 2007). mTORC1 substrates include the eukaryotic translation initiation factor 4E binding protein 1 (4EBP1),

and ribosomal protein S6 kinase, 70 kDa, polypeptide 1 (S6K1), which, in turn, phosphorylates the ribosomal protein S6 (S6/RPS6), promoting protein synthesis and cellular proliferation (Figure 1).

Phosphorylation of Akt at S473 in the carboxy-terminal hydrophobic motif, either by mTOR (Sarbasov et al. 2005) or by DNA-PK (Feng et al. 2004), stimulates full Akt activity.

Full activation of Akt leads to additional substrate-specific phosphorylation events in both the cytoplasm and nucleus, including inhibitory phosphorylation of the pro-apoptotic FOXO proteins (Guertin et al. 2006). Fully active PKB/Akt mediates numerous cellular functions including angiogenesis, metabolism, growth, proliferation, survival, protein synthesis, transcription, and apoptosis (as shown in Fig. 2). Dephosphorylation of T308 by PP2A (Andjelković et al. 1996), and S473 by PHLPP1/2 (Brognaard et al. 2007), and the conversion of PIP₃ to PIP₂ by PTEN (Stambolic et al. 1998) antagonize Akt signaling. Figure 2.

Figures adapted, with kind permission, from Cell Signaling Technology (<http://www.cellsignal.com>.)

REFERENCES

- Alessi DR. 2001. Discovery of PDK1, one of the missing links in insulin signal transduction. Colworth Medal Lecture. *Biochem Soc Trans* **29**: 1–14.
- Alessi DR, James SR, Downes CP, Holmes AB, Gaffney PR, Reese CB, Cohen P. 1997. Characterization of a 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase B α . *Curr Biol* **7**: 261–269.
- Altomare DA, Testa JR. 2005. Perturbations of the AKT signaling pathway in human cancer. *Oncogene* **24**: 7455–7464.
- Andjelković M, Jakubowicz T, Cron P, Ming XF, Han JW, Hemmings BA. 1996. Activation and phosphorylation of a pleckstrin homology domain containing protein kinase (RAC-PK/PKB) promoted by serum and protein phosphatase inhibitors. *Proc Natl Acad Sci* **93**: 5699–5704.
- Bozulic L, Hemmings BA. 2009. PIKKing on PKB: Regulation of PKB activity by phosphorylation. *Curr Opin Cell Biol* **21**: 256–261.
- Brazil DP, Hemmings BA. 2001. Ten years of protein kinase B signalling: A hard Akt to follow. *Trends Biochem Sci* **26**: 657–664.
- Brognaard J, Sierceki E, Gao T, Newton AC. 2007. PHLPP and a second isoform, PHLPP2, differentially attenuate the amplitude of Akt signaling by regulating distinct Akt isoforms. *Mol Cell* **25**: 917–931.
- Feng J, Park J, Cron P, Hess D, Hemmings BA. 2004. Identification of a PKB/Akt hydrophobic motif Ser-473 kinase as DNA-dependent protein kinase. *J Biol Chem* **279**: 41189–41196.
- Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, Brown M, Fitzgerald KJ, Sabatini DM. 2006. Ablation in mice of the mTORC components raptor, rictor, or mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKC α , but not S6K1. *Dev Cell* **11**: 859–871.
- Manning BD, Cantley LC. 2007. AKT/PKB signaling: Navigating downstream. *Cell* **129**: 1261–1274.
- Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. 2005. Phosphorylation and regulation of Akt/PKB by the rictor–mTOR complex. *Science* **307**: 1098–1101.
- Stambolic V, Suzuki A, de la Pompa JL, Brothers GM, Mirtsos C, Sasaki T, Ruland J, Penninger JM, Siderovski DP, Mak TW. 1998. Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN. *Cell* **95**: 29–39.
- Vander Haar E, Lee SI, Bandhakavi S, Griffin TJ, Kim DH. 2007. Insulin signalling to mTOR mediated by the Akt/PKB substrate PRAS40. *Nat Cell Biol* **9**: 316–323.



Cold Spring Harbor Perspectives in Biology

PI3K-PKB/Akt Pathway

Brian A. Hemmings and David F. Restuccia

Cold Spring Harb Perspect Biol 2012; doi: 10.1101/cshperspect.a011189

Subject Collection [Signal Transduction](#)

Cell Signaling and Stress Responses

Gökhan S. Hotamisligil and Roger J. Davis

Protein Regulation in Signal Transduction

Michael J. Lee and Michael B. Yaffe

Synaptic Signaling in Learning and Memory

Mary B. Kennedy

Vertebrate Reproduction

Sally Kornbluth and Rafael Fissore

Signaling in Lymphocyte Activation

Doreen Cantrell

Signaling in Muscle Contraction

Ivana Y. Kuo and Barbara E. Ehrlich

Toll-Like Receptor Signaling

Kian-Huat Lim and Louis M. Staudt

Signaling Pathways that Regulate Cell Division

Nicholas Rhind and Paul Russell

Second Messengers

Alexandra C. Newton, Martin D. Bootman and John D. Scott

Signals and Receptors

Carl-Henrik Heldin, Benson Lu, Ron Evans, et al.

Cell Death Signaling

Douglas R. Green and Fabien Llambi

Signaling Networks that Regulate Cell Migration

Peter Devreotes and Alan Rick Horwitz

Signaling Networks: Information Flow, Computation, and Decision Making

Evren U. Azeloglu and Ravi Iyengar

Signal Transduction: From the Atomic Age to the Post-Genomic Era

Jeremy Thorner, Tony Hunter, Lewis C. Cantley, et al.

Signaling by the TGF β Superfamily

Jeffrey L. Wrana

Subversion of Cell Signaling by Pathogens

Neal M. Alto and Kim Orth

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>



**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

