

1 Evolution along allometric lines of least resistance: Morphological
2 differentiation in *Pristurus* geckos

3
4 **Héctor Tejero-Cicuéndez^{1,2,*}, Iris Menéndez³, Adrián Talavera², Gabriel Riaño²,**
5 **Bernat Burriel-Carranza², Marc Simó-Riudalbas², Salvador Carranza², and Dean C.**
6 **Adams⁴**

7 ~~11~~¹⁵ April, 2023

8 ¹Evolution and Conservation Biology research Group, Department of Biodiversity, Ecology and
9 Evolution. Faculty of Biology. Universidad Complutense de Madrid, 28040, Madrid, Spain

10 ²Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la
11 Barceloneta 37-49, Barcelona 08003, Spain

12 ³Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

13 ⁴Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa,
14 50010 USA

15 *Correspondence: Héctor Tejero-Cicuéndez cicuendez93@gmail.com

Abstract

Species living in distinct habitats often experience unique ecological selective pressures, which can drive phenotypic divergence. However, how ecophenotypic patterns are affected by allometric trends and trait integration levels is less well understood. Here we evaluate the role of allometry in shaping body size and ~~shape~~body form diversity in *Pristurus* geckos utilizing differing habitats. We found that patterns of ~~body shape~~-allometry and integration in body form were distinct in species with different habitat preferences, with ground-dwelling *Pristurus* displaying the most divergent allometric trend and ~~the strongest~~high levels of integration. There was also strong concordance between static allometry across individuals and evolutionary allometry among species, revealing that ~~body shape differences~~differences in body form among individuals were predictive of evolutionary changes across the phylogeny at macroevolutionary scales. This suggested that phenotypic evolution occurred along allometric lines of least resistance, with allometric trajectories imposing a strong influence on the magnitude and direction of size and shape changes across the phylogeny. When viewed in phylomorphospace, the largest rock-dwelling species were most similar ~~in body shape~~ to the smallest ground-dwelling species, and vice versa. Thus, in *Pristurus*, phenotypic evolution along the differing habitat-based allometric trajectories resulted in similar body ~~shapes~~forms at differing body sizes in distinct ecological habitats.

1. Introduction

Understanding how phenotypic diversity evolves, and elucidating the forces that generate and maintain this diversity, are major goals in evolutionary biology. Because adaptive evolution is the product of natural selection, changes in ecological selection pressures are expected to affect the evolutionary trajectory of phenotypic traits that facilitate an organism’s survival and reproduction in their habitat. ~~Evolutionary theory~~ The theory of natural selection predicts that differing habitats will exert unique ecological selection pressures on organisms, resulting in associations between ecological and phenotypic traits. Indeed, species inhabiting differing habitats often display functional, behavioral, or phenotypic differences, that have presumably been the result of adaptive diversification in their respective ecological contexts (Collar et al. 2010; Kaliontzopoulou et al. 2015; Price et al. 2015; Martinez et al. 2021; Kolmann et al. 2022).

One possible evolutionary outcome of ecological specialization is that organisms inhabiting similar environments display common phenotypic characteristics. When such patterns occur repeatedly (Losos 1992; Schluter and McPhail 1992), this convergent evolution is treated as strong evidence of adaptation. Indeed the ecomorphological paradigm (Arnold 1983) is predicated, in part, on such cases, which emphasize the strong association between the phenotypic traits that organisms display (morphological, behavioral, or physiological) and the ecological characteristics of their habitat that mediate organismal performance. In vertebrates, ecomorphological trends have been well studied in numerous taxonomic groups, and include the emblematic ‘ecomorphs’ of Caribbean *Anolis* lizards that exploit different microhabitats (Losos 1992, 2009; Mahler et al. 2013), differential beak morphology in species of Darwin’s finches (Schluter and Grant 1984; Grant and Grant 2006; Reaney et al. 2020), the recurring phenotypes of African lake cichlids across ecological regimes (Albertson and Kocher 2001; Urban et al. 2022), and the distinct body forms of freshwater fishes in benthic and limnetic habitats (Jastrebski and Robinson 2004; Berner et al. 2008; Stuart et al. 2017), among others.

However, while the patterns of morphological differences in distinct ecological contexts have been

well documented, less-well understood is how this differentiation has been influenced by trait covariation associated with body size differences (i.e., allometry). Evaluating allometric trends across ~~hierarchical~~ levels (e.g., comparing allometry at the individual level, ~~or static allometry, and among species, or evolutionary allometry~~ and at the evolutionary level among species) may aid in our understanding of how adaptive morphological change occurs at macroevolutionary scales (Klingenberg and Zimmermann 1992). DEAN: NEW ALLOMETRY STUFF HERE. Levels of allometry: ‘schools’ of allometric thought (see Intro of Klingenberg 2022), how allometry leads to patterns of phenotypic evolution, etc. Look at Klingenberg review on levels of allometry, and more recent one on allometry. Cite and discuss work by authors mentioned by reviewer: Klingenberg, Goswami, Adams... Cooney and Bright?? <https://www.nature.com/articles/nature21074> :), etc.

separated this part: probably needs a transition sentence It has long been recognized that the interrelationships among traits can exert a strong influence on how phenotypic evolution proceeds, as trait correlations influence the degree to which phenotypic variation is exposed to selection (Wagner and Altenberg 1996). Thus, the integration among traits can constrain phenotypic change in certain directions, or enhance variation along other phenotypic axes (Schluter 1996; Wagner and Altenberg 1996; Wagner and Zhang 2011; Klingenberg and Marugán-Lobón 2013; Goswami et al. 2014, 2016; Felice et al. 2018; Navalón et al. 2020). Further, because nearly all linear traits covary strongly with overall body size (Jolicoeur 1963; Bookstein 2022), allometric trends could be considered the quintessential expression of phenotypic integration. Thus, identifying whether allometric patterns differ across habitats, and how such patterns of trait covariation affect ecomorphological trends among species utilizing those habitats, remains an important question worthy of investigation.

The Afro-Arabian geckos in the genus *Pristurus* afford the opportunity to elucidate the interdigitating effects of allometry and habitat specialization on clade-level patterns of phenotypic diversity. ~~Prior~~ From an evolutionary perspective, *Pristurus* geckos are an ideal system to investigate the role of different factors in evolutionary history. They are found both insular and continental settings, which are known to impose differential ecological selection pressures resulting

in distinct evolutionary trajectories of species and clades (Losos and Ricklefs 2009). They are also distributed in the contact zone between Africa and Eurasia, a region of high biogeographic interest which has been the epicenter of major faunal interchanges and complex geologic and environmental processes (Kappelman et al. 2003; Tejero-Cicuéndez et al. 2022). The study of evolutionary dynamics in *Pristurus* and other Afro-Arabian taxa is also important biological, as our understanding of the dynamics of biological systems in remote arid regions is generally neglected and understudied (Durant et al. 2012). Furthermore, prior work on this system (Tejero-Cicuéndez et al. 2021a) has revealed that the colonization of ground habitats has been a trigger of morphological change, specifically reflected in an increase in body size and shape disparity. Interestingly, some ground-dwelling species are among the largest of the genus and also show increased relative head sizes and limb proportions, while some other species with this ecological specialization have evolved to be among the smallest of the group. Additionally, among the species exploiting rocky habitats (the most common ecological feature in *Pristurus*), there are also species with both considerably large and small body sizes (Tejero-Cicuéndez et al. 2021a). What remains unexplored, however, is how the evolution of body ~~shape-form~~ is related to differences in body size and whether habitat specialization has an impact in this shape-size relationship.

In this study, we employed a combination of multivariate morphometric and phylogenetic comparative analyses to interrogate macroevolutionary patterns of evolutionary allometry in *Pristurus* geckos of Afro-Arabia. Using phenotypic, phylogenetic, and ecological data, we first characterized allometric trends in body form in the group, to discern the extent to which evolutionary allometric trends across the phylogeny aligned with habitat-based static allometry for species occupying distinct ecological regimes. We then examined changes in allometric trends across the phylogeny, and linked these patterns to overall phenotypic integration, diversification in morphospace, and habitat utilization among taxa. Our analyses reveal that patterns of evolutionary allometry across species align with allometric trends ~~within habitats, demonstrating that the~~ among individuals, and that differing habitat-based allometric trajectories have resulted in similar body forms at differing body sizes in distinct ecological regimes. Thus, patterns of phenotypic diversification in *Pristurus* are the outcome of an interplay between ecological specialization and size-form changes evolving along

habitat-specific allometric trajectories in ~~species with disparate body size may play a determinant~~
~~role in shaping the phenotypic evolution and hence in adaptive dynamics in this clade~~the group.

2. Materials and Methods

(a) Data

We used a combination of phenotypic, phylogenetic, and ecological data to characterize and evaluate intra- and interspecific allometric trends in *Pristurus*. The data utilized here were obtained from our prior work on this system (Tejero-Cicuéndez et al. 2021a, 2022), and are briefly described here. First we used a ~~time-dated~~, time-calibrated molecular phylogeny of squamates that included all members of the genus *Pristurus*, including several currently undescribed taxa. The tree was estimated in a Bayesian framework, using five mitochondrial markers, six nuclear markers, and 21 calibration points (Tejero-Cicuéndez et al. 2022). Next we categorized each species as belonging to one of three ecological ~~groups (ground, rock, or tree), based on descriptions of habitat use found in the literature (Tejero-Cicuéndez et al. 2021a)~~ habitat groups (ground-dwelling, rock-dwelling, or tree-arboreal). Habitat designations were based on substrate preferences and habitat use as found through extensive field observations described in the primary literature (Arnold 1993; Arnold 2009, and references therein). Finally, we obtained a phenotypic data set containing body size (snout-vent length: SVL) and eight linear measurements (Figure 1) that described overall body form: trunk length (TL), head length (HL), head width (HW), head height (HH), humerus length (Lhu), ulna length (Lun), femur length (Lfe), and tibia length (Ltb) (Tejero-Cicuéndez et al. 2021a). We restricted our study to those species represented by nine or more individuals; resulting in a dataset of 687 individuals from 25 species (individuals per species: ~~$\mu = 27$~~ mean=27; min = 9, max = 56). All specimens used in this study were adults, and thus patterns of ontogenetic allometry could not be explored. Species in the phenotypic dataset were then matched to the phylogeny, which was subsequently pruned to the final topology. All measurements were log-transformed prior to statistical analyses. Additional details regarding data collection and formal descriptions of each linear measurement may be found in the original sources (Tejero-Cicuéndez et al. 2021a, 2022). The data are available on DRYAD: <https://doi.org/10.5061/dryad.xwdbrv1f6> (Tejero-Cicuéndez et al. 2021b).

(b) Statistical and Comparative Analyses

We conducted a series of analyses to interrogate allometric trends, patterns of integration, and macroevolutionary changes in allometry, relative to differentiation in body form. First we characterized evolutionary allometry in the genus by performing a phylogenetic multivariate regression of body form on body size (i.e., SVL), using the species means as data. We then ~~performed an analogous procedure at the individual level, regressing body form on body size using our entire dataset~~ evaluated patterns of static allometry among individuals using a pooled within-species regression (sensu Klingenberg 2016). Here, the residual values of individuals from their respective species mean were obtained, which were then pooled across species and used in a multivariate regression to obtain an overall estimate of static allometry among individuals. By first removing species-specific differences, this procedure partials out trends of evolutionary allometry from the data, enabling patterns of static and evolutionary allometry to be disentangled. From both the species-level (~~phylogenetic~~) and the individual-level analyses, we obtained the set of regression coefficients, ~~and which respectively described the trajectories of evolutionary and static allometry in morphospace.~~ We then calculated the difference in their ~~angular direction to describe direction in morphospace to discern~~ the extent to which patterns of static allometry at the individual level were concordant with evolutionary allometric trends across species.

Next we used the ~~dataset containing all individuals~~ pooled within-species dataset to determine whether trends in static allometry differed across habitat groups. This was accomplished by performing a multivariate analysis of covariance, with body size (*SVL*), *habitat*, and $SVL \times habitat$ as model effects. Significance of model effects was evaluated using 999 iterations of a permutation procedure, where residuals from a reduced model were randomly permuted in each permutation (RRPP), model statistics were recalculated, and used to generate empirical null sampling distributions to evaluate the observed test statistics (following Freedman and Lane 1983; Collyer and Adams 2007; Collyer et al. 2015). ~~We then~~ In this analysis, no variation was attributable to the *habitat* effect, as the pooled-within species data are mean-centered for both the dependent and independent variables. However, any differences in multivariate allometric slopes among habitats will be discernable, and revealed by a significant $SVL \times habitat$ interaction. To evaluate this

possibility, we compared the multivariate allometric vectors for each habitat group to one another, and to a vector representing multivariate isometry, by calculating pairwise angular differences in their ~~angular~~ direction in morphospace, and evaluating these relative to empirical sampling distributions obtained through RRPP (Collyer and Adams 2007; Adams and Collyer 2009; Collyer and Adams 2013). Here, residuals were obtained from a common isometry reduced model, whose common slope component described a pattern of multivariate isometry, and whose intercepts allowed for differences in least-squares means among groups. Patterns of multivariate allometry relative to body size were visualized via regression scores (Drake and Klingenberg 2008) and predicted lines (Adams and Nistri 2010), based on the coefficients and fitted values from the linear model described above.

Additionally, because allometry describes the extent to which traits covary with body size and with each other (i.e., integration), we conducted an analysis of integration. Here we characterized the extent of morphological integration in body form for individuals within each habitat group using the pooled within-species dataset, and by summarizing the dispersion of eigenvalues of their respective trait covariance matrix (Pavlicev et al. 2009). This measure (V_{rel}) was subsequently converted to an effect size (a Z-score), which quantified the strength of morphological integration (Conaway and Adams 2022). We then performed a series of two-sample tests to compare the strength of morphological integration across habitat groups, following the procedures of Conaway and Adams (2022). Additionally and for comparison, we repeated these analyses on the set of size-standardized trait data, found as a set of shape ratios (Mosimann 1970) where each trait was divided by body size (Supplementary Material).

To determine the extent to which static and evolutionary allometry were concordant, we evaluated the directions in morphospace of both the evolutionary (species-level) and static (habitat-based) allometric trends. Specifically, we obtained the set of regression coefficients from both the phylogenetic multivariate regression and the multivariate analysis of covariance analyses above, and calculated the ~~differences in angular~~ angular difference in direction between the evolutionary allometry trajectory and the static allometry trend for each habitat group. The observed angles

were then statistically evaluated relative to empirical sampling distributions obtained through permutation (RRPP), based on the common isometry model described above.

Next, to discern how allometric trends resulted in the evolution of distinct body forms, we examined changes in ~~the body shape proportions~~ relative body form across the phylogeny. Here we treated the head dimensions and limb dimensions separately, as allometric trends could potentially differ between these body regions due to differential functional or selective constraints (Kaliontzopoulou et al. 2010). Because both the head and limb data were multivariate, we ~~first performed a partial least squares (PLS) analysis (Rohlf and Corti 2000) of the~~ used regression scores (sensu Drake and Klingenberg 2008) of a multivariate regression of head traits versus SVL ~~and the~~ and limb traits versus SVL ~~to describe the direction of maximal covariation between each body region and size~~ to represent the allometric trends in each dataset. We then measured the mean residuals of each species to the inferred allometric trend, which described the extent to which head and limb proportions of species were greater or smaller than expected for their body size. The species residuals were then mapped on the phylogeny of *Pristurus* using a Brownian motion model of evolution, to qualitatively evaluate shifts in head and limbs proportionality across the phylogeny for the group. Similarly, within-species patterns of static allometry were visualized by plotting ~~regressions of PLS scores on~~ regression scores versus SVL for both head and limb traits separately.

Finally, to relate within-species allometric trends with patterns of phenotypic diversification in the group we generated a phylomorphospace (sensu Sidlauskas 2008), based on a ~~phylogenetic~~ principal component analyses (PCA) ~~on~~ of the size-standardized species means (i.e., relative body proportions) obtained from a phylogenetic regression (see Tejero-Cicuéndez et al. 2021a). Here, phenotypic similarities among species, relative to their phylogenetic relationships and habitat affiliations, were observed. ~~Additionally, representative specimens (sealed to unit size) were also visually compared to aid in describing these trends.~~ A similar phylomorphospace was constructed for species means not corrected for body size, and the phenotypic disparity among species means in each habitat was calculated and subsequently compared (Supplementary Material). Additionally, anatomical changes associated with allometric trends across taxa were visually depicted via representative specimens

from the largest and smallest ground-dwelling species (scaled to unit size), and specimens from a large and small rock-dwelling species, to aid in describing these allometric trends. All analyses were conducted in R 4.2.1 (R Core Team 2022), using RRPP version 1.3.1.2 (Collyer and Adams 2018; Collyer and Adams 2022) and geomorph 4.0.4.5 (Baken et al. 2021a) for statistical analyses and the tidyverse version 1.3.0 (Wickham et al. 2019), phytools version 0.7-77 (Revell 2012), and a modified version of the function ggphylomorpho [https://github.com/wabarr/ggphylomorpho] for data manipulation and visualization, as well as scripts written by the authors (Supplementary Material).

3. Results

Using phylogenetic regression, we found significant evolutionary allometry in body form across species ($N_{sp} = 25$; $F = 217.9$; $Z = 5.53$; $P < 0.001$). Likewise, when static allometry in body form was examined across individuals, a similar pattern was observed ($N_{ind} = 687$; ~~$F = 7910.8$~~ ; ~~$Z = 9.20$~~ $F = 1176.9$; $Z = 8.24$; $P < 0.001$). Further, the vectors of regression coefficients between the two analyses were ~~highly correlated ($\rho = 0.94$) and were oriented in nearly parallel directions~~ oriented in a similar direction and were nearly parallel in morphospace (~~$\theta = 1.49^\circ$~~ $\theta = 5.64^\circ$). This revealed that the pattern of multivariate allometry across individuals was largely concordant with macroevolutionary trends of interspecific static allometry among species of *Pristurus* across the phylogeny.

Our analyses also exposed significant differences in the allometry of body form among *Pristurus* utilizing distinct habitats (Table 1). Further, pairwise comparisons of multivariate allometric vectors revealed that patterns of static allometry in each habitat differed significantly from isometry, indicating the presence of multivariate allometry in each (Table 2). Additionally, comparisons identified that ground-dwelling *Pristurus* displayed the most distinct allometric trend as compared with *Pristurus* occupying both the rock and tree habitats (Table 2; Figure 2). Here, regression coefficients of each trait versus size (Supplementary Material) revealed that ground-dwelling *Pristurus* exhibited ~~strong positive allometry for all head and limb traits (i.e., $\beta > 1.0$)~~ higher coefficients for head traits as compared with rock-dwelling and tree-dwelling taxa

($\beta_{ground} > \beta_{rock}; \beta_{ground} > \beta_{tree}$). By contrast, ~~rock and tree-dwelling *Pristurus* displayed negative~~
~~allometry (i.e., $\beta < 1.0$) for head traits, and were more varied for limb traits; with rock-dwelling~~
~~*Pristurus* displaying positive limb allometry (though less extreme than that of~~
~~coefficients for limb~~
~~traits were somewhat smaller for~~ ground-dwelling taxa), whereas most limb traits in tree-dwelling
~~taxa showed negative allometry or near-isometry (Supplementary Material~~
~~*Pristurus* as compared~~
~~with other taxa ($\beta_{ground} < \beta_{rock}; \beta_{ground} < \beta_{tree}$)). Thus, these findings implied that~~
~~within species,~~
~~larger individuals of ground-dwelling *Pristurus* species displayed disproportionately~~
~~displayed~~
~~proportionately~~ larger heads and limbs, slightly smaller limbs as compared with large individuals in
taxa utilizing other habitat types. ~~Multivariate visualizations of these multivariate~~
~~Visualizations~~
~~of the~~ allometric trends (Figure 2) confirmed these statistical findings, and indicated that the
allometric trajectory in ground-dwelling *Pristurus* was more extreme as compared with either rock-
or tree-dwelling *Pristurus*.

Examination of patterns of trait covariation ~~for the pooled within-species data~~ revealed strong levels
of morphological integration ~~within each habitat type ($Z_{ground} = 3.97$; $Z_{rock} = 3.72$; $Z_{tree} = 2.15$).~~
~~Further, in the ground and tree ecotypes, with lower levels of integration displayed in the rock~~
~~habitat. Subsequent~~ two-sample tests revealed that the strength of morphological integration
was significantly greater in ~~both~~ ground-dwelling ~~and tree-dwelling~~ *Pristurus* than ~~either in~~
those utilizing rock (~~$Z_{ground-rock} = 6.59$~~ $Z_{ground-rock} = 6.05$; $P < 0.001$) ~~or tree habitats~~
(~~$Z_{ground-tree} = 11.17$; $Z_{tree-rock} = 4.07$~~ $Z_{tree-rock} = 4.07$; $P < 0.001$). ~~Arboreal *Pristurus* displayed the lowest~~
~~levels of integration, which were also significantly lower than in the rock habitat ($Z_{rock-tree} = 7.19$;~~
 ~~$P < 0.001$)). When size was accounted for in the data~~
~~Levels of morphological integration did not~~
~~differ between ground and tree-dwelling *Pristurus* ($Z_{tree-rock} = 0.38$; $P = 0.702$)). Finally, when~~
~~body size was taken into account,~~ levels of integration dropped considerably, though the overall
pattern and differences among habitat groups remained the same (Supplementary Material).

Comparisons of evolutionary allometry with static allometry in each habitat revealed substantial
concordance between allometric trends at these hierarchical levels. Here, vectors of regression
coefficients representing static allometry within habitat groups were oriented in very similar

directions with the regression vector representing evolutionary allometry, with small pairwise angles between them ($\theta: 2.3^\circ \rightarrow 5.9^\circ$ $\theta: 5.8^\circ \rightarrow 7.2^\circ$). Subsequent permutation tests indicated no differences ~~between the static allometry vectors and the~~ in direction between the regression vector representing evolutionary allometry and the static allometry vectors for *Pristurus* in both the ground or tree habitats, indicating strong congruence between them (Table 3). By contrast, rock-dwelling *Pristurus* differed most in their static allometry trend relative to patterns of evolutionary allometry. Notably, static allometry in ground-dwelling *Pristurus* was most similar to trends of evolutionary allometry, displaying the smallest angular difference ~~and largest effect size.~~ ~~Thus, static and evolutionary allometry trends were essentially parallel in this group, indicating a direct correspondence between the two. This result~~ in direction when compared to evolutionary allometry. Overall, these findings implied that phenotypic evolution across species aligned closely with directions of allometric variation within habitat groups at the individual level; ~~namely that larger individuals~~, describing a trend where larger individuals – and larger ground-dwelling species – exhibited disproportionately larger heads and limbs, while smaller individuals – and smaller ground-dwelling species – displayed disproportionately smaller heads and limbs.

Mapping the residuals of ~~species into~~ phylogenetic regression onto the phylogeny showed that large ground-dwelling species displayed greater head proportions than large rock-dwelling species, who exhibited smaller heads relative to body size (Figure 3A). Conversely, the opposite pattern was observed when comparing small species utilizing these habitats: ground-dwelling species showed small relative head proportions while rock-dwelling species displayed generally larger head proportions. In contrast, limb shape showed more variable patterns. Although all large ground-dwelling species consistently displayed large relative limb proportions, large rock-dwelling species were more variable in this trait, with *P. insignis* exhibiting large and *P. insignoides* small limb proportions. For small species, shifts in relative limb proportions seemed more independent of habitat utilization, since there were differences in limb residuals both within rock- and ground-dwelling species (Figure 3B). ~~Visual inspection of~~ Likewise, static allometry trends within species ~~(Figure 4) largely confirmed these patterns, illustrating~~ revealed that ground-dwelling species generally displayed steeper allometric patterns in head proportions as compared with

rock-dwelling species (Figure 4). Overall there was general concordance across taxa in terms of trends of multivariate allometry, affirming that the association between evolutionary allometry and habitat-based static allometry was robust.

Viewing ~~body shape differentiation in~~ differentiation in relative body proportions in *Pristurus* in phylomorphospace (Figure 5) revealed broad overlap among habitat groups in the first few dimensions, though arboreal (tree-dwelling) species were somewhat more ~~separated~~ constrained in morphospace. Rock-dwelling species occupied a slightly larger region of morphospace as compared with the other groups, though this pattern was not statistically significant (Supplementary Material). Intriguingly, when viewed in relation to body size, large *Pristurus* species were not localized to a particular region of morphospace, nor were smaller species. Instead, the largest rock-dwelling species were found in close proximity to the smallest ground-dwelling species, indicating that they were similar in ~~overall body shape~~ relative body proportions. Likewise, the smallest rock-dwelling species were found close to large ground-dwelling species in morphospace, indicating they displayed similar body ~~shapes~~ proportions as well.

Finally, when representative specimens were scaled to a similar body size (Figure 6), the anatomical consequences of differences in allometric trends on body ~~proportions~~ form became apparent. Here, larger ground-dwelling *Pristurus* species displayed disproportionately larger heads and limbs as compared with large *Pristurus* species utilizing other habitat types. Conversely, smaller rock-dwelling species were found to have disproportionately larger heads and limbs as compared with smaller ground-dwelling species. These patterns corresponded closely with those identified in morphospace (Figure 5), where large ground-dwelling species were similar in body form to small rock-dwelling species, while small ground-dwelling species were similar in body form to large rock-dwelling species (Figure 6). Thus, synthesizing the patterns revealed in the phylomorphospace with those from the other analyses revealed that the same body ~~shape~~ proportions could be obtained in different ways, as determined by subtle differences in allometric slope across habitats, combined with body size differences. As such, species with similar body ~~shapes~~ proportions displayed differing overall size, were found in distinct habitats, and exhibited different allometric trends.

4. Discussion

Elucidating the selective forces that generate patterns of phenotypic diversity is a major goal in evolutionary biology. For species that utilize distinct habitats, disentangling the causes of phenotypic differentiation across those habitats is essential for our understanding of how natural selection operates and how evolution proceeds. In this study, we evaluated the role of potential drivers of body ~~shape~~ form differentiation in the geckos of the genus *Pristurus*. To this end, we compared allometric trends and levels of integration among *Pristurus* occupying distinct habitats, interrogated allometric patterns at both the static and evolutionary levels, and related these trends to diversification in body form. Our findings have several important implications for how ecological specialization, phenotypic integration, and body form evolution along allometric trajectories relate to patterns of phenotypic diversity generally, and the evolution of phenotypic diversification in *Pristurus* in particular.

REv 1: “much of this rehash from above”. REWORD??? First, our analyses revealed that patterns of ~~body-shape-allometry~~ allometry in body form and morphological integration are relatively distinct in ground-dwelling *Pristurus* lizards, as compared with *Pristurus* occupying other habitats. Specifically, we found that multivariate vectors of regression coefficients differed significantly from what was expected under isometry (Table 2) for taxa utilizing all habitat types (ground, rock, tree), indicating that in *Pristurus*, allometric scaling patterns predominate. Further, our interrogation of allometric trends revealed differences between habitat types, where ground-dwelling *Pristurus* displayed steeper ~~(i.e., positively allometric) trends for both head and limb traits~~ while allometric trends for head traits as compared with rock and tree-dwelling taxa ~~displayed shallower (negatively allometric) trends for head traits and more varied patterns for limb proportions~~. Biologically, these patterns revealed that not only does shape differ between large and small *Pristurus*, but this pattern differs across habitat types. Specifically, large ground-dwelling *Pristurus* present disproportionately larger heads ~~and longer limbs~~ relative to large individuals in other habitats, while small ground-dwelling *Pristurus* exhibit disproportionately smaller heads ~~and shorter limbs~~ (Figure 3). These findings are consistent with previous work at the macroevolutionary

level (Tejero-Cicuéndez et al. 2021a), where large ground species were also found to display disproportionately large heads and long limbs. NOTE: DCA REMOVED THE LIMBS PART

Second, our findings revealed that within species, rock-dwelling *Pristurus* show a converse pattern, where smaller individuals displayed relatively larger heads, while larger individuals have smaller heads relative to their body size. These allometric patterns also corresponded with findings at macroevolutionary scales (Tejero-Cicuéndez et al. 2021a), where similar patterns at the species level were observed. Regarding relative limb proportions, we found a high variability among small rock-dwelling species rather than a common pattern (Figure 3B). Indeed, earlier work in the subclade comprising several of these species (the *P. rupestris* species complex) found two well-differentiated phenotypes in populations of these lineages segregated by elevation (Garcia-Porta et al. 2017). These two ecotypes, defined as ‘slender’ and ‘robust’, differed in their head and limb characteristics. Our work is consistent with this, and extends these patterns to the allometric realm. Tejero-Cicuéndez et al. (2021a) also performed habitat ancestral estimation, finding that the rock habitat was the most likely ancestral condition in the group, with subsequent colonization by *Pristurus* of ground habitats. When patterns of allometry are viewed through this lens, it suggests the hypothesis that habitat shifts from rock-dwelling to ground-dwelling incurred a concomitant evolutionary shift in allometric trajectories as well (Adams and Nistri 2010). Indeed, our analyses are consistent with this hypothesis, as allometric trends are inferred to be more rock-like towards the root of the *Pristurus* phylogeny (Figure 3), with subsequent shifts along branches leading to ground-dwelling species. This further suggests that the segregation in body size and shape through differential allometric relationships across habitats responds to adaptive dynamics concerning the colonization of new habitats. Thus, in *Pristurus*, there is support for the hypothesis that colonization of ground habitats has been a trigger for morphological change (Tejero-Cicuéndez et al. 2021a), as there appears to be a link between shifts in allometric trajectories as a result of habitat-induced selection, and differential patterns of body ~~shape~~-form observed across taxa. More broadly, these findings are consistent with prior discoveries in other lizards, where the differential selective pressures imposed by rocky and ground habitats have resulted in the

differentiation of head and limb morphology (Goodman et al. 2008; Kaliontzopoulou et al. 2010; Garcia-Porta et al. 2017; Foster et al. 2018). Indeed, such phenotypic differences resulting from the effects of habitat-based ecological selection have been extensively documented in reptiles as well as in other vertebrates (Losos 2009; Reynolds et al. 2016; Hipsley and Müller 2017; Samuels and Hopkins 2017; Watanabe et al. 2019; Busschau and Boissinot 2022; Navalón et al. 2022; Friedman et al. 2022), and our work in *Pristurus* thus contributes to this growing body of literature.

Biologically, the findings revealed here may reflect ecological and behavioral changes linked to the adoption of a new lifestyle. For lizards, the transition to utilizing ground habitats implies adopting an existence in more open environments than in rocky substrates. As such, numerous aspects of daily existence (including activity patterns, climatic factors, prey availability, abundance of predators, etc.) are expected to exert a differential influence on an organism's phenotype when compared with life in their ancestral environment (Fuentes and Cancino (1979)). Indeed, the largest ground-dwelling *Pristurus* species (*P. carteri*, *P. ornitocephalus*, and *P. collaris*) differ from the rest of the genus in having developed partially nocturnal habits, which would presumably have major ecological consequences for their survival and reproduction. In this sense, these species might have been subjected to evolutionary processes selecting for larger relative head proportions, which would allow them to accommodate larger or modified eyes, a clear advantage in animals with nocturnal and semi-nocturnal habits (Hall and Ross 2006; Ross et al. 2007; Hall et al. 2012). Likewise, the steep allometric patterns found in the limbs of large ground-dwelling species (Figure 3B) might be related to selective processes favoring longer limbs in large species present in this new ecological context. Longer limbs in open habitats might be advantageous for active prey foraging and, in hyper-arid areas such as the Arabian Peninsula, this morphology might contribute to thermoregulation separating the body from the ground (Avery et al. (1982); Huey (1974)). A more detailed examination of behavioral and morphological traits (e.g., eye shape) might shed light on the factors driving this pattern.

Another important finding of our study was the strong concordance between static allometry across individuals and evolutionary allometry among *Pristurus* species. Our analyses revealed

small pairwise angles between static and evolutionary allometry vectors, indicating that allometric trends at these two ~~hierarchical~~ levels were oriented in similar directions and were ~~essentially~~ parallel largely concordant. As such, size-associated changes in body ~~shape-form~~ among individuals were predictive of evolutionary shifts across taxa at higher macroevolutionary scales. This in turn, suggests that body ~~shape-form~~ evolution in *Pristurus* ~~follows~~ may follow an allometric line of least resistance (Marroig and Cheverud 2005). In other empirical systems, a similarly tight correspondence between static and evolutionary allometry has also been observed (Marroig and Cheverud 2005; Firmat et al. 2014; Voje et al. 2014; Brombacher et al. 2017; Marcy et al. 2020), though the trend is not universal across all taxa or traits (see Klingenberg and Zimmermann 1992; Voje et al. 2022). Nonetheless, when such trends are present, they imply that allometric trajectories impose a prevailing influence on the magnitude, direction, and rate of phenotypic change across the phylogeny. Our work in *Pristurus* contributes to the growing literature on this topic, and suggests that perhaps such patterns may be more widespread.

Given the observation that static and evolutionary allometry in *Pristurus* are ~~so~~ largely concordant, an obvious question is: why might this be the case? One possible explanation is that when genetic covariation remains relatively constant, selection on body size will generate an evolutionary allometric trajectory along the trend described by static allometry (Lande 1979, 1985). Here, allometry effectively acts as a constraint on evolutionary change, as size-associated shape changes at one hierarchical level are linked to changes at another level (Voje et al. 2014, 2022; Pélabon et al. 2014). Further, when this is the case, one may also expect high levels of phenotypic integration in traits associated with body size changes. Indeed, our analyses reveal precisely this pattern in *Pristurus*, with the highest levels of integration in the group (ground-dwelling) whose static allometry is most similar to that of evolutionary allometry. Thus, our results reveal that patterns of trait covariation are more constrained in ground-dwelling species, such that their differences in body form are most likely found along the primary allometric axis. When viewed in this light, integration and allometry may thus be interpreted as potential drivers that facilitate morphological change, as they provide a phenotypic pathway through adaptive lines of least resistance that enable rapid evolutionary changes in particular phenotypic directions but not in others (Felice et al. 2018;

Navalón et al. 2020). The fact that ground-dwelling species in *Pristurus* have been found to have the widest phenotypic disparity, greatest range of body sizes, and highest rates of morphological evolution (Tejero-Cicuéndez et al. 2021a) are all consistent with this hypothesis, and suggest that in this group, integration describes the path of morphological evolution along allometric lines of least resistance.

Finally, interpreting the observed patterns of phenotypic integration and allometry relative to habitat-specific differences helps to shed light on the possible pathways by which phenotypic diversity in *Pristurus* has evolved. For instance, prior work on this system (Tejero-Cicuéndez et al. 2021a) revealed that the colonization of new ecological habitats elicited strong ecological selection and phenotypic responses. This was particularly true of the invasion of ground habitats, where ground-dwelling species displayed the largest variation in body size in the genus. This observation implies some level of ecological selection on body size. In lizards, the ecological context in which species exist is known to play a pervasive role in body size evolution (James and M'closkey 2004; Meiri 2008; Tamar et al. 2019), as it does in other animal groups (Bergmann 1847; Calder 1983; Peters 1983; LaBarbera 1989; Olson et al. 2009). While to date this has not been thoroughly explored in *Pristurus*, the evolutionary patterns revealed by our analyses suggest that the body size diversity in this clade conforms, at least in part, with patterns expected under ecological selection on body size. Intriguingly, such patterns are not only observed in ground- and rock-dwelling taxa, but also in arboreal species, whose restricted phenotypic diversity in both size and shape (Figures 3 & 5) is consistent with strong ecological selection in the arboreal habit (Baken and Adams 2019; Baken et al. 2021b). Furthermore, our study identified the presence of strong integration and allometric trajectories, such that evolutionary changes in body size elicit corresponding changes in body [shapeform](#). However, these trends differed significantly across habitats, implying that, at evolutionary scales, these trends serve to channel phenotypic responses to selection, but do so in differing directions for the different habitat groups. This, in turn, suggests that *Pristurus* species occupying different habitats display differing combinations of body size with body [shapeform](#). The evolutionary consequence of ecological selection is that species have evolved similar shapes (Figure 6), but do so in differing habitats, and at different body sizes (Figure 5). Therefore, the

493 phenotypic diversity observed in *Pristurus* is best explained as the result of a complex interplay
494 between ecological selection, body size differentiation, and differing allometric trajectories across
495 ecological habitats.

References

- Adams, D. C., and M. L. Collyer. 2009. A general framework for the analysis of phenotypic trajectories in evolutionary studies. *Evolution* 63:1143–1154.
- Adams, D. C., and A. Nistri. 2010. Ontogenetic convergence and evolution of foot morphology in European cave salamanders (Family: Plethodontidae). *BMC Evolutionary Biology* 10:1–10. BioMed Central.
- Albertson, R. C., and T. D. Kocher. 2001. Assessing morphological differences in an adaptive trait: A landmark-based morphometric approach. *Journal of Experimental Zoology* 289:385–403.
- [Arnold, E. N. 1993. Historical changes in the ecology and behaviour of semaphore geckos \(*Pristurus*, Gekkonidae\) and their relatives. *Journal of Zoology* 229:353–384.](#)
- [Arnold, E. N. 2009. Relationships, evolution and biogeography of semaphore geckos, *Pristurus* \(Squamata, Sphaerodactylidae\) based on morphology. *Zootaxa* 2060:1–21.](#)
- [Arnold, S. J. 1983. Morphology, performance, fitness. *American Zoologist* 23:347–361.](#)
- [Avery, R. A., Bedford J. D., and C. P. Newcombe. 1982. The role of thermoregulation in lizard biology: Predatory efficiency in a temperate basker. *Behavioral Ecology and Sociobiology* 11:261–267.](#)
- [Baken, E. K., and D. C. Adams. 2019. Macroevolution of arboreality in salamanders. *Ecology and Evolution* 9:7005–7016.](#)
- [Baken, E. K., M. L. Collyer, A. Kaliontzopoulou, and D. C. Adams. 2021a. Geomorph 4.0 and gmShiny: Enhanced analytics and a new graphical interface for a comprehensive morphometric experience. *Methods in Ecology and Evolution* 12:2355–2363.](#)
- [Baken, E. K., L. E. Mellenthin, and D. C. Adams. 2021b. Is salamander arboreality limited by broad-scale climatic conditions? *PLoS ONE* 16:e0255393.](#)
- [Bergmann, C. 1847. Über die verhältnisse der warmekonomie der thiere zu ihrer grosse. *Göttinger Studien* 1:595–708.](#)
- [Berner, D., D. C. Adams, A.-C. Grandchamp, and A. P. Hendry. 2008. Natural selection drives patterns of lake-stream divergence in stickleback foraging morphology. *Journal of*](#)

Evolutionary Biology 21:1653–1665.

Bookstein, F. L. 2022. Dimensions of morphological integration. *Evolutionary Biology* 49:342–372.

Brombacher, A., P. A. Wilson, I. Bailey, and T. H. G. Ezard. 2017. The breakdown of static and evolutionary allometries during climatic upheaval. *The American Naturalist* 190:350–362.

Busschau, T., and S. Boissinot. 2022. Habitat determines convergent evolution of cephalic horns in vipers. *Biological Journal of the Linnean Society* 135:652–664.

Calder, W. A. 1983. Ecological scaling: Mammals and birds. *Annual Review of Ecology and Systematics* 14:213–230.

Collar, D. C., J. A. Schulte, B. C. O’Meara, and J. B. Losos. 2010. Habitat use affects morphological diversification in dragon lizards. *Journal of Evolutionary Biology* 23:1033–1049.

Collyer, M. L., and D. C. Adams. 2007. Analysis of two-state multivariate phenotypic change in ecological studies. *Ecology* 88:683–692.

Collyer, M. L., and D. C. Adams. 2013. Phenotypic trajectory analysis: Comparison of shape change patterns in evolution and ecology. *Hystrix* 24:75–83.

Collyer, M. L., and D. C. Adams. 2022. R: RRPP: Linear model evaluation with randomized residuals in a permutation procedure. Vsn. 1.3-~~1~~.2. R Foundation for Statistical Computing, Vienna, Austria.

Collyer, M. L., and D. C. Adams. 2018. RRPP: An R package for fitting linear models to high-dimensional data using residual randomization. *Methods in Ecology and Evolution* 9:1772–1779.

Collyer, M. L., D. J. Sekora, and D. C. Adams. 2015. A method for analysis of phenotypic change for phenotypes described by high-dimensional data. *Heredity* 115:357–365.

Conaway, M. A., and D. C. Adams. 2022. An effect size for comparing the strength of morphological integration across studies. *Evolution* 76:2244–2259.

Drake, A. G., and C. P. Klingenberg. 2008. The pace of morphological change: Historical transformation of skull shape in St Bernard dogs. *Proceedings of the Royal Society B: Biological Sciences* 275:71–76.

Durant, S. M., N. Pettorelli, S. Bashir, R. Woodroffe, T. Wachter, P. De Ornellas, C. Ransom,

- T. Abáigar, M. Abdelgadir, H. El Alqamy, M. Beddiaf, F. Belbachir, A. Belbachir-Bazi, A. A. Berbash, R. Beudels-Jamar, L. Boitani, C. Breitenmoser, M. Cano, P. Chardonnet, B. Collen, W. A. Cornforth, F. Cuzin, P. Gerngross, B. Haddane, M. Hadjeloum, A. Jacobson, A. Jebali, F. Lamarque, D. Mallon, K. Minkowski, S. Monfort, B. Ndoassal, J. Newby, B. E. Ngakoutou, B. Niagate, G. Purchase, S. Samaïla, A. K. Samna, C. Sillero-Zubiri, A. E. Sulttan, M. R. Stanley Price, and J. E. M. Baillie. 2012. Forgotten biodiversity in desert ecosystems. *Science* 336:1379–1380.
- Felice, R. N., M. Randau, and A. Goswami. 2018. A fly in a tube: Macroevolutionary expectations for integrated phenotypes. *Evolution* 72:2580–2594.
- Firmat, C., I. Lozano-Fernández, J. Agustí, G. H. Bolstad, G. Cuenca-Bescós, T. F. Hansen, and C. Pélabon. 2014. Walk the line: 600000 years of molar evolution constrained by allometry in the fossil rodent *Mimomys savini*. *Philosophical Transactions of the Royal Society B: Biological Sciences* 369:20140057.
- Foster, K. L., T. Garland, L. Schmitz, and T. E. Higham. 2018. Skink ecomorphology: Forelimb and hind limb lengths, but not static stability, correlate with habitat use and demonstrate multiple solutions. *Biological Journal of the Linnean Society* 125:673–692.
- Freedman, D., and D. Lane. 1983. A nonstochastic interpretation of reported significance levels. *Journal of Business & Economic Statistics* 1:292–298.
- Friedman, S. T., M. L. Collyer, S. A. Price, and P. C. Wainwright. 2022. Divergent Processes Drive Parallel Evolution in Marine and Freshwater Fishes. *Systematic biology* 71:1319–1330.
- Fuentes, E. R., and J. Cancino. 1979. Rock-ground patchiness in a simple liolaemus lizard community (reptilia, lacertilia, iguanidae). *Journal of Herpetology* 13:343.
- Garcia-Porta, J., M. Simó-Riudalbas, M. Robinson, and S. Carranza. 2017. Diversification in arid mountains: Biogeography and cryptic diversity of *Pristurus rupestris rupestris* in Arabia. *Journal of Biogeography* 44:1694–1704.
- Goodman, B. A., D. B. Miles, and L. Schwarzkopf. 2008. Life on the rocks: Habitat use drives morphological and performance evolution in lizards. *Ecology* 89:3462–3471.
- Goswami, A., M. Randau, P. D. Polly, V. Weisbecker, C. Verity Bennett, L. Hautier, and M. R. Sánchez-Villagra. 2016. Do developmental constraints and high integration limit the

evolution of the marsupial oral apparatus? Integrative and Comparative Biology 56:404–415.

Goswami, A., J. B. Smaers, C. Soligo, and P. D. Polly. 2014. The macroevolutionary consequences of phenotypic integration: From development to deep time. Philosophical Transactions of the Royal Society B: Biological Sciences 369:20130254.

Grant, P. R., and B. R. Grant. 2006. Evolution of character displacement in darwin’s finches. Science 313:224–226.

Hall, M. I., J. M. Kamilar, and E. C. Kirk. 2012. Eye shape and the nocturnal bottleneck of mammals. Proceedings of the Royal Society B: Biological Sciences 279:4962–4968.

Hall, M. I., and C. F. Ross. 2006. Eye shape and activity pattern in birds. Journal of Zoology 271:437–444.

Hipsley, C. A., and J. Müller. 2017. Developmental dynamics of ecomorphological convergence in a transcontinental lizard radiation. Evolution 71:936–948.

Huey, R. B. 1974. Behavioral thermoregulation in lizards: Importance of associated costs. Science 184:1001–1003.

James, S. E., and R. T. M’closkey. 2004. Patterns of body size and habitat use in a lizard assemblage. Ecoscience 11:160–167.

Jastrebski, C. J., and B. W. Robinson. 2004. Natural selection and the evolution of replicated trophic polymorphisms in pumpkinseed sunfish (*Lepomis gibbosus*). Evolutionary Ecology Research 6:285–305.

Jolicoeur, P. 1963. The multivariate generalization of the allometry equation. Biometrics 19:497–499.

Kaliontzopoulou, A., M. A. Carretero, and D. C. Adams. 2015. Ecomorphological variation in male and female wall lizards and the macroevolution of sexual dimorphism in relation to habitat use. Journal of Evolutionary Biology 28:80–94.

Kaliontzopoulou, A., M. A. Carretero, and G. A. Llorente. 2010. Intraspecific ecomorphological variation: Linear and geometric morphometrics reveal habitat-related patterns within *Podarcis bocagei* wall lizards. Journal of Evolutionary Biology 23:1234–1244.

Kappelman, J., D. T. Rasmussen, W. J. Sanders, M. Feseha, T. Bown, P. Copeland, J.

- Crabaugh, J. Fleagle, M. Glantz, A. Gordon, B. Jacobs, M. Maga, K. Muldoon, A. Pan,
L. Pyne, B. Richmond, T. Ryan, E. R. Seiffert, S. Sen, L. Todd, M. C. Wiemann, and
A. Winkler. 2003. Oligocene mammals from Ethiopia and faunal exchange between
Afro-Arabia and Eurasia. *Nature* 426:549–552.
- Klingenberg, C. P. 2016. Size, shape, and form: Concepts of allometry in geometric
morphometrics. *Development Genes and Evolution* 226:113–137.
- Klingenberg, C. P., and J. Marugán-Lobón. 2013. Evolutionary covariation in geometric
morphometric data: Analyzing integration, modularity, and allometry in a phylogenetic
context. *Systematic Biology* 62:591–610.
- Klingenberg, C. P., and M. Zimmermann. 1992. Static, ontogenetic, and evolutionary allometry:
A multivariate comparison in nine species of water striders. *American Naturalist* 140:601–620.
- Kolmann, M. A., F. P. L. Marques, J. C. Weaver, M. N. Dean, J. P. Fontenelle, and N. R.
Lovejoy. 2022. Ecological and phenotypic diversification after a continental invasion in
neotropical freshwater stingrays. *Integrative and Comparative Biology* 62:424–440.
- LaBarbera, M. 1989. Analyzing body size as a factor in ecology and evolution. *Annual Review
of Ecology and Systematics* 20:97–117.
- Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain-body
size allometry. *Evolution* 33:402–416.
- Lande, R. 1985. Size and scaling in primate biology. Pp. 21–32 in W. L. Jungers, ed. Plenum
Press.
- Losos, J. B. 2009. Lizards in an evolutionary tree: Ecology and adaptive radiation of anoles.
University of California Press.
- Losos, J. B. 1992. The evolution of convergent structure in Caribbean *Anolis* communities.
Systematic Biology 41:403–420.
- Losos, J. B., and R. E. Ricklefs. 2009. Adaptation and diversification on islands. *Nature*
457:830–836.
- Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the
macroevolutionary landscape in island lizard radiations. *Science* 341:292–295.

- 639 Marcy, A. E., T. Guillaume, E. Sherratt, K. C. Rowe, M. J. Phillips, and V. Weisbecker. 2020.
640 Australian rodents reveal conserved cranial evolutionary allometry across 10 million years of
641 murid evolution. *The American Naturalist* 196:755–768.
- 642 Marroig, G., and J. M. Cheverud. 2005. Size as a line of least evolutionary resistance: Diet and
643 adaptive morphological radiation in New World monkeys. *Evolution* 59:1128–1142.
- 644 Martinez, C. M., S. T. Friedman, K. A. Corn, O. Larouche, S. A. Price, and P. C. Wainwright.
645 2021. The deep sea is a hot spot of fish body shape evolution. *Ecology Letters* 24:1788–1799.
- 646 Meiri, S. 2008. Evolution and ecology of lizard body sizes. *Global Ecology and Biogeography*
647 17:724–734.
- 648 Mosimann, J. E. 1970. Size allometry: Size and shape variables with characterizations of
649 the lognormal and generalized gamma distributions. *Journal of the American Statistical*
650 *Association* 65:930–945.
- 651 Navalón, G., A. Bjarnason, E. Griffiths, and R. B. J. Benson. 2022. Environmental signal in the
652 evolutionary diversification of bird skeletons. *Nature* 611:306–311.
- 653 Navalón, G., J. Marugán-Lobón, J. A. Bright, C. R. Cooney, and E. J. Rayfield. 2020. The
654 consequences of craniofacial integration for the adaptive radiations of Darwin’s finches and
655 Hawaiian honeycreepers. *Nature Ecology & Evolution* 4:270–278. Nature Publishing Group.
- 656 Olson, V. A., R. G. Davies, C. D. L. Orme, G. H. Thomas, S. Meiri, T. M. Blackburn, K. J.
657 Gaston, I. P. F. Owens, and P. M. Bennett. 2009. Global biogeography and ecology of body
658 size in birds. *Ecology Letters* 12:249–259.
- 659 Pavlicev, M., J. M. Cheverud, and G. P. Wagner. 2009. Measuring morphological integration
660 using eigenvalue variance. *Evolutionary Biology* 36:157–170.
- 661 Pélabon, C., G. H. Bolstad, C. K. Egset, J. M. Cheverud, M. Pavlicev, and G. Rosenqvist. 2014.
662 On the relationship between ontogenetic and static allometry. *The American Naturalist*
663 181:195–212.
- 664 Peters, R. H. 1983. The ecological implications of body size. Cambridge University Press.
- 665 Price, S. A., S. T. Friedman, and P. C. Wainwright. 2015. How predation shaped fish: The
666 impact of fin spines on body form evolution across teleosts. *Proceedings of the Royal Society*
667 *B: Biological Sciences* 282:20151428.
- 668 R Core Team. 2022. R: A language and environment for statistical computing. Version 4.2.1. R

- Foundation for Statistical Computing, Vienna, Austria.
- Reaney, A. M., Y. Bouchenak-Khelladi, J. A. Tobias, and A. Abzhanov. 2020. Ecological and morphological determinants of evolutionary diversification in Darwin's finches and their relatives. *Ecology and Evolution* 10:14020–14032.
- Revell, L. J. 2012. Phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3:217–223.
- Reynolds, R. G., D. C. Collar, S. A. Pasachnik, M. L. Niemiller, A. R. Puente-Rolón, and L. J. Revell. 2016. Ecological specialization and morphological diversification in Greater Antillean boas. *Evolution* 70:1882–1895.
- ~~Rohlf, F. J., and M. Corti. 2000. Use of two-block partial least-squares to study covariation in shape. *Systematic Biology* 49: 740–753.~~
- ~~Ross, C. F., M. I. Hall, and C. P. Heesy. 2007. Were basal primates nocturnal? Evidence from eye and orbit shape. Pp. 233–256 in *PRIMATE ORIGINS: Adaptations and evolution*. Springer.~~
- Samuels, J. X., and S. S. B. Hopkins. 2017. The impacts of Cenozoic climate and habitat changes on small mammal diversity of North America. *Global and Planetary Change* 149:36–52.
- Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. *Evolution* 50:1766–1774.
- Schluter, D., and P. R. Grant. 1984. Determinants of morphological patterns in communities of Darwin's finches. *The American Naturalist* 123:175–196.
- Schluter, D., and J. D. McPhail. 1992. Ecological character displacement and speciation in sticklebacks. *The American Naturalist* 140:85–108.
- Sidlauskas, B. 2008. Continuous and arrested morphological diversification in sister clades of characiform fishes: A phylomorphospace approach. *Evolution* 62:3135–3156.
- Stuart, Y. E., T. Veen, J. N. Weber, D. Hanson, M. Ravinet, B. K. Lohman, C. J. Thompson, T. pre
- Tasneem, A. Doggett, R. Izen, N. Ahmed, R. D. H. Barrett, A. P. Hendry, C. L. Peichel, and D. I. Bolnick. 2017. Contrasting effects of environment and genetics generate a continuum of parallel evolution. *Nature Ecology and Evolution* 1:158.
- Tamar, K., P. Mitsi, M. Simó-Riudalbas, H. Tejero-Cicuéndez, T. Al-Sariri, and S. Carranza. 2019. Systematics, biogeography, and evolution of *Pristurus minimus* (Squamata, Sphaero-

- dactylidae) with the discovery of the smallest Arabian vertebrate. *Systematics and Biodiversity* 17:349–366.
- Tejero-Cicuéndez, H., A. H. Patton, D. S. Caetano, J. Šmíd, L. J. Harmon, and S. Carranza. 2022. Reconstructing squamate biogeography in Afro-Arabia reveals the influence of a complex and dynamic geologic past. *Systematic Biology* 71:261–272.
- Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021a. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. *Proceedings of the Royal Society B: Biological Sciences* 288:20211821.
- Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021b. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. Dryad digital repository. (Doi:10.5061/dryad.xwdbvr1f6).
- Urban, S., J. Gerwin, C. D. Hulsey, A. Meyer, and C. F. Kratochwil. 2022. The repeated evolution of stripe patterns is correlated with body morphology in the adaptive radiations of East African cichlid fishes. *Ecology and Evolution* 12:e8568.
- Voje, K. L., M. A. Bell, and Y. E. Stuart. 2022. Evolution of static allometry and constraint on evolutionary allometry in a fossil stickleback. *Journal of Evolutionary Biology* 35:423–438.
- Voje, K. L., T. F. Hansen, C. K. Egset, G. H. Bolstad, and C. Pélabon. 2014. Allometric constraints and the evolution of allometry. *Evolution* 68:866–885.
- Wagner, G. P., and J. Zhang. 2011. The pleiotropic structure of the genotype–phenotype map: The evolvability of complex organisms. *Nature Reviews Genetics* 12:204–213.
- Wagner, G., and L. Altenberg. 1996. Perspective: Complex adaptations and the evolution of evolvability. *Evolution* 50:967–976.
- Watanabe, A., A. C. Fabre, R. N. Felice, J. A. Maisano, J. Müller, A. Herrel, and A. Goswami. 2019. Ecomorphological diversification in squamates from conserved pattern of cranial integration. *Proceedings of the National Academy of Sciences of the United States of America* 116:14688–14697.
- Wickham, H., M. Averick, J. Bryan, W. Chang, L. D. McGowan, R. François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. L. Pedersen, E. Miller, S. M. Bache, K. Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, and H. Yutani. 2019. Welcome to the tidyverse. *Journal of Open Source Software* 4:1686.

728

729

730 **Acknowledgments:** We are very grateful to J. Roca, M. Metallinou, K. Tamar, J. Šmíd, R.
731 Vasconcelos, R. Sindaco, F. Amat, Ph. de Pous, L. Machado, J. Garcia-Porta, J. Els, T. Mazuch,
732 T. Papenfuss, and all the people from the Environment Authority, Oman, for their help in
733 different aspects of the work.

734 **Funding Statement:** This work was funded in part by PGC2018-098290-B-I00 (MCI-
735 U/AEI/FEDER, UE) and PID2021-128901NB-I00 (MCIN/AEI/10.13039/501100011033 and by
736 ERDF, A way of making Europe), Spain to SC. HT-C is supported by a “Juan de la Cierva -
737 Formación” postdoctoral fellowship (FJC2021-046832-I). IM was funded by the Alexander von
738 Humboldt Foundation through a Humboldt Research Fellowship. AT is supported by the “la
739 Caixa” doctoral fellowship programme (LCF/BQ/DR20/11790007). GR was funded by an FPI
740 grant from the Ministerio de Ciencia, Innovación y Universidades, Spain (PRE2019-088729).
741 BB-C was funded by FPU grant from Ministerio de Ciencia, Innovación y Universidades, Spain
742 (FPU18/04742). DCA was funded in part by National Science Foundation Grant DBI-1902511.

743 **Data availability statement:** All the data used in this study are available on DRYAD from a
744 previous study: <https://doi.org/10.5061/dryad.xwdbrv1f6> (Tejero-Cicuéndez et al. 2021b). The
745 scripts for implementing all analyses and generating the figures in this manuscript can be found
746 in the Supplementary Material and in a GitHub repository (and on DRYAD upon acceptance).

747 **Competing interests:** The authors declare no competing interests.

Table 1: Multivariate analysis of covariance describing variation in body form in *Pristurus*. Note that there is no variation explained by the 'habitat' term, as the pooled-within species data are mean-centered.

	Df	SS	MS	Rsqr	F	Z	Pr(>F)
SVL <u>svl</u>	1	516.04 <u>36.04</u>	516.04 <u>36.04</u>	0.92 <u>0.63</u>	10188.70 <u>1177.2</u>	9.49 <u>8.24</u>	0.001
habitat	2	6.22 <u>0.00</u>	3.11 <u>0.00</u>	0.01 <u>0.00</u>	61.39 <u>0.0</u>	9.32	0.001
SVL <u>svl</u> :habitat	2	3.97 <u>0.13</u>	1.99 <u>0.06</u>	0.01 <u>0.00</u>	39.23 <u>2.1</u>	7.08 <u>1.90</u>	0.001 <u>0.025</u>
Residuals	681	34.49 <u>20.85</u>	0.05 <u>0.03</u>	0.06 <u>0.37</u>			
Total	686	560.72 <u>57.02</u>					

Table 2: Pairwise comparisons of multivariate static allometry for each habitat group. Comparisons with the vector of multivariate isometry are included. Displayed are: pairwise angular differences (θ_{12}), their associated effect sizes ($Z_{\theta_{12}}$), and significance levels obtained via permutation (RRPP).

	Ground	Rock	Tree	Isometry
Angle				
Ground	0			
Rock	6.629 <u>6.316</u>	0		
Tree	8.095 <u>6.549</u>	3.628 <u>3.37</u>	0	
Isometry	5.034 <u>5.87</u>	5.901 <u>9.319</u>	7.189 <u>8.774</u>	0
Effect Size				
Ground	0			
Rock	7.004 <u>3.112</u>	0		
Tree	2.1 <u>1.9</u>	-0.408 <u>-0.454</u>	0	
Isometry	7.673 <u>4.461</u>	7.357 <u>6.567</u>	1.779 <u>3.727</u>	0
P-value				
Ground	1			
Rock	0.001 <u>0.003</u>	1		
Tree	0.027 <u>0.026</u>	0.673 <u>0.67</u>	1	
Isometry	0.001	0.001	0.042 <u>0.001</u>	1

Table 3: Pairwise comparisons of multivariate evolutionary allometry versus static allometry for each habitat group. Pairwise angular differences between evolutionary and static allometry (θ_{ES}), their associated effect sizes ($Z_{\theta_{ES}}$), and significance levels are displayed.

	θ_{ES}	$Z_{\theta_{ES}}$	P-value
Evol. vs. Ground	2.37 <u>5.85</u>	-4.26 <u>-1.61</u>	1.000 <u>0.063</u>
Evol. vs. Rock	4.55 <u>7.23</u>	0.87 <u>-2.54</u>	0.191 <u>0.009</u>
Evol. vs. Tree	5.96 <u>6.79</u>	0.21 <u>-1.11</u>	0.405 <u>0.139</u>

Figures

Figure 1. Linear Measurements used in this study. SVL = snout-vent length, TL = trunk length, HL = head length, HW = head width, HH = head height, Lhu = humerus length, Lun = ulna length, Lfe = femur length, Ltb = tibia length (for details see Tejero-Cicuéndez et al. 2021a).

Figure 2. Plot of regression scores and predicted lines representing the relationship between linear body measurements and size (SVL). Individuals are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta). Isometric trend represented by the dashed line.

Figure 3. Traitgrams showing the evolution of body size (SVL) through time based on the phylogenetic tree of *Pristurus*. Colors represent an evolutionary mapping of residuals from phylogenetic regressions describing the relationship of (A) head morphology versus body size, and (B) limb proportions versus body size (see text for descriptions). Species names are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta).

Figure 4. Patterns of static allometry for each species for head traits (upper panel) and limb traits (lower panel). Species are separated by their habitat groups and colored by the magnitude of their regression slope (purple: steeper slopes, yellow: shallower slopes).

Figure 5. Phylomorphospace of *Pristurus*, based on residuals from a phylogenetic regression of body measurements on size (SVL). Species means are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta). Large and small rock-dwelling and ground-dwelling are highlighted with darker colors to highlight their differentiation and relative positions in morphospace. Point size is proportional to mean species body size. 79% of the total variation is displayed in the first two PC axes (PC1 = 62.8%; PC2 = 16.3%).

Figure 6. Representative specimens (based on real specimens) from large and small *Pristurus* species, colored by habitat use: ground (beige) and rock (dark purple). Specimens are scaled to a common body size (SVL, gray rectangles) to emphasize the relative differences in limb and head proportions. Relatively slender-headed and short-limbed species shown on the left. Original scale shown as the gray bar.




Figure 1: Linear Measurements used in this study. SVL = snout-vent length, TL = trunk length, HL = head length, HW = head width, HH = head height, Lhu = humerus length, Lun = ulna length, Lfe = femur length, Lti = tibia length (for details see Tejero-Cicuéndez et al. 2021a).




Figure 2: Plot of regression scores and predicted lines representing the relationship between linear body measurements and size (SVL). Individuals are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta). Isometric trend represented by the dashed line.




Figure 3: Traitgrams showing the evolution of body size (SVL) through time based on the phylogenetic tree of *Pristurus*. Colors represent an evolutionary mapping of residuals from phylogenetic regressions describing the relationship of (A) head morphology versus body size, and (B) limb proportions versus body size (see text for descriptions). Species names are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta).





Figure 4: Patterns of static allometry for each species for head traits (upper panel) and limb traits (lower panel). Species are separated by their habitat groups and colored by the magnitude of their regression slope (purple: steeper slopes, yellow: shallower slopes).

`\begin{figure}`



Phylomorphospace of *Pristurus*, based on residuals from a phylogenetic regression of body measurements on size (SVL). Species means are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta). Large and small rock-dwelling and ground-dwelling are highlighted with darker colors to highlight their differentiation and relative positions in morphospace.

Phylomorphospace of *Pristurus*, based on residuals from a phylogenetic regression of body measurements on size (SVL). Species means are colored by habitat use: ground (beige), rock (dark purple), and tree (magenta). Large and small rock-dwelling and ground-dwelling are highlighted with darker colors to highlight their differentiation and relative positions in morphospace. Point size is proportional to mean species body size. 79% of the total variation is displayed in the first two PC axes (PC1 = 62.8%; PC2 = 16.3%).




Figure 5: Representative specimens (based on real specimens) from large and small *Pristurus* species, colored by habitat use: ground (beige) and rock (dark purple). Specimens are scaled to a common body size (SVL, gray rectangles) to emphasize the relative differences in limb and head proportions. Relatively slender-headed and short-limbed species shown on the left. Original scale shown as the gray bar.