

XXX

Héctor Tejero-Cicuéndez^{1,*}, Iris Menéndez^{2,3}, Salvador Carranza¹, and Dean C. Adams⁴

16 September, 2022

¹Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49, Barcelona 08002, Spain

²Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias Geológicas, Universidad Complutense de Madrid, C/José Antonio Novais 12, Madrid 28040, Spain

³Departamento de Cambio Medioambiental, Instituto de Geociencias (UCM, CSIC), C/Severo Ochoa 7, Madrid 28040, Spain

⁴Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa, 50010 USA

*Correspondence: Héctor Tejero-Cicuéndez cicuendez93@gmail.com

Keywords: Phenotypic Evolution, Morphospace, Allometry, *Pristurus* geckos

Short Title: XXX

Author Contributions: All authors collaboratively developed the concept and contributed to all portions of this manuscript. HT-C, IM, and DCA performed the analyses. All authors approve of the final product and are willingly accountable for any portion of the content.

Conflicts of Interests: The authors declare no conflicts of interest.

Data Archiving: Data are available on DRYAD ([doi:10.5061/dryad.xwdbrv1f6](https://doi.org/10.5061/dryad.xwdbrv1f6)) (Tejero-Cicuéndez et al.

26 2021b)). R-scripts are found in the Supplemental Information.

27

28 **Acknowledgments:** We thank XYZPDQ... This work was sponsored in part by XXX (to SC) DCA was
29 funded in part by National Science Foundation Grant DEB-2140720, and a Fulbright Senior Scholar Grant.

³⁰ **Abstract**

³¹ asdf

Introduction

some general paragraph on the evolution of phenotypic diversity

when organisms colonize new and unique habitats, they are subjected to novel ecological selection pressures in those habitats. Often these selective pressures elicit changes in body form, as organisms adapt to their new habitats (examples: some comment on ecomorphs, etc.). . . . leads to so-called ecomorphs, with such well known examples in Anolis lizards, cichlid fishes, etc. It follows that . . . Some comment on the fact that clades living in diverse ecological conditions often display greater diversity in form and function (REFS).

However, while the above patterns have been well documented in a variety of vertebrate taxa, what remains less known is how allometry plays a role in this phenotypic diversification. We know that XYZPDQ (about allometry). Then links to diversity..

The Afro-Arabian geckos in the genus *Pristurus* afford the opportunity to elucidate the interdigitating effects of allometry and habitat specialization on clade-level patterns of phenotypic diversity. Prior work on this system (Tejero-Cicuéndez et al. 2021a) has revealed that . . . (sentence or 2 about your prior study, getting to diversity and . . . Importantly, . . . something about habitat. . . . What remains unexamined however, is XYZPDQ. . .

In this study, we . . .

Materials and Methods

Data

Phylogenetic, ecological, phenotypic. . . . (describe briefly). Data from (Tejero-Cicuéndez et al. 2021a).

Statistical Analyses

To test the hypothesis. . .

- Mancova $\text{body} \sim \text{SVL} * \text{hab.gp}$
 - PW of slopes, and inspected reg. coefficients to identify biological trends
 - Visualized multivariate regressions via regression scores (sensu Drake and Klingenberg 2008) and predicted lines (sensu Adams and Nistri 2010)
- Examine allometry phylogenetically.

- PLS of head vs. SVL and limb vs SVL. Obtained scores on 1st axis for each.
- within-species regressions of Head.sc \sim SVL & limb.sc \sim SVL; obtained regression coefficients (slopes)
- mapped slopes on phylogeny under BM and generated traitgrams to identify changes in allometric relationships across the phylogeny
- Finally, to link allometric patterns with trends in phenotypic diversification we obtained size-standardized species means, following procedures in H TC paper (residuals from phylo-regressions of traits on SVL, residuals). We then performed an ordination to obtain a phylomorphospace, where habitat types and species could be observed.

Results

Discussion

References

- Adams, D. C., and A. Nistri. 2010. Ontogenetic convergence and evolution of foot morphology in european cave salamanders (family: plethodontidae). *BMC Evolutionary Biology* 10:1–10. BioMed Central.
- Drake, A. G., and C. P. Klingenberg. 2008. The pace of morphological change: Historical transformation of skull shape in st bernard dogs. *Proceedings of the Royal Society B: Biological Sciences* 275:71–76.
- Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021a. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. *Proceedings of the Royal Society B: Biological Sciences* 288:20211821.
- Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021b. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. Dryad digital repository. (Doi:10.5061/dryad.xwdbrv1f6).

Figures

Figure 1. Linear Measures

Figure 2. Plot of regression scores and predicted lines representing the relationship between linear body measurements and size (SVL). Individuals occupying differing habitats are denoted by distinct colors as: rock (beige), ground (dark purple), and tree (magenta).

Figure 3. Traitgrams showing the evolution of body size (SVL) through time mapped on the phylogenetic tree of *Pristurus*. Colors represent evolutionary mapping of colored by (A) mapped by the discrete categories of presence in Socotra or the continent (left) and by ecological specialization (right).

Figure 4. Phylomorphospace




Figure 1: asdf.




Figure 2: Plot of regression scores and predicted lines representing the relationship body proportions and body size (SVL). Individuals occupying differing habitats are denoted by distinct colors as: rock (beige), ground (dark purple), and tree (magenta).




Figure 3: xxx.




Figure 4: asdf.