

Title: Evolution along allometric lines of least resistance: Morphological differentiation in *Pristurus* geckos

Running title: Allometric patterns in *Pristurus*

Héctor Tejero-Cicuéndez^{1,2,*}, Iris Menéndez³, Adrián Talavera², Gabriel Riaño², Bernat Burriel-Carranza², Marc Simó-Riudalbas², Salvador Carranza², Dean C. Adams⁴

¹Evolution and Conservation Biology research Group, Department of Biodiversity, Ecology and Evolution. Faculty of Biology. Universidad Complutense de Madrid, 28040, Madrid, Spain

²Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49, Barcelona 08003, Spain

³Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

⁴Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa, 50010 USA

*Correspondence: Héctor Tejero-Cicuéndez cicuendez93@gmail.com; C/ José Antonio Novais, 12, 28040 Madrid, Spain.

Author contributions. Conceptualization: HT-C, IM, and DCA. Supervision: DCA. Project Administration: HT-C and DCA. Investigation: HT-C, IM, and DCA. Formal Analysis: HT-C, IM, and DCA. Software: HT-C, IM, and DCA. Methodology: HT-C, IM, and DCA. Validation: HT-C, IM, AT, GR, BB-C, MS-R, SC, and DCA. Data Curation: HT-C, IM, MS-R, SC, and DCA. Resources: HT-C, IM, AT, SC, and DCA. Funding Acquisition: HT-C, IM, AT, GR, BB-C, SC, and DCA. Writing - Original Draft Preparation: HT-C, IM, and DCA. Writing - Review & Editing: HT-C, IM, AT, GR, BB-C, MS-R, SC, and DCA. Visualization: HT-C, IM, and AT.

Acknowledgments: We are very grateful to J. Roca, M. Metallinou, K. Tamar, J. Šmíd, R. Vasconcelos, R. Sindaco, F. Amat, Ph. de Pous, L. Machado, J. Garcia-Porta, J. Els, T. Mazuch, T. Papenfuss, and all the people from the Environment Authority, Oman, for their help in different aspects of the work.

Funding Statement: This work was funded in part by PGC2018-098290-B-I00 (MCIU/AEI/FEDER, UE) and PID2021-128901NB-I00 (MCIN/AEI/10.13039/501100011033 and by ERDF, A way of making Europe), Spain to SC. HT-C is supported by a “Juan de la Cierva - Formación” postdoctoral fellowship (FJC2021-046832-I). IM was funded by the Alexander von Humboldt Foundation through a Humboldt Research Fellowship. AT is supported by the “la Caixa” doctoral fellowship programme (LCF/BQ/DR20/11790007). GR was funded by an FPI grant from the Ministerio de Ciencia, Innovación y Universidades, Spain (PRE2019-088729). BB-C was funded by FPU grant from Ministerio de Ciencia, Innovación y Universidades, Spain (FPU18/04742). DCA was funded in part by National Science Foundation Grant DBI-1902511.

Data availability statement: All the data used in this study are available on DRYAD from a previous study: <https://doi.org/10.5061/dryad.xwdbvr1f6> (Tejero-Cicuéndez et al. 2021b). The scripts for implementing all analyses and generating the figures in this manuscript can be found in the Supplementary Material and in a GitHub repository (and on DRYAD upon acceptance: <https://doi.org/10.5061/dryad.w0vt4b8w1>).

Competing interests: The authors declare no competing interests.