

1 The role of habitat and evolutionary allometry in the morphological
2 differentiation of *Pristurus* geckos

3
4 **ORDER TBD: Héctor Tejero-Cicuéndez^{1,*}, Iris Menéndez^{2,3}, Marc Simó Riudalbas¹,**
5 **Salvador Carranza¹, and Dean C. Adams⁴**

6 29 September, 2022

7 ¹Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la
8 Barceloneta 37-49, Barcelona 08002, Spain

9 ²Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias Geológicas,
10 Universidad Complutense de Madrid, C/José Antonio Novais 12, Madrid 28040, Spain

11 ³Departamento de Cambio Medioambiental, Instituto de Geociencias (UCM, CSIC), C/Severo
12 Ochoa 7, Madrid 28040, Spain

13 ⁴Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa,
14 50010 USA

15 *Correspondence: Héctor Tejero-Cicuéndez cicuendez93@gmail.com

16
17
18 **Keywords:** Phenotypic Evolution, Morphospace, Allometry, *Pristurus* geckos

19
20 **Short Title:** XXX

21
22 **Author Contributions:** All authors collaboratively developed the concept and contributed to all

portions of this manuscript. HT-C, IM, and DCA performed the analyses. All authors approve of the final product and are willingly accountable for any portion of the content.

Conflicts of Interests: The authors declare no conflicts of interest.

Data Archiving: Data are available on DRYAD (doi:10.5061/dryad.xwdbrv1f6 (Tejero-Cicuéndez et al. 2021b)). R-scripts are available at **XXX**.

Acknowledgments: We thank XYZPDQ... This work was sponsored in part by XXX (to SC) DCA was funded in part by National Science Foundation Grant DBI-1902511.

33 **Abstract**

34 asdf

Introduction

Understanding how phenotypic diversity evolves, and elucidating the forces that generate and maintain this diversity, are major goals in evolutionary biology. Because adaptive evolution is the product of natural selection, changes in ecological selection pressures are expected to affect the evolutionary trajectory of phenotypic traits that facilitate an organism’s survival in their habitat. Evolutionary theory predicts that differing habitats will exert unique ecological selection pressures on organisms, resulting in associations between ecological and phenotypic traits. Indeed, species inhabiting differing habitats often display functional, behavioral, or phenotypic differences, that have presumably been the result of adaptive diversification in their respective ecological habitats (Collar et al. 2010; Kaliontzopoulou et al. 2015; Price et al. 2015; Martinez et al. 2021; Kolmann et al. 2022).

One possible evolutionary outcome of ecological specialization is that organisms inhabiting similar environments display common phenotypic characteristics. When such patterns occur repeatedly (e.g., Losos 1992; Schluter and McPhail 1992), this convergent evolution is treated as strong evidence of adaptation. Indeed the ecomorphological paradigm (sensu Arnold 1983) is predicated, in part, on such cases, which emphasize the strong association between the phenotypic traits that organisms display (morphological, behavioral, or physiological), and the ecological characteristics of their habitat that mediate organismal performance. In vertebrates, ecomorphological trends have been well-studied in numerous taxonomic groups, and include the emblematic ‘ecomorphs’ of Caribbean *Anolis* lizards that exploit different microhabitats (Losos 1992, 2009; Mahler et al. 2013), differential beak morphology in species of Darwin’s finches (Schluter and Grant 1984; Grant and Grant 2006; Reaney et al. 2020), the recurring phenotypes of African lake cichlids across ecological regimes (Albertson and Kocher 2001; Urban et al. 2022), and the distinct body forms of freshwater fishes in benthic and limnetic habitats (Jastrebski and Robinson 2004; Berner et al. 2008; Stuart et al. 2017) among others.

However, while the patterns of morphological differences in distinct ecological contexts have been well

documented, less-well understood is how this differentiation has been influenced by the covariance between body parts resulting from body size variation (i.e., allometry). It has long been recognized that the interrelationships among traits can have a strong influence on how phenotypic evolution proceeds, as trait correlations influence the degree to which phenotypic variation is exposed to selection (Wagner and Altenberg 1996). Thus, the integration among traits can constrain phenotypic change in certain directions, or enhance variation along other phenotypic axis (Schluter 1996; Wagner and Altenberg 1996; Wagner and Zhang 2011; Klingenberg and Marugán-Lobón 2013; Goswami et al. 2014, 2016; Felice et al. 2018). Further, because nearly all linear traits covary strongly with overall body size (Jolicœur 1963; Bookstein 2022), allometric trends could be considered the quintessential measure of phenotypic integration. Thus, identifying whether allometric patterns differ across habitats, and how such patterns of trait covariation affect ecomorphological trends among species utilizing those habitats, remains an important understudied question worthy of investigation.

The Afro-Arabian geckos in the genus *Pristurus* afford the opportunity to elucidate the interdigitating effects of allometry and habitat specialization on clade-level patterns of phenotypic diversity. Prior work on this system (Tejero-Cicuéndez et al. 2021a) revealed that the colonization of ground habitats has been a trigger of morphological change, specifically reflected in an increase in body size and shape disparity. Interestingly, some ground-dwelling species are among the largest of the genus and also show increased relative head sizes and limb proportions, while some other species with this ecological specialization have evolved to be among the smallest of the group. Additionally, among the species exploiting rocky habitats (the most common ecological feature in *Pristurus*), there are also species with both considerably large and small body sizes (Tejero-Cicuéndez et al. 2021a). What remains unexplored, however, is how the evolution of body shape is related to differences in body size and whether habitat specialization has an impact in this relationship shape-size. (how the size-shape relationship differs among habitats.)

In this study, we employed a combination of multivariate morphometric and phylogenetic comparative analysis to interrogate macroevolutionary patterns of evolutionary allometry in *Pristurus* geckos of Afro-Arabia. Using phenotypic, phylogenetic, and ecological data, we first characterized allometric

trends in body form in the group, to discern the extent to which allometric patterns differed across species occupying distinct ecological habitats. We then examined changes in allometric trends across the phylogeny, and linked these patterns to overall phenotypic diversification in morphospace, and in relation to habitat utilization. Overall our results demonstrate that differing allometric trajectories across habitats can result in similar overall phenotypes, even when those species display differing body sizes across habitat types **Reword: I don't like the last sentence... or have some more general sentence here....**

Materials and Methods

Data

We used a combination of phenotypic, phylogenetic, and ecological data to characterize and evaluate intra- and interspecific allometric trends. The data utilized here were obtained from our prior work on this system (Tejero-Cicuéndez et al. 2021a, 2022), and are briefly described here. First we used a time-dated, molecular phylogeny that included all members of the genus *Pristurus*, including several currently undescribed taxa. The tree was estimated in a Bayesian framework, using five mitochondrial markers, six nuclear markers, and 21 calibration points (for details see Tejero-Cicuéndez et al. 2022). Next we categorized each species as belonging to one of three ecological groups (ground, rock, or tree), based on descriptions of habitat use found in the literature (see Tejero-Cicuéndez et al. 2021a). Finally, we obtained a phenotypic data set containing body size (snout-vent length: SVL) and eight linear measurements (Figure 1) that described overall body form: trunk length (TrL), head length (HL), head width (HW), head height (HH), humerus length (Lhu), ulna length (Lun), femur length (Lfe), and tibia length (Ltb) (Tejero-Cicuéndez et al. 2021a). We restricted our study to those species represented by nine or more individuals; resulting in a dataset of 687 individuals from 25 species (individuals per species: $\mu = 27$; min = 9, max = 56). Species in the phenotypic dataset were then matched to the phylogeny, which was subsequently pruned to arrive at the final topology. All measurements were log-transformed prior to statistical analyses. Additional details regarding data collection and formal descriptions of each linear measurement may be found in the original sources (see Tejero-Cicuéndez et al. 2021a, 2022). The data are found on DRYAD: <https://doi.org/10.5061/dryad.xwdbrv1f6> (Tejero-Cicuéndez et al. 2021b).

Statistical and Comparative Analyses

We conducted a series of analyses to interrogate allometric trends and macroevolutionary changes in allometry, relative to differentiation in body form. First we characterized evolutionary allometry in the genus by performing a phylogenetic multivariate regression of body form on size, using the species means as data. We then performed an analogous procedure at the individual level, regressing body form on size using our entire dataset. From both the species-level (phylogenetic) and the individual-level regression models, we obtained the set of regression coefficients, and calculated the difference between them to describe the extent to which patterns of allometry at the individual level were concordant with evolutionary allometric trends across species.

Next we used the individual dataset to determine whether allometric trends in body form differed across habitat groups. This was accomplished by performing a multivariate analysis of covariance, with body size (*SVL*), *habitat*, and $SVL \times habitat$ as model effects. Significance was evaluated using 999 iterations of a permutation procedure, where residuals from a reduced model were randomly permuted in each permutation (RRPP), model statistics were recalculated, and used to generate empirical null sampling distributions to evaluate the observed test statistics (following Freedman and Lane 1983; Collyer and Adams 2007; Collyer et al. 2015). We then compared the multivariate allometric vectors for each habitat group by calculating pairwise differences in their angular direction in morphospace, and evaluating these relative to empirical sampling distributions obtained through RRPP (Collyer and Adams 2007; Adams and Collyer 2009; Collyer and Adams 2013). Patterns of multivariate allometry relative to body size were visualized via regression scores (Drake and Klingenberg 2008) and predicted lines (Adams and Nistri 2010), based on the coefficients and fitted values from the linear model described above.

We then examined changes in allometric trends across the phylogeny. Here we treated the head dimensions and limb dimensions separately, as allometric trends could potentially differ between these body regions due to differential functional or selective constraints (Kaliontzopoulou et al. 2010). Because both the head and limb data were multivariate, we first performed a partial least squares analysis (Rohlf and Corti 2000) of the head traits versus SVL, and the limb traits versus

SVL, to describe the direction of maximal covariation between each body region and size. PLS scores from each analysis were obtained, and species-specific slopes describing the extent of head and limb allometry within each species were extracted from an analysis of covariance modeled as: $PLS1_{head} \sim SVL * species$ and $PLS1_{limb} \sim SVL * species$ respectively. The species-specific allometric slopes were then mapped on the phylogeny of *Pristurus* using a Brownian motion model of evolution, to qualitatively evaluate shifts in allometry across the phylogeny for the group (for a similar approach see Adams and Nistri 2010).

Finally, to relate within-species allometric trends with patterns of phenotypic diversification in the group we generated a phylomorphospace, based on the size-standardized species means obtained from a phylogenetic regression (see Tejero-Cicuéndez et al. 2021a). Here, phenotypic similarities among species, relative to their phylogenetic relationships and habitat affiliations, were observed. All analyses were conducted in R 4.2.1 (R Core Team 2022), using RRPP version 1.3.1 (Collyer and Adams 2018; Collyer and Adams 2022), and scripts written by the authors (available at **XXX**).

Results

Using phylogenetic regression, we found significant evolutionary allometry in body form across species ($N_{sp} = 25$; $F = 217.9$; $Z = 5.53$; $P < 0.001$). Likewise, when allometry in body form was examined across individuals, a similar pattern was observed ($N_{ind} = 687$; $F = 7910.8$; $Z = 9.20$; $P < 0.001$). Further, the vectors of regression coefficients between the two analyses were highly correlated ($\rho = 0.94$) and were oriented in nearly parallel directions in morphospace ($\theta = 1.49^\circ$). This revealed that the pattern of multivariate allometry across individuals was concordant with macroevolutionary trends of interspecific allometry among species of *Pristurus* across the phylogeny.

Our analyses also exposed significant differences in the allometry of body form among *Pristurus* utilizing distinct habitats (Table 1). Here, comparisons of multivariate allometric vectors identified that ground-dwelling *Pristurus* displayed a distinct allometric trend as compared with *Pristurus* occupying both the rock and tree habitats (Table 2). Inspection of the regression coefficients for

each trait (Supplemental Information) found steeper allometric coefficients for all head and limb traits in ground-dwelling *Pristurus* as compared with rock and tree-dwelling taxa, corroborating this result. These findings implied that larger individuals of ground-dwelling *Pristurus* species displayed proportionately larger heads and limbs, as compared with large individuals in taxa utilizing other habitat types. A visualization of multivariate allometric trends (Fig. 2) confirmed these statistical findings, and indicated that the allometric trajectory in rock-dwelling animals was more extreme as compared with either ground or tree-dwelling *Pristurus*.

When allometric patterns were mapped on the phylogeny, traitgrams elucidated that changes in allometric trends were not concentrated to specific regions of the phylogeny (Figure 3). Rather, increases and decreases in allometry of both the head traits and the limb traits occurred repeatedly (see also Supplemental Information). When these patterns were viewed relative to body size, large ground-dwelling species tended to display steeper head allometry as compared to large rock-dwelling species; who displayed shallower head allometry (Figure 3A). Likewise, a similar pattern was observed when comparing small species utilizing these habitats. In contrast, large ground-dwelling species exhibited intermediate patterns of limb allometry, and small species in general were more variable in their patterns of limb allometry (Figure 3B). Inspection of traitgrams thus revealed some degree of discordance in patterns of allometry across body regions; a pattern evidenced by the relatively low correlation between slopes representing head and limb allometry respectively ($\rho = 0.42$).

When body shape differentiation in *Pristurus* was viewed in phylomorphospace (Figure 4), we found broad overlap among habitat groups, though arboreal (tree-dwelling) species were somewhat more separated in morphospace. Rock-dwelling species occupied a slightly larger region of morphospace as compared with the other groups, though this pattern was not statistically significant (Supplemental Information). Intriguingly, when viewed in relation to body size, large *Pristurus* species were not localized to a particular region of morphospace, nor were smaller species. Instead, the largest rock-dwelling species were found in close proximity to the smallest ground-dwelling species, indicating that they were similar in overall body shape. Likewise, the smaller rock-dwelling species were found

close to large ground-dwelling species in morphospace, indicating they displayed similar body shapes as well. Finally, integrating the patterns revealed in the phylomorphospace with those identified in our earlier analyses revealed a complex interplay between body shape, body size, habitat use, and multivariate allometry; where species with similar body shapes displayed differing overall size, were found in distinct habitats, and exhibited different allometric trends.

Discussion

- First paragraph; restate topic/questions (see last par of Intro); summarize general findings ...
 - something about linking allometry and phenotypic diversification?
- result 1: implication
- result 2: implication
- result 3: implication
 - morphospace: Thus there was a reciprocal relationship between body shape and body size across ground-dwelling and rock-dwelling species.
- one interesting... head vs. (correlation of head vs. limb slopes: 0.42. Pretty low. Implies some sort of differential something here, resulting in distinct allometric patterns for these two body regions. Similar to Antigoni's work (and refs therein). IMPLICATION: tie this into integration/modularity. Less integrated across the whole organism, and more modular... Future studies should examine this.

closing paragraph

References

- Adams, D. C., and M. L. Collyer. 2009. A general framework for the analysis of phenotypic trajectories in evolutionary studies. *Evolution* 63:1143–1154.
- Adams, D. C., and A. Nistri. 2010. Ontogenetic convergence and evolution of foot morphology in european cave salamanders (family: plethodontidae). *BMC Evolutionary Biology* 10:1–10. BioMed Central.
- Albertson, R. C., and T. D. Kocher. 2001. Assessing morphological differences in an adaptive trait: A landmark-based morphometric approach. *Journal of Experimental Zoology* 289:385–403.
- Arnold, S. J. 1983. Morphology, performance, fitness. *American Zoologist* 23:347–361.
- Berner, D., D. C. Adams, A.-C. Grandchamp, and A. P. Hendry. 2008. Natural selection drives patterns of lake-stream divergence in stickleback foraging morphology. *Journal of Evolutionary Biology* 21:1653–1665.
- Bookstein, F. L. 2022. Dimensions of morphological integration. *Evolutionary Biology* 49:342–372.
- Collar, D. C., J. A. Schulte, B. C. O’Meara, and J. B. Losos. 2010. Habitat use affects morphological diversification in dragon lizards. *Journal of Evolutionary Biology* 23:1033–1049.
- Collyer, M. L., and D. C. Adams. 2007. Analysis of two-state multivariate phenotypic change in ecological studies. *Ecology* 88:683–692.
- Collyer, M. L., and D. C. Adams. 2013. Phenotypic trajectory analysis: Comparison of shape change patterns in evolution and ecology. *Hystrix* 24:75–83.
- Collyer, M. L., and D. C. Adams. 2022. R: RRPP: Linear model evaluation with randomized residuals in a permutation procedure. Vsn. 1.3.1. R Foundation for Statistical Computing, Vienna, Austria.
- Collyer, M. L., and D. C. Adams. 2018. RRPP: An r package for fitting linear models to high-dimensional data using residual randomization. *Methods in Ecology and Evolution* 9:1772–1779.
- Collyer, M. L., D. J. Sekora, and D. C. Adams. 2015. A method for analysis of phenotypic

- change for phenotypes described by high-dimensional data. *Heredity* 115:357–365.
- Drake, A. G., and C. P. Klingenberg. 2008. The pace of morphological change: Historical transformation of skull shape in st bernard dogs. *Proceedings of the Royal Society B: Biological Sciences* 275:71–76.
- Felice, R. N., M. Randau, and A. Goswami. 2018. A fly in a tube: Macroevolutionary expectations for integrated phenotypes. *Evolution* 72:2580–2594.
- Freedman, D., and D. Lane. 1983. A nonstochastic interpretation of reported significance levels. *Journal of Business & Economic Statistics* 1:292–298.
- Goswami, A., M. Randau, P. D. Polly, V. Weisbecker, C. Verity Bennett, L. Hautier, and M. R. Sánchez-Villagra. 2016. Do developmental constraints and high integration limit the evolution of the marsupial oral apparatus? *Integrative and Comparative Biology* 56:404–415.
- Goswami, A., J. B. Smaers, C. Soligo, and P. D. Polly. 2014. The macroevolutionary consequences of phenotypic integration: From development to deep time. *Philosophical Transactions of the Royal Society B: Biological Sciences* 369:20130254.
- Grant, P. R., and B. R. Grant. 2006. Evolution of character displacement in darwin’s finches. *Science* 313:224–226.
- Jastrebski, C. J., and B. W. Robinson. 2004. Natural selection and the evolution of replicated trophic polymorphisms in pumpkinseed sunfish (*Lepomis gibbosus*). *Evolutionary Ecology Research* 6:285–305.
- Jolicoeur, P. 1963. The multivariate generalization of the allometry equation. *Biometrics* 19:497–499.
- Kaliontzopoulou, A., M. A. Carretero, and D. C. Adams. 2015. Ecomorphological variation in male and female wall lizards and the macroevolution of sexual dimorphism in relation to habitat use. *Journal of Evolutionary Biology* 28:80–94.
- Kaliontzopoulou, A., M. A. Carretero, and G. A. Llorente. 2010. Intraspecific ecomorphological variation: Linear and geometric morphometrics reveal habitat-related patterns within *Podarcis bocagei* wall lizards. *Journal of Evolutionary Biology* 23:1234–1244.
- Klingenberg, C. P., and J. Marugán-Lobón. 2013. Evolutionary covariation in geometric morphometric data: Analyzing integration, modularity, and allometry in a phylogenetic context. *Systematic Biology* 62:591–610.

- Kolmann, M. A., F. P. L. Marques, J. C. Weaver, M. N. Dean, J. P. Fontenelle, and N. R. Lovejoy. 2022. Ecological and phenotypic diversification after a continental invasion in neotropical freshwater stingrays. *Integrative and Comparative Biology* 62:424–440.
- Losos, J. B. 2009. *Lizards in an evolutionary tree: Ecology and adaptive radiation of anoles*. University of California Press.
- Losos, J. B. 1992. The evolution of convergent structure in Caribbean *Anolis* communities. *Systematic Biology* 41:403–420.
- Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the macroevolutionary landscape in island lizard radiations. *Science* 341:292–295.
- Martinez, C. M., S. T. Friedman, K. A. Corn, O. Larouche, S. A. Price, and P. C. Wainwright. 2021. The deep sea is a hot spot of fish body shape evolution. *Ecology Letters* 24:1788–1799.
- Price, S. A., S. T. Friedman, and P. C. Wainwright. 2015. How predation shaped fish: The impact of fin spines on body form evolution across teleosts. *Proceedings of the Royal Society B: Biological Sciences* 282:20151428.
- R Core Team. 2022. *R: A language and environment for statistical computing*. Version 4.2.1. R Foundation for Statistical Computing, Vienna, Austria.
- Reaney, A. M., Y. Bouchenak-Khelladi, J. A. Tobias, and A. Abzhanov. 2020. Ecological and morphological determinants of evolutionary diversification in darwin's finches and their relatives. *Ecology and Evolution* 10:14020–14032.
- Rohlf, F. J., and M. Corti. 2000. Use of two-block partial least-squares to study covariation in shape. *Systematic Biology* 49:740–753.
- Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. *Evolution* 50:1766–1774.
- Schluter, D., and P. R. Grant. 1984. Determinants of morphological patterns in communities of darwin's finches. *The American Naturalist* 123:175–196.
- Schluter, D., and J. D. McPhail. 1992. Ecological character displacement and speciation in sticklebacks. *The American Naturalist* 140:85–108.
- Stuart, Y. E., T. Veen, J. N. Weber, D. Hanson, M. Ravinet, B. K. Lohman, C. J. Thompson, T. Tasneem, A. Doggett, R. Izen, N. Ahmed, R. D. H. Barrett, A. P. Hendry, C. L. Peichel, and D. I. Bolnick. 2017. Contrasting effects of environment and genetics generate a continuum

of parallel evolution. *Nature Ecology and Evolution* 1:158.

Tejero-Cicuéndez, H., A. H. Patton, D. S. Caetano, J. Šmíd, L. J. Harmon, and S. Carranza. 2022. Reconstructing squamate biogeography in afro-arabia reveals the influence of a complex and dynamic geologic past. *Systematic Biology* 71:261–272.

Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021a. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. *Proceedings of the Royal Society B: Biological Sciences* 288:20211821.

Tejero-Cicuéndez, H., M. Simó-Riudalbas, I. Menéndez, and S. Carranza. 2021b. Ecological specialization, rather than the island effect, explains morphological diversification in an ancient radiation of geckos. Dryad digital repository. (Doi:10.5061/dryad.xwdbrv1f6).

Urban, S., J. Gerwin, C. D. Hulsey, A. Meyer, and C. F. Kratochwil. 2022. The repeated evolution of stripe patterns is correlated with body morphology in the adaptive radiations of east african cichlid fishes. *Ecology and Evolution* 12:e8568.

Wagner, G. P., and J. Zhang. 2011. The pleiotropic structure of the genotype–phenotype map: The evolvability of complex organisms. *Nature Reviews Genetics* 12:204–213.

Wagner, G., and L. Altenberg. 1996. Perspective: Complex adaptations and the evolution of evolvability. *Evolution* 50:967–976.

Table 1: Multivariate analysis of covariance describing variation in body form in extitPristurus.

	Df	SS	MS	Rsqr	F	Z	Pr(>F)
svl	1	516.036559	516.0365588	0.9203096	10188.69842	9.490057	0.001
habitat	2	6.218510	3.1092552	0.0110902	61.38957	9.322480	0.001
svl:habitat	2	3.974307	1.9871536	0.0070879	39.23464	7.077264	0.001
Residuals	681	34.491245	0.0506479	0.0615124			
Total	686	560.720622					

Table 2: Pairwise comparisons of multivariate allometry vectors. Effect sizes ($Z_{\theta_{12}}$) based on pairwise differences in angular direction are below the diagonal, and their corresponding significance levels are above diagonal. Significant values in bold.

	Ground	Rock	Tree
Ground	0	0.001	0.001
Rock	6.872	0	0.261
Tree	3.657	0.649	0

Figures

Figure 1. Linear Measurements used in this study. SVL = snout-vent length, TL = trunk length, HL = head length, HW = head width, HH = head height, Lhu = humerus length, Lun = ulna length, Lfe = femur length, Ltb = tibia length (for details see Tejero-Cicuéndez et al. 2021a).

Figure 2. Plot of regression scores and predicted lines representing the relationship between linear body measurements and size (SVL). Individuals are colored by habitat use: rock (beige), ground (dark purple), and tree (magenta).

Figure 3. Traitgrams showing the evolution of body size (SVL) through time based on the phylogenetic tree of *Pristurus*. Colors represent an evolutionary mapping of regression slopes describing the relationship of (A) head morphology versus body size, and (B) limb proportions versus body size (see text for descriptions). Species names are colored by habitat use: rock (beige), ground (dark purple), and tree (magenta).

Figure 4. Phylomorphospace of *Pristurus*, based on residuals from a phylogenetic regression of body measurements on size (SVL). Species means are colored by habitat use: rock (beige), ground (dark purple), and tree (magenta). Large and small rock-dwelling and ground-dwelling are highlighted with darker colors to highlight their differentiation and relative positions in morphospace.




Figure 1: Linear Measurements used in this study. SVL = snout-vent length, TL = trunk length, HL = head length, HW = head width, HH = head height, Lhu = humerus length, Lun = ulna length, Lfe = femur length, Ltb = tibia length (for details see Tejero-Cicu'endez et al. 2021a).




Figure 2: Plot of regression scores and predicted lines representing the relationship between linear body measurements and size (SVL). Individuals re colored by habitat use: rock (beige), ground (dark purple), and tree (magenta).




Figure 3: Traitgrams showing the evolution of body size (SVL) through time based on the phylogenetic tree of *Pristurus*. Colors represent an evolutionary mapping of regression slopes describing the relationship of (A) head morphology versus body size, and (B) limb proportions versus body size (see text for descriptions). Species names are colored by habitat use: rock (beige), ground (dark purple), and tree (magenta).




Figure 4: Phylomorphospace of *Pristurus*, based on residuals from a phylogenetic regression of body measurements on size (SVL). Species means are colored by habitat use: rock (beige), ground (dark purple), and tree (magenta). Large and small rock-dwelling and ground-dwelling are highlighted with darker colors to highlight their differentiation and relative positions in morphospace.