

Extreme Values in Financial Statistics

Master Thesis Project done at EPFL
towards the French 'Diplôme d'Ingénieur' degree
under the exchange agreement EPFL-ENSEEIH



Thèse de Master
présentée le 15 Août 2015
sous la supervision de
Prof. Stephan Morgenthaler,
Chaire de Statistique Appliquée
École Polytechnique Fédérale de Lausanne
pour l'obtention du Diplôme d'Ingénieur ENSEEIHT
en Mathématiques Appliquées et informatique
par

Killian Martin–Horgassan

Lausanne, EPFL, 2015

Mihi cura futuri
— Ovide, *Métamorphoses*, 13, 363

To my friends and loved ones...



Acknowledgements

TO BE FILLED

Lausanne, 14 Août 2015

K. M–H.



Preface

A preface is not mandatory. It would typically be written by some other person (eg your thesis director).

TO BE FILLED

Lausanne, 14 Août 2015

K. M–H.



Abstract

Key words:



Résumé

Mots clefs :

Contents

Acknowledgements	i
Preface	iii
Abstract (English/Français/Deutsch)	v
List of figures	xi
List of tables	xiii
1 Introduction	1
Introduction	1
1.1 A few words to set the scene	1
1.2 Formalising the settings	1
2 Investigating results on the limiting distribution	3
2.1 Playing with the (original) sequence of maxima	3
2.1.1 Sample following a Normal distribution	3
2.1.2 Sample following a Cauchy distribution	5
2.1.3 Sample following an Exponential Distribution	7
2.1.4 Why does this work ?	8
2.2 Making appear extreme value distributions	9
2.2.1 Making it appear in our context	9
3 Looking into real-world data	11
Looking into real-world data	11
3.1 Five real-world stocks and their evolution over 15 years	11
3.2 A détour around Brownian Motion	14
3.3 Back to the data	14
A An appendix	15
Bibliography	17

List of Figures

2.1	Below, a realisation of the sequence of maxima for i.i.d. standard unit Gaussian RVs	4
2.2	Scatter Plot of the Maxima, $n = 10000$	4
2.3	Maxima against the time steps	4
2.4	Maxima against the time steps and function $n \rightarrow \sqrt{2 * \log(n)}$	5
2.5	The correction becomes negligible compared to the starting term as n grows large	5
2.6	Below, a realisation of the sequence of maxima for i.i.d. <i>Cauchy</i> (0,1) RVs	6
2.7	Scatter Plot of the Maxima, $n = 10000$	6
2.8	Maxima against the time steps	6
2.9	Maxima against the time steps and function $n \rightarrow \tan(\pi * \frac{2-n}{2*n})$	7
2.10	Below, a realisation of the sequence of maxima for i.i.d. <i>Exp</i> (1) RVs	7
2.11	Scatter Plot of the Maxima, $n = 10000$	8
2.12	Maxima against the time steps	8
2.13	Maxima against the time steps and function $n \rightarrow \log(n)$	8
3.1	15 years of weekly BNP Stock Price Data	11
3.2	15 years of weekly Carrefour Stock Price Data	11
3.3	15 years of weekly LVMH Stock Price Data	12
3.4	15 years of weekly Sanofi Stock Price Data	12
3.5	15 years of weekly Total Stock Price Data	12
3.6	$\exp(R_t)$ and $1 + R_t$ for BNP Stock Price Data, residual : 3.22E-15	13
3.7	$\exp(R_t)$ and $1 + R_t$ for Carrefour Stock Price Data, residual : 1.44E-15	13
3.8	$\exp(R_t)$ and $1 + R_t$ for LVMH Stock Price Data, residual : 4.75E-15	14
3.9	$\exp(R_t)$ and $1 + R_t$ for Sanofi Stock Price Data, residual : 1.55E-15	14
3.10	$\exp(R_t)$ and $1 + R_t$ for Total Stock Price Data, residual : 3.86E-15	14



List of Tables

1 Introduction

1.1 A few words to set the scene

In real life, it is not uncommon to have at one's disposal data about a phenomenon occurring through time. It may be as simple as daily rainfall data in a city for the past two years, or it could be the weekly opening prices of a stock for the past decade.

Most of the time, people would like to use the data at their disposal to make predictions to answer questions, from the prosaic ones such as 'Will it rain tomorrow?' to more consequential ones such as 'Will I make a profit if I cling to my shares today and sell them only tomorrow?'. Of course, those are only vaguely worded questions : it is impossible to answer them satisfactorily without knowing the context, the objectives etc. behind them.

Yet, what these questions have in common is that they focus on the normal 'behaviour' that is to be expected in the future. Depending on the specific issue that is considered, the 'average behaviour' may not be the most interesting thing. For instance, suppose that a government wants to build a network of dams¹. The dams are meant to protect the country from future floods for the next one hundred years, therefore the question that needs to be answered is one of 'worst case event' : "Over the next century, how severe may be the worst flood?".

Extreme events are the kind of events we will be interested in this master thesis project. Although Extreme Value Theory has applications in many fields², we will here apply it more specifically to financial data.

1.2 Formalising the settings

Let $(X_n)_{n \geq 0}$ be a sequence of independent identically distributed random variables with common cumulative distribution function F_X . The sequence of maxima is defined by $M_0 = X_0$ and $\forall n \geq 1, M_n = \max_{0 \leq i \leq n} (X_i)$. We would like to determine the limiting distribution of the

¹As was done in The Netherlands beginning in the fifties

²including climate science, seismology, insurance etc.

Chapter 1. Introduction

sequence $(M_n)_{n \geq 0}$.³ This is a matter that will keep us busy quite a long time but the first thing to do is to re-formulate it.

Indeed, let us do a quick and simple computation :

$$\begin{aligned} F_n(t) &= \Pr(\{M_n \leq t\}) \\ &= \Pr(\{\max_{0 \leq i \leq n} (X_i) \leq t\}) \\ &= \Pr(\{X_1 \leq t\} \cap \cdots \cap \{X_n \leq t\}) \\ &= (F_X(t))^n \end{aligned} \tag{1.1}$$

Here we see that not much information will be drawn from this result by taking the limit $n \rightarrow +\infty$. The limiting distribution will be degenerate. Indeed, let us consider the upper end-point of F_X ⁴, z^+ . Then,

$$\begin{aligned} \forall z < z^+ \quad \lim_{n \rightarrow \infty} F_n(z) &= 0 \\ \forall z \geq z^+ \quad \lim_{n \rightarrow \infty} F_n(z) &= 1 \end{aligned} \tag{1.2}$$

It turns out we cannot use the limiting distribution directly. A common approach⁵ is to consider a sequence of the maxima, centred and normalised.

We will thus consider in all what follows the sequence defined by $(M_n^*)_{n \geq 0} = (\frac{M_n - b_n}{a_n})_{n \geq 0}$ where $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ are a sequence of real numbers and positive real numbers respectively. Finding a result on whether such a sequence admits a limiting distributions, and the conditions under which the result holds, will be one of our goals.

³If we can determine the limiting distribution of the maxima from the data, then we will have a means to make predictions on the occurrence of future extreme events.

⁴that is the smallest z such that $F_X(z)$ be equal to one. For the Normal distribution, z will be $+\infty$, by contrast for a continuous Uniform Distribution $U([a, b])$ it will be b .

⁵adopted by the mathematicians that laid the grounds for EVT

2 Investigating results on the limiting distribution

2.1 Playing with the (original) sequence of maxima

Here, we will generate finite size sequences ($N = 10000$) of independent identically distributed random variables following respectively :

- a standard Normal Distribution $\mathcal{N}(0, 1)$
- a Cauchy Distribution $Cauchy(0, 1)$
- an Exponential Distribution $Exp(1)$

We will compute the sequence of maxima, neither centred nor normalised, and draw the scatter plot as well as the plot of the maxima M_n as a function of the time steps n . We will also draw the $\frac{1}{n}$ -quantiles of the distributions (distributions of the sample, not of the maxima) as a function of the time steps n . This will lead us to make an interesting observation.

2.1.1 Sample following a Normal distribution

Computing the quantiles The Normal distribution is a particular case because, unlike in the cases of the Cauchy and the Exponential distribution, there is no explicit form to the cumulative distribution function. We will thus use a "well-known"¹ inequality, holding $\forall t > 0$:

$$\left(\frac{1}{t} - \frac{1}{t^3}\right) * \frac{\exp(-\frac{t^2}{2})}{\sqrt{2 * \pi}} < 1 - \Phi(t) < \frac{1}{t} * \frac{\exp(-\frac{t^2}{2})}{\sqrt{2 * \pi}} \quad (2.1)$$

¹Many textbooks mention it, though it is not necessarily what springs to the mind when thinking about the properties of Gaussian RVs.

Chapter 2. Investigating results on the limiting distribution

From there, it is easy to see that the following holds :

$$1 - \Phi(t) \sim_{t \rightarrow +\infty} \frac{1}{t} * \frac{\exp(-\frac{t^2}{2})}{\sqrt{2 * \pi}} \quad (2.2)$$

When n grows large, the $\frac{1}{n}$ -quantile grows very large so it is valid to replace $1 - \Phi(t)$ by its equivalent in the equation satisfied by the quantiles :

$$\begin{aligned} F_X(q_{\frac{1}{n}}) &= 1 - \frac{1}{n} \\ \Leftrightarrow \quad \frac{1}{q_{\frac{1}{n}}} * \frac{\exp(-\frac{q_{\frac{1}{n}}^2}{2})}{\sqrt{2 * \pi}} &= \frac{1}{n} \\ \Leftrightarrow \quad \log(q_{\frac{1}{n}}) + \log(\exp(-\frac{q_{\frac{1}{n}}^2}{2})) + \log(\sqrt{2 * \pi}) &= \log(n) \end{aligned} \quad (2.3)$$

This equation cannot be solved analytically, we will resolve it iteratively. The starting point is $\log(n) = \frac{t_0^2}{2}$, which gives us $t_0 = \sqrt{(2 * \log(n))}$. If we then run the Newton-Raphson algorithm, we see that the corrections to t_0 from the next iterations are small enough that we can keep t_0 as solution.².

Figure 2.1 – Below, a realisation of the sequence of maxima for i.i.d. standard unit Gaussian RVs




Figure 2.2 – Scatter Plot of the Maxima, $n = 10000$




Figure 2.3 – Maxima against the time steps

²See the fourth of the figures below

2.1. Playing with the (original) sequence of maxima




Figure 2.4 – Maxima against the time steps and function $n \rightarrow \sqrt{2 \cdot \log(n)}$




Figure 2.5 – The correction becomes negligible compared to the starting term as n grows large

2.1.2 Sample following a Cauchy distribution

Computing the quantiles Let X_1, \dots, X_n be i.i.d. RVs $\sim \text{Cauchy}(0, 1)$. The distribution function is $F_X(t) = \frac{1}{\pi} * \arctan(x) - \frac{1}{2}$. The $\frac{1}{n}$ -quantiles satisfy the equation :

$$\begin{aligned}
 & F_X(q_{\frac{1}{n}}) = 1 - \frac{1}{n} \\
 \Leftrightarrow & \frac{\arctan(q_{\frac{1}{n}})}{\pi} + \frac{1}{2} = 1 - \frac{1}{n} \\
 \Leftrightarrow & \frac{\arctan(q_{\frac{1}{n}})}{\pi} = \frac{2-n}{n} \\
 \Leftrightarrow & q_{\frac{1}{n}} = \tan\left(\frac{\pi}{2} * \frac{2-n}{n}\right)
 \end{aligned} \tag{2.4}$$

Figure 2.6 – Below, a realisation of the sequence of maxima for i.i.d. *Cauchy(0,1)* RVs




Figure 2.7 – Scatter Plot of the Maxima, $n = 10000$




Figure 2.8 – Maxima against the time steps

2.1. Playing with the (original) sequence of maxima




Figure 2.9 – Maxima against the time steps and function $n \rightarrow \tan(\pi i * \frac{2-n}{2*n})$

2.1.3 Sample following an Exponential Distribution

Computing the quantiles Let X_1, \dots, X_n be i.i.d. RVs $\sim \text{Exponential}(\lambda)$. The distribution function is $F_X(t) = 1 - \exp(-\lambda * t)$. The $\frac{1}{n}$ -quantiles satisfy the equation :

$$\begin{aligned}
 F_X(q_{\frac{1}{n}}) &= 1 - \frac{1}{n} \\
 \Leftrightarrow 1 - \exp(-\lambda * q_{\frac{1}{n}}) &= 1 - \frac{1}{n} \\
 \Leftrightarrow q_{\frac{1}{n}} &= \frac{1}{\lambda} * \log(n)
 \end{aligned} \tag{2.5}$$

Figure 2.10 – Below, a realisation of the sequence of maxima for i.i.d. $\text{Exp}(1)$ RVs



Figure 2.11 – Scatter Plot of the Maxima, $n = 10000$
 Figure 2.12 – Maxima against the time steps




Figure 2.13 – Maxima against the time steps and function $n \rightarrow \log(n)$

2.1.4 Why does this work ?

(The underlying idea) Why have we made a link between the $\frac{1}{n}$ -quantiles of the common distribution of the X_i and the sequence of the M_n ? Actually, the $\frac{1}{n}$ -quantiles are a good proxy for the M_n .

The idea behind this is as follows, that the maxima will get closer to the upper end-point of the distribution of the X_i , F_{X_i} . That is also the case for the $\frac{1}{n}$ -quantiles of F_{X_i} . Of course, it is only an intuition ! In the next section, we will show that it turns out to be valid (and in what precise sense it does so).

A more rigorous approach

2.2 Making appear extreme value distributions

The Extreme Value Theorem The following result is fundamental in the field of Extreme Value Theory³, it will be found in any textbook under the name of Fisher-Tippett theorem or Extreme Value theorem.

Let $(X_n)_{n \geq 1}$ be a sequence of independent identically distributed random variables and $(M_n)_{n \geq 1}$ be the sequence of the maxima as previously defined. If there exist a sequence of nonnegative real numbers $(a_n)_{n \geq 1}$ and a sequence of real numbers $(b_n)_{n \geq 1}$ such that $(\frac{M_n - b_n}{a_n})_{n \geq 1}$ converges in distribution to a non-degenerate distribution function F , then F belongs to the Extreme Value distribution family.

Extreme Value distribution The Extreme Value distribution sub-divides into three families of distribution :

- the Fréchet distribution family, **Fréchet(a,b,α)**
 $G(z) = \exp(-(\frac{z-b}{a})^{-\alpha}) \mathbb{1}(\{z > b\})$, $a > 0$, $b \in \mathbb{R}$, $\alpha > 0$.
- the Gumbel distribution family, **Gumbel(a,b)**
 $G(z) = \exp(-\exp(-(\frac{z-b}{a})))$, $-\infty < z < +\infty$, $a > 0$, $b \in \mathbb{R}$.
- the Weibull distribution family, **Weibull(a,b,α)**
 $G(z) = \exp(-(\frac{z-b}{a})^{-\alpha}) \mathbb{1}(\{z > b\}) + \mathbb{1}(\{z \leq b\})$, $a > 0$, $b \in \mathbb{R}$, $\alpha > 0$.

Those three families can be combined into a single family of models, the Generalized Extreme Value Distribution.

$$G(z) = \exp(-(1 + \xi(\frac{z-\mu}{\sigma}))^{-\frac{1}{\xi}})^4$$

2.2.1 Making it appear in our context

Fréchet distribution Let X_1, \dots, X_n be i.i.d. RVs $\sim \text{Exponential}(\lambda)$. Let us remember from the previous section the form of the $\frac{1}{n}$ -quantile $q_n = \frac{1}{\lambda} \log(n)$. Let us evaluate F_{M_n} in $q_{\frac{1}{n}} - t$:

³it is the analogue in terms of extremes of the Central Limit Theorem in classical probability theory.

⁴However, the domain of the distribution depends on the parameters $\mu, \sigma, \xi : \{z / 1 + \xi \frac{z-\mu}{\sigma} > 0\}$, with μ, ξ in \mathbb{R} and $\sigma > 0$

$$\begin{aligned} F_{M_n}(q_{\frac{1}{n}} - t) &= (1 - \exp(-\lambda(q_{\frac{1}{n}} - t)))^n \\ &= (1 - \exp(\lambda t) \exp(-\lambda q_{\frac{1}{n}}))^n \\ &= (1 - \frac{\exp(\lambda t)}{n})^n \\ &\longrightarrow_{n \rightarrow +\infty} \exp(-\exp(\lambda t)) \end{aligned} \tag{2.6}$$

Gumbel distribution

Weibull distribution

3 Looking into real-world data

3.1 Five real-world stocks and their evolution over 15 years

We have chosen to study five stocks listed on the Paris Stock Exchange : BNP Paribas, Carrefour, LVMH, Sanofi and Total stocks.¹ The evolution of the stock prices has been studied over the past 15 years, on a weekly basis. We first draw the data itself, then the net returns and the gross log returns on the stocks²




Figure 3.1 – 15 years of weekly BNP Stock Price Data

Figure 3.2 – 15 years of weekly Carrefour Stock Price Data

¹We have chosen companies positioned on different domains, otherwise, information from different stock might more easily be redundant.

²Both quantities are widely used in Finance.

Chapter 3. Looking into real-world data




Figure 3.3 – 15 years of weekly LVMH Stock Price Data Figure 3.4 – 15 years of weekly Sanofi Stock Price Data




Figure 3.5 – 15 years of weekly Total Stock Price Data

Let X_t be the price of a stock at time t , the gross return at time $t + 1$ is defined as the ratio $\frac{X_{t+1}}{X_t}$, the net return at time $t + 1$ is defined as the ratio $r_t = \frac{X_{t+1} - X_t}{X_t}$ and the log gross return at time $t + 1$ is defined as the log of the gross return at time $t + 1$ i.e. $R_t = \log(\frac{X_{t+1}}{X_t})$. The latter two quantities are of particular interest in Finance.

Let us observe that the relationship between R_t , X_t and X_{t+1} can be rewritten as $X_{t+1} = \exp(R_{t+1}) * X_t$. An approximation would be to take $X_{t+1} = (1 + R_{t+1}) * X_t$ by taking the expansion of the exponential, cut at order 1. Below are the plots of the quantities $\exp(R_t)$ and $1 + R_t$ for the five stocks previously considered. As we can see from the value of the residuals, this is in practice a very good approximation !

3.1. Five real-world stocks and their evolution over 15 years




Figure 3.6 – $\exp(R_t)$ and $1 + R_t$ for BNP Stock Price Data, residual : $3.22E-15$
Figure 3.7 – $\exp(R_t)$ and $1 + R_t$ for Carrefour Stock Price Data, residual : $1.44E-15$




Figure 3.8 – $\exp(R_t)$ and $1 + R_t$ for LVMH Stock Price Data, residual : $4.75\text{E-}15$

Figure 3.9 – $\exp(R_t)$ and $1 + R_t$ for Sanofi Stock Price Data, residual : $1.55\text{E-}15$




Figure 3.10 – $\exp(R_t)$ and $1 + R_t$ for Total Stock Price Data, residual : $3.86\text{E-}15$

3.2 A détour around Brownian Motion

3.3 Back to the data

A An appendix

Bibliography

[Applied Technology Council(1985)] Applied Technology Council. Earthquake damage evaluation data for California. Technical report, Seismic Safety Commission, Applied Technology Council (ATC), California, 1985.

- Modelling Extremal Events, Embrechts, Kluppelberg and Mikosch
- Statistics of Extremes, Theory and Application, Beirlant, Goegebeur, Segers and Teugels
- An Introduction to Statistical Modeling of Extreme Values, Coles
- Introduction to Scientific Programming and Simulation Using R, Jones, Maillardet and Robinson