

## MA1200 TAKE HOME PROBLEM SET 2

The following is the second take-home assignment of MA1200, which counts 3 points of total 100 of your final score of the course.

Please submit it via canvas in a pdf file (you can handwrite the answers and take photos by your phone, then make it into a pdf file, see for example, <https://www.wikihow.com/Convert-JPG-to-PDF>) for how to combine jpg files to a pdf; you can also do it by note-taking apps on an iPad or an Surface)

Q1.  $f(x) = 3x^2 + 18x + 8$ , find the range, and sketch the graph.

Q2. Factorize  $3x^3 + 4x^2 - 17x - 6$  into product of three linear factors (hint, try  $x = 2$ )

Q3. Express  $\frac{x^2 + 11x + 20}{(x - 1)(x + 3)^2}$  into partial fractions.

Q4. Rewrite  $3 \cos(x) - 4 \sin(x)$  as  $r \cos(x + \alpha)$ , where  $r > 0$  and  $\alpha \in (0, \pi/2)$ . (hint,  $\cos(a + b) = \cos(a) \cos(b) - \sin(a) \sin(b)$ )

Q5. Solve  $3 \cos(x) - 4 \sin(x) = 5/2$

Q6. Solve  $\sin(3\theta) = \cos(2\theta)$


The assignment is due on 23:59 of Oct 23, Friday.

You will lose 1 point for each day of late submission. All submissions after the midnight of Oct 26 will be marked as 0.

MA 200. HW2. 0.5 point each.

Q1  $f(x) = 3x^2 + 18x + 8 = 3(x^2 + 6x + 9) + 8 - 27$   
 $= 3(x+3)^2 - 19.$

Range  $[-19, \infty)$



Q2.  $f(x) = 3x^3 + 4x^2 - 17x - 6.$

$$f(x) = 0 \Rightarrow 3x^3 + 4x^2 - 17x - 6 = (x-2)(3x^2 + 10x + 3)$$
$$= (x-2)(x+3)(3x+1)$$

Q3.  $\frac{x^2 + 11x + 20}{(x-1)(x+3)^2} = \frac{A}{x-1} + \frac{B}{x+3} + \frac{C}{(x+3)^2}$

$$x^2 + 11x + 20 = A(x+3)^2 + B(x-1)(x+3) + C(x-1)$$

$$x=1 \Rightarrow 1+11+20 = 16A \Rightarrow A=2$$

$$x=-3 \Rightarrow 9-33+20 = -4C \Rightarrow C=1$$

$$x=0 \Rightarrow 20 = 9A - 3B - C = 18 - 1 - 3B$$

$$B=-1$$

$$\Rightarrow \frac{2}{x-1} - \frac{1}{x+3} + \frac{1}{(x+3)^2}$$

Q4.  $3 \cos x - 4 \sin x = 5 \left( \frac{3}{5} \cos x - \frac{4}{5} \sin x \right)$

let  $A = \cos^{-1}\left(\frac{3}{5}\right)$ . then  $\cos A = \frac{3}{5}$ ,  $\sin A = \frac{4}{5}$

$$\Rightarrow 3 \cos x - 4 \sin x = 5 (\cos A \cos x - \sin A \sin x)$$

$$= 5 \cos (x+A) = 5 \cos \left(x + \cos^{-1}\left(\frac{3}{5}\right)\right)$$

$$\text{or } = 5 \cos \left(x + \sin^{-1}\left(\frac{4}{5}\right)\right)$$

Q5.  $3 \cos x - 4 \sin x = \frac{5}{2}$

$$\Rightarrow 5 \cos \left(x + \cos^{-1}\left(\frac{3}{5}\right)\right) = \frac{5}{2}$$

$$\Rightarrow \cos \left(x + \cos^{-1}\left(\frac{3}{5}\right)\right) = \frac{1}{2} = \cos\left(\frac{\pi}{3}\right)$$

$$x + \cos^{-1}\left(\frac{3}{5}\right) = 2n\pi \pm \frac{\pi}{3}$$

$$x = 2n\pi - \underbrace{\cos^{-1}\left(\frac{3}{5}\right)}_{\sin^{-1}\left(\frac{4}{5}\right)} \pm \frac{\pi}{3}, \quad n \in \mathbb{Z}$$

Q6  $\sin(3\theta) = \cos(2\theta)$

method 1

$$\cos\left(\frac{\pi}{2} - 3\theta\right) = \cos(2\theta)$$

$$\frac{\pi}{2} - 3\theta = 2n\pi \pm 2\theta$$

" + "

$$5\theta = -2n\pi + \frac{\pi}{2}$$

$$\theta = \frac{2}{5}n\pi + \frac{\pi}{10}$$

" - "

$$\theta = 2n\pi + \frac{\pi}{2}$$

method 2.

$$\sin(3\theta) = \sin\left(\frac{\pi}{2} - 2\theta\right)$$

$$\text{or } 3\theta = 2n\pi + \frac{\pi}{2} - 2\theta$$

$$3\theta = 2n\pi + \pi - \left(\frac{\pi}{2} - 2\theta\right)$$

$$\Rightarrow \theta = \frac{2}{5}n\pi + \frac{\pi}{10}$$

or

$$\theta = 2n\pi + \frac{\pi}{2}$$

$n \in \mathbb{Z}$