

Chapter 3. Multiple Integral

1 Single-Variable Case:

- Definition and Interpretation:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} [f(x_1^*) \Delta x + f(x_2^*) \Delta x + \cdots + f(x_n^*) \Delta x] = \lim_{n \rightarrow \infty} \sum_{j=1}^n \underbrace{f(x_j^*)}_{\text{height}} \underbrace{\Delta x_j}_{\text{base}} = \text{Area of the region between the curve } f \text{ and } x\text{-axis.}$$

- Computation:

$$\int_a^b f(x) dx = F(b) - F(a),$$

where F is an antiderivative of f , i.e. $F'(x) = \underline{f(x)}$.


eg. $F(x) = \frac{x^3}{3}$ is an antiderivative of x^2

$$\int_1^2 x^2 dx = \left(\frac{x^3}{3} \right) \Big|_{x=1}^{x=2} - \left(\frac{x^3}{3} \right) \Big|_{x=1} = \frac{2^3}{3} - \frac{1}{3} = \frac{7}{3}$$

2 Two-Variable Case (Double Integral):

$$z = f(x, y).$$

2.1 Definition and Interpretation:





$$V_{ij} = \underbrace{f(x_i^*, y_j^*)}_{\text{height.}} \cdot \underbrace{\Delta x_i \Delta y_j}_{\text{base}}$$



$$\iint_R f(x, y) dx dy = \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \sum_{i=1}^n \sum_{j=1}^m \underbrace{f(x_i^*, y_j^*) \Delta x_i \Delta y_j}$$

R ↑


double integral
of f on region R .

= Volume of the solid below
the surface of f and above xy -plane.






(a) $m = n = 4$, $V \approx 41.5$



(b) $m = n = 8$, $V \approx 44.875$



(c) $m = n = 16$, $V \approx 46.46875$

2.2 Computation of Double Integrals: How to evaluate $\iint_R f(x,y) dx dy$

Case 1: R is an rectangle: $R = \{ (x,y) : a \leq x \leq b, c \leq y \leq d \}$

$$\begin{aligned} \iint_R f(x,y) dx dy &= \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \sum_{i=1}^n \sum_{j=1}^m f(x_i^*, y_j^*) (\Delta x_i) (\Delta y_j) \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\lim_{m \rightarrow \infty} \sum_{j=1}^m f(x_i^*, y_j^*) \Delta y_j \right] \Delta x_i \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\int_c^d f(x_i^*, y) dy \right] \Delta x_i \\ &= \int_a^b \left[\int_c^d f(x, y) dy \right] dx \quad \leftarrow \text{iterated integral in } dy \text{ then } dx \end{aligned}$$

Similarly,


$$\iint_R f(x,y) dx dy = \lim_{n \rightarrow \infty} \sum_{j=1}^m \left[\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*, y_j^*) \Delta x_i \right] \Delta y_j = \int_c^d \int_a^b f(x, y) dx dy$$

iterated integral in dx then dy

Example Use iterated integral in two different orders to evaluate

$$\iint_R (2xy + y^2) dx dy$$

with $R = [1, 2] \times [0, 1]$.



$$\begin{aligned} \iint_R (2xy + y^2) dx dy &= \int_0^1 \int_1^2 (2xy + y^2) dx dy \\ &\stackrel{\text{treat } y \text{ as constant}}{=} \int_0^1 \left[x^2 y + x y^2 \right]_{x=1}^{x=2} dy \end{aligned}$$

$$\begin{aligned} &= \int_0^1 [2^2 y + 2y^2 - y - y^2] dy \\ &= \int_0^1 (3y^2 + y^2) dy = \left(\frac{3y^3}{3} + \frac{y^3}{3} \right) \Big|_{y=0}^{y=1} \end{aligned}$$

$$= \frac{3}{2} + \frac{1}{3} = \frac{11}{6}$$

$$\iint_R (2xy + y^2) \, dx \, dy = \int_1^2 \left[\int_0^1 \underline{(2xy + y^2)} \, dy \right] dx$$

$$\begin{array}{l} \text{treat } x \\ \text{as constant} \end{array} = \int_1^2 \left[\left(xy^2 + \frac{y^3}{3} \right) \Big|_{y=0}^{y=1} \right] dx$$

$$= \int_1^2 \left(x + \frac{1}{3} - 0 \right) dx$$


$$= \left(\frac{x^2}{2} + \frac{1}{3}x \right) \Big|_{x=1}^{x=2}$$

$$= \left(\frac{2^2}{2} + \frac{2}{3} \right) - \left(\frac{1}{2} + \frac{1}{3} \right)$$

$$= \frac{3}{2} + \frac{1}{3} = \frac{11}{6}$$

Case 2. R is vertically simple (vertical segments are easily bounded in x)


$$R = \{ (x, y) : a \leq x \leq b, g_2(x) \leq y \leq g_1(x) \}$$



$$\iint_R f(x, y) dx dy = \int_a^b \left[\int_{g_2(x)}^{g_1(x)} f(x, y) dy \right] dx$$

Example Evaluate $\iint_R 2x^2 y dx dy$, with R as a region bounded by $x = 1$, $x = 3$, $y = x^2$ and $y = -x^2 + 18$.

$$R = \{ (x, y) : 1 \leq x \leq 3, x^2 \leq y \leq 18 - x^2 \}$$




$$\begin{aligned} \iint_R 2x^2 y dx dy &= \int_1^3 \left[\int_{x^2}^{18-x^2} 2x^2 y dy \right] dx \\ &= \int_1^3 \left[x^2 y^2 \Big|_{y=x^2}^{y=18-x^2} \right] dx \\ &= \int_1^3 \left[x^2 (18-x^2)^2 - x^2 \cdot x^2 \right] dx \\ &= \dots = \frac{5328}{2} \end{aligned}$$

$$\begin{aligned} \iint_R 2x^2 y dx dy &= \int_1^3 \left[\int_0^9 2x^2 y dx \right] dy \\ &= \int_1^9 \left[\int_1^{\sqrt{y}} 2x^2 y dx \right] dy \\ &+ \int_9^{17} \left[\int_1^{\sqrt{18-y}} 2x^2 y dx \right] dy \\ &= \dots = \end{aligned}$$


Exercise Evaluate $\iint_R xy^2 dx dy$, with R as a region bounded by $x = 1$, $y = 0$ and $y = x^2$.

$$\iint_R xy^2 dx dy = \int_0^1 \int_{y=0}^{y=x^2} xy^2 dy dx$$



$$= \int_0^1 \int_{x=\sqrt{y}}^{\sqrt{1-y}} x y^2 \, \underline{dx} \, dy$$

Case 3. R is horizontally simple (horizontal segments are easily bounded in y)




$$R = \{(x, y) : c \leq y \leq d, x_1(y) \leq x \leq x_2(y)\}$$

$$\iint_R f(x, y) dx dy = \int_c^d \left[\int_{x_1(y)}^{x_2(y)} f(x, y) dx \right] dy$$

Example Evaluate

$$\iint_R xy^2 dx dy$$



, with R as a region bounded by $x = y^2$ and $x = -y^2 + 1$.

$$R = \{(x, y) : -\frac{1}{\sqrt{2}} \leq y \leq \frac{1}{\sqrt{2}}, y^2 \leq x \leq 1 - y^2\} \Rightarrow y^2 = -y^2 + 1 \Rightarrow y = \pm \frac{1}{\sqrt{2}}$$

$$\iint_R xy^2 dx dy = \int_{-\frac{1}{\sqrt{2}}}^{\frac{1}{\sqrt{2}}} \left[\int_{y^2}^{1-y^2} xy^2 dx \right] dy$$


$$= \int_{-\frac{1}{\sqrt{2}}}^{\frac{1}{\sqrt{2}}} \left[\frac{x^2}{2} y^2 \Big|_{x=y^2}^{x=1-y^2} \right] dy$$

$$= \int_{-\frac{1}{\sqrt{2}}}^{\frac{1}{\sqrt{2}}} \left[\frac{(1-y^2)^2}{2} y^2 - \frac{(y^2)^2}{2} y^2 \right] dy = \dots$$

Example Show that

$$R = \{ 0 \leq x \leq 1, \quad x \leq y \leq \sqrt{x} \}.$$

$$\int_0^1 \int_x^{\sqrt{x}} f(x, y) \, dy \, dx = \int_0^1 \int_{y^2}^y f(x, y) \, dx \, dy.$$



$$R = \{ 0 \leq y \leq 1, \quad y^2 \leq x \leq y \}$$

