



TOPIC 4. LOGISTIC REGRESSION

Introduction

- Logistic regression is a method that allows the prediction (classification) of discrete variables (outputs) based on a mixture of continuous and discrete predictors (inputs).
- It offers a non-linear relationship between the outputs and inputs (but linear after certain transformation– generalized linear models)
 - *Example: the probability of heart disease changes very little with a ten-point difference among people with low-blood pressure, but a ten point change can mean a drastic change in the probability of heart disease in people with high blood-pressure.*

Introduction

- Application criteria: the limitation on using the logistic regression is that the output should be discrete.

Terms used in logistic regression

- Odds - like probability, a measure of the likelihood of an event.
 - *Odds are usually written as “1 to 4 odds” which is equivalent to 1 out of five or .20 probability or 20% chance.*
- Odds ratio – the ratio of the odds over 1 – the odds. The probability of winning over the probability of losing. 1 to 4 odds equates to an odds ratio of $.20/.80 = .25$.
- Logit – this is the natural log of an odds ratio; often called a log odds.

Introduction

- $Y = \text{A BINARY RESPONSE (Discrete Variable)}$, p and $1 - p$
 - $1 \text{ POSITIVE RESPONSE (Success)} \rightarrow p$
 - $0 \text{ NEGATIVE RESPONSE (failure)} \rightarrow 1 - p$
- $\text{MEAN}(Y) = p$, observed proportion of successes
- $\text{VAR}(Y) = p(1 - p)$, maximized when $p = .50$, variance depends on mean
- $X = \text{ANY TYPE OF PREDICTOR} \rightarrow \text{Continuous, Dichotomous, Polytomous.}$
- p depends on X .

The Logistic Function

$$p(X) = \frac{e^u}{1 + e^u}$$


- Where $p(X)$ is the estimated probability that Y is in the success category and u is the regular linear regression equation:

$$u = b_0 + b_1X$$

The Logistic Function

- Simple case

$$p(X) = \frac{e^{b_0 + b_1 X}}{1 + e^{b_0 + b_1 X}}$$



The Logit Function

- By algebraic manipulation, the logistic regression equation can be written in terms of an odds ratio for success:

$$\frac{p(X)}{1 - p(X)} = e^{b_0 + b_1 X}$$

- Taking the natural log of both sides, we can write the equation in terms of logits (log-odds):

$$\ln \frac{p(X)}{1 - p(X)} = b_0 + b_1 X$$

Logistic Regression

- The response (binary)

$$Y \sim \text{Bernoulli}(p(X))$$

- Probability of success:


$$p(X) = \frac{e^{b_0 + b_1 X}}{1 + e^{b_0 + b_1 X}} \qquad \ln \frac{p(X)}{1 - p(X)} = b_0 + b_1 X$$

Latent Variable Model

$$Z = b_0 + b_1X + \varepsilon$$

$$Y = \begin{cases} 1, & \text{if } Z > 0 \\ 0, & \text{if } Z \leq 0 \end{cases}$$

- Where the random error ε follows a logistic distribution.
- This latent variable model is equivalent to the standard logistic regression.



Interpretation

$$p(X) = \frac{e^{b_0 + b_1 X}}{1 + e^{b_0 + b_1 X}} \qquad \ln \frac{p(X)}{1 - p(X)} = b_0 + b_1 X$$

- If $b_1 = 0$, there is no relationship between the response and the predictor.
- If $b_1 > 0$, when X gets larger so does the probability of success.
- If $b_1 < 0$, when X gets larger, the probability of success gets smaller.
- How much bigger or smaller depends on value of the slope.

Estimating Coefficients

- Given a dataset $\{(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)\}$, find estimates for b_0, b_1 .
- Maximum likelihood estimation (MLE): find \hat{b}_0, \hat{b}_1 that maximize the likelihood function

$$L(b_0, b_1) = \prod_{i=1}^n P(y_i) = \prod_{i:y_i=1} P(y_i = 1) \prod_{i':y_{i'}=0} P(y_{i'} = 0)$$

$P(y_i = 1) = \frac{e^{b_0 + b_1 x_i}}{1 + e^{b_0 + b_1 x_i}}$

$P(y_{i'} = 0) = \frac{1}{1 + e^{b_0 + b_1 x_{i'}}}$

Example

- Assume the dataset has 5 samples

Patient	Risk score X	Surgery Outcome Y (survival/0;death/1)
1	3	0
2	10	1
3	6	1
4	8	1
5	0	0

$$P(y_1 = 0) = \frac{1}{1 + e^{b_0 + 3b_1}}$$

$$P(y_5 = 0) = \frac{1}{1 + e^{b_0 + 10b_1}}$$

$$P(y_2 = 1) = \frac{e^{b_0 + 10b_1}}{1 + e^{b_0 + 10b_1}}$$

$$P(y_3 = 1) = \frac{e^{b_0 + 6b_1}}{1 + e^{b_0 + 6b_1}}$$

$$P(y_4 = 1) = \frac{e^{b_0 + 8b_1}}{1 + e^{b_0 + 8b_1}}$$

$$L(b_0, b_1) = \frac{1}{1 + e^{b_0 + 3b_1}} \cdot \frac{1}{1 + e^{b_0 + 10b_1}} \cdot \frac{e^{b_0 + 10b_1}}{1 + e^{b_0 + 10b_1}} \cdot \frac{e^{b_0 + 6b_1}}{1 + e^{b_0 + 6b_1}} \cdot \frac{e^{b_0 + 8b_1}}{1 + e^{b_0 + 8b_1}}$$

Significance Test

- Hypothesis testing

$H_0: b_1 = 0$ (there is a relationship between X and Y)

$H_1: b_1 \neq 0$ (no relationship between X and Y)

- Z (Wald) test

$$W = \frac{\hat{b}_1}{SD(\hat{b}_1)} \sim N(0, 1)$$

Extension

- Multiple logistic regression

$$\ln \frac{p(X)}{1 - p(X)} = b_0 + b_1 X_1 + b_2 X_2 + \dots + b_k X_k$$

- Multinomial logistic regression ($Y = 0, 1, 2, \dots$)

$$\ln \frac{p_{Y=1}(X)}{p_{Y=0}(X)} = \dots \quad \ln \frac{p_{Y=2}(X)}{p_{Y=0}(X)} = \dots \quad \ln \frac{p_{Y=3}(X)}{p_{Y=0}(X)} = \dots$$

- Ordinal (ordered) logistic regression ($Y = 0, 1, 2, \dots$)

$$\ln \frac{p_{Y \leq 0}(X)}{1 - p_{Y \leq 0}(X)} = \dots \quad \ln \frac{p_{Y \leq 1}(X)}{1 - p_{Y \leq 1}(X)} = \dots \quad \ln \frac{p_{Y \leq 2}(X)}{1 - p_{Y \leq 2}(X)} = \dots$$

Generalized Linear Models

- Logistic regression is a special type of GLM.
- General form of GLMs:

$$Y \sim \text{Probability Distribution}(\mu)$$
$$g(\mu) = b_0 + b_1 X$$

Example: Logistic regression

$$Y \sim \text{Bernoulli}(p)$$

$$g(p) = \ln \frac{p}{1-p} = b_0 + b_1 X$$

- g is called link function. It transforms the mean of the probability distribution to a linear function of X .
- $E(Y|X) = g^{-1}(b_0 + b_1 X)$ vs. $E(Y|X) = b_0 + b_1 X$ (linear regression)

Probit Regression

- Probit regression is another popular model for binary responses.
- Probit model:

$$Y \sim \text{Bernoulli}(p)$$

$$p = \Phi(b_0 + b_1 X)$$

$$g(p) = \Phi^{-1}(p) = b_0 + b_1 X$$

- Φ is the cumulative distribution of standard normal distribution.

Latent Variable Model of Probit Regression


$$Z = b_0 + b_1X + \varepsilon$$

$$Y = \begin{cases} 1, & \text{if } Z > 0 \\ 0, & \text{if } Z \leq 0 \end{cases}$$

- Where the random error ε follows a standard normal distribution. So the model of Z is a linear regression model.
- This latent variable model is equivalent to the probit regression.

Logit vs. Probit Regression

- Logit function and probit function have similar shapes.



(from datacamp.com)

Summary

- Generalized linear models including logistic regression is an extension of probability models to incorporate the effects of predictors.
- These models are easy to understand and have good interpretation.