

MA1201 Calculus and Basic Linear Algebra II
Problem Set 2 Techniques of Integration
Problem 1 (Method of substitution)

Compute the following integral using method of substitution:

$$(b) \int x^2 \sec(1 - 2x^3) dx$$

$$\text{let } y = 1 - 2x^3 \Rightarrow \frac{dy}{dx} = -6x^2 \Rightarrow dx = -\frac{1}{6x^2} dy$$

$$\begin{aligned} \int x^2 \sec(1 - 2x^3) dx &= \int x^2 \sec(1 - 2x^3) \left(-\frac{1}{6x^2} dy\right) = -\frac{1}{6} \int \sec y dy \\ &= -\frac{1}{6} \ln |\sec y + \tan y| + C \\ &= -\frac{1}{6} \ln |\sec(1 - 2x^3) + \tan(1 - 2x^3)| + C. \end{aligned}$$

$$(d) \int x \cos^2(x^2) dx$$

$$\text{let } y = x^2 \Rightarrow \frac{dy}{dx} = 2x \Rightarrow dx = \frac{1}{2x} dy$$

$$\begin{aligned} \int x \cos^2(x^2) dx &= \int x \cos^2(x^2) \left(\frac{1}{2x} dy\right) = \frac{1}{2} \int \cos^2 y dy \\ &= \frac{1}{2} \int \frac{1}{2} [\cos(y+y) + \cos(y-y)] dy = \frac{1}{4} \int (\cos(2y) + 1) dy \\ &= \frac{1}{4} \cdot \left(\frac{1}{2} \sin(2y) + y\right) + C \\ &= \frac{1}{8} \sin(2x^2) + \frac{1}{4} x^2 + C. \end{aligned}$$

$$(f) \int \frac{e^{2x}}{(1+e^x)^3} dx$$

$$\text{let } y = 1 + e^x \Rightarrow \frac{dy}{dx} = e^x \Rightarrow dx = \frac{1}{e^x} dy$$

$$\begin{aligned} \int \frac{e^{2x}}{(1+e^x)^3} dx &= \int \frac{e^{2x}}{(1+e^x)^3} \cdot \left(\frac{1}{e^x} dy\right) = \int \frac{e^x}{(1+e^x)^3} dy = \int \frac{y-1}{y^3} dy \\ &= \int (y^{-2} - y^{-3}) dy = \frac{y^{-2+1}}{-2+1} - \frac{y^{-3+1}}{-3+1} + C \\ &= -\frac{1}{y} + \frac{1}{2y^2} + C = -\frac{1}{1+e^x} + \frac{1}{2(1+e^x)^2} + C. \end{aligned}$$

$$(h) \int_1^5 \frac{\sin^2(\ln x)}{x} dx$$

$$\text{let } y = \ln x \Rightarrow \frac{dy}{dx} = \frac{1}{x} \Rightarrow dx = x dy$$

$$x=5, y=\ln 5, \quad x=1, y=\ln 1=0$$

$$\begin{aligned} \int_1^5 \frac{\sin^2(\ln x)}{x} dx &= \int_0^{\ln 5} \frac{\sin^2(\ln x)}{x} \cdot x dy = \int_0^{\ln 5} \sin^2 y dy \\ &= \int_0^{\ln 5} -\frac{1}{2} [\cos(y+y) - \cos(y-y)] dy \\ &= \int_0^{\ln 5} -\frac{1}{2} [\cos(2y) - 1] dy = -\frac{1}{2} \cdot \left[\frac{1}{2} \sin(2y) - y \right] \Big|_0^{\ln 5} \\ &= -\frac{1}{4} \sin(2 \ln 5) + \frac{1}{2} \ln 5 \end{aligned}$$

$$(i) \int \frac{2x+1}{x^2-2x+5} dx$$

$$\text{let } y = x^2 - 2x + 5 \Rightarrow \frac{dy}{dx} = 2x - 2 \Rightarrow dx = \frac{1}{2x-2} dy$$

$$\int \frac{2x+1}{x^2-2x+5} dx = \int \frac{2x-2}{x^2-2x+5} dx + \int \frac{3}{x^2-2x+5} dx \quad \rightarrow (\tan^{-1} x)' = \frac{1}{1+x^2}$$

$$= \int \frac{2x-2}{x^2-2x+5} \left(\frac{1}{2x-2} dy \right) + 3 \int \frac{1}{(x-1)^2+4} dx$$

$$= \int \frac{1}{y} dy + \frac{3}{4} \int \frac{1}{1 + \left(\frac{x-1}{2}\right)^2} dx$$

$$= \ln|y| + C_1 + \frac{3}{4} \cdot \frac{1}{\frac{1}{2}} \tan^{-1}\left(\frac{x-1}{2}\right) + C_2$$

$$= \ln|x^2-2x+5| + \frac{3}{2} \tan^{-1}\left(\frac{x-1}{2}\right) + C$$

$$(l) \int \frac{1}{x^2 \sqrt{1-x^2}} dx \quad \sin^2 \theta + \cos^2 \theta = 1.$$

$$\text{let } x = \sin \theta \Rightarrow \frac{dx}{d\theta} = \cos \theta \Rightarrow dx = \cos \theta d\theta$$

$$\int \frac{1}{x^2 \sqrt{1-x^2}} dx = \int \frac{1}{\sin^2 \theta \cdot \cos \theta} \cdot \cos \theta d\theta = \int \frac{1}{\sin^2 \theta} d\theta$$

$$\text{let } u = \frac{\cos \theta}{\sin \theta} \Rightarrow \frac{du}{d\theta} = \frac{-\sin \theta \sin \theta - \cos \theta \cos \theta}{\sin^2 \theta} = -\frac{1}{\sin^2 \theta} \Rightarrow d\theta = -\sin^2 \theta du$$

$$= \int \frac{1}{\sin^2 \theta} \cdot (-\sin^2 \theta du) = \int -1 du = -u + C = -\cot \theta + C$$

$$= -\frac{\sqrt{1-x^2}}{x} + C$$

$$x = \sin \theta$$

$$\cos \theta = \sqrt{1-x^2}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{\sqrt{1-x^2}}{x}$$

$$(n) \int \frac{3x}{\sqrt{4x^2+1}} dx$$

$$\text{let } y = 4x^2 + 1 \Rightarrow \frac{dy}{dx} = 8x \Rightarrow dx = \frac{1}{8x} dy$$

$$\begin{aligned} \int \frac{3x}{\sqrt{4x^2+1}} dx &= \int \frac{3x}{\sqrt{4x^2+1}} \cdot \left(\frac{1}{8x} dy\right) = \frac{3}{8} \int \frac{1}{\sqrt{y}} dy = \frac{3}{8} \int y^{-\frac{1}{2}} dy = \frac{3}{8} \cdot 2y^{\frac{1}{2}} + C \\ &= \frac{3}{4} \sqrt{4x^2+1} + C \end{aligned}$$

$$(p) \int \frac{1}{(x^2+6x+10)^{\frac{3}{2}}} dx$$

$$\text{let } y = x^2 + 6x + 10 \Rightarrow \frac{dy}{dx} = 2x + 6 \Rightarrow dx = \frac{1}{2x+6} dy \cdot x$$

$$\int \frac{1}{(x^2+6x+10)^{\frac{3}{2}}} dx = \int \frac{1}{((x+3)^2+1)^{\frac{3}{2}}} dx = \int \frac{1}{(\tan^2 \theta + 1)^{\frac{3}{2}}} \cdot \sec^2 \theta d\theta$$

$$\text{let } x+3 = \tan \theta \Rightarrow \frac{dx}{d\theta} = \sec^2 \theta \Rightarrow dx = \sec^2 \theta d\theta$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$= \int \frac{1}{(\sec^2 \theta)^{\frac{3}{2}}} \cdot \sec^2 \theta d\theta = \int \frac{1}{\sec \theta} d\theta = \int \cos \theta d\theta = \sin \theta + C$$

$$= \frac{x+3}{\sqrt{x^2+6x+10}} + C$$

$$\left. \begin{aligned} \tan \theta &= x+3 \\ \sec \theta &= \sqrt{x^2+6x+10} \end{aligned} \right\} \Rightarrow \sin \theta = \frac{\tan \theta}{\sec \theta}$$

$$(r) \int \sin^3 x \cos^5 x dx$$

$$\text{let } y = \sin x \Rightarrow \frac{dy}{dx} = \cos x \Rightarrow dx = \frac{1}{\cos x} dy. \rightarrow \cos^2 x = 1 - \sin^2 x.$$

$$\begin{aligned} \int \sin^3 x \cos^5 x dx &= \int \sin^3 x \cos^4 x \cdot \left(\frac{1}{\cos x} dy\right) = \int \sin^3 x \cos^4 x dy \\ &= \int y^3 (1-y^2)^2 dy = \int y^3 (1-2y^2+y^4) dy = \int (y^3 - 2y^5 + y^7) dy \\ &= \frac{y^4}{4} - \frac{2}{6} y^6 + \frac{1}{8} y^8 + C. \\ &= \frac{\sin^4 x}{4} - \frac{\sin^6 x}{3} + \frac{\sin^8 x}{8} + C. \end{aligned}$$

Problem 2 (Integration by parts)

Compute the following integrals using integration by parts:

$$(b) \int_1^e \sqrt{x} \ln x dx$$

$$u = \ln x, dv = \sqrt{x} dx \Rightarrow v = \int dv = \int \sqrt{x} dx = \frac{2}{3} x^{\frac{3}{2}}$$

$$\begin{aligned} \int_1^e \sqrt{x} \ln x dx &= \int_1^e \ln x \sqrt{x} dx = \frac{2}{3} x^{\frac{3}{2}} \ln x \Big|_1^e - \int_1^e \frac{2}{3} x^{\frac{3}{2}} d(\ln x) \\ &= \frac{2}{3} e^{\frac{3}{2}} \ln e - \frac{2}{3} \ln 1 - \int_1^e \frac{2}{3} x^{\frac{3}{2}} \cdot \frac{1}{x} dx \\ &= \frac{2}{3} e^{\frac{3}{2}} - \int_1^e \frac{2}{3} x^{\frac{1}{2}} dx = \frac{2}{3} e^{\frac{3}{2}} - \frac{2}{3} \cdot \frac{2}{3} x^{\frac{3}{2}} \Big|_1^e \\ &= \frac{2}{3} e^{\frac{3}{2}} - \frac{4}{9} e^{\frac{3}{2}} + \frac{4}{9} \cdot 1 = \frac{2}{9} e^{\frac{3}{2}} + \frac{4}{9}. \end{aligned}$$

$$(d) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x \sin^2 x dx = \int x \left[-\frac{1}{2} [\cos(x+x) - \cos(x-x)] \right] dx$$

$$= -\frac{1}{2} \int (x \cos 2x - x) dx = -\frac{1}{2} \int x \cos 2x dx + \frac{1}{2} \int x dx$$

$$u = x$$

$$dv = \cos 2x dx$$

$$v = \int dv = \int \cos 2x dx$$

$$= \frac{1}{2} \sin 2x$$

$$= -\frac{1}{2} \left[\frac{1}{2} x \sin 2x - \int \frac{1}{2} \sin 2x \cdot dx \right] + \frac{1}{2} \int x dx$$

$$= -\frac{1}{4} x \sin 2x + \frac{1}{2} \cdot \frac{1}{2} \cdot \left(-\frac{1}{2} \cos 2x\right) + C_1 + \frac{1}{2} \cdot \frac{1}{2} x^2 + C_2$$

$$= -\frac{1}{4} x \sin 2x - \frac{1}{8} \cos 2x + \frac{1}{4} x^2 + C.$$

