

Feedback — Quiz 2

[Help](#)

Thank you. Your submission for this quiz was received.

You submitted this quiz on **Thu 15 Jan 2015 6:22 AM PST**. You got a score of **10.00** out of **10.00**.

Question 1

Consider the following data with x as the predictor and y as the outcome.

```
x <- c(0.61, 0.93, 0.83, 0.35, 0.54, 0.16, 0.91, 0.62, 0.62)
y <- c(0.67, 0.84, 0.6, 0.18, 0.85, 0.47, 1.1, 0.65, 0.36)
```

Give a P-value for the two sided hypothesis test of whether β_1 from a linear regression model is 0 or not.

Your Answer	Score	Explanation
<input checked="" type="radio"/> 0.05296	✓ 1.00	
<input type="radio"/> 0.025		
<input type="radio"/> 2.325		
<input type="radio"/> 0.391		
Total	1.00 / 1.00	

Question 2

Consider the previous problem, give the estimate of the residual standard deviation.

Your Answer	Score	Explanation
<input type="radio"/> 0.05296		
<input type="radio"/> 0.4358		

☐ 0.3552

☒ 0.223



1.00

Total

1.00 / 1.00

Question 3

In the `mtcars` data set, fit a linear regression model of weight (predictor) on mpg (outcome). Get a 95% confidence interval for the expected mpg at the average weight. What is the lower endpoint?

Your Answer

Score

Explanation

☐ 21.190

☐ -6.486

☐ -4.00

☒ 18.991



1.00

Total

1.00 / 1.00

Question 4

Refer to the previous question. Read the help file for `mtcars`. What is the weight coefficient interpreted as?

Your Answer

Score

Explanation

☐ The estimated expected change in mpg per 1 lb increase in weight.

☐ The estimated 1,000 lb change in weight per 1 mpg increase.

☒



1.00

The estimated expected change in mpg per 1,000 lb increase in weight.

☐ It can't be interpreted without further information

Total	1.00 / 1.00
-------	----------------

Question 5

Consider again the `mtcars` data set and a linear regression model with mpg as predicted by weight (1,000 lbs). A new car is coming weighing 3000 pounds. Construct a 95% prediction interval for its mpg. What is the upper endpoint?

Your Answer	Score	Explanation
<input type="radio"/> -5.77		
<input checked="" type="radio"/> 27.57	✓ 1.00	
<input type="radio"/> 14.93		
<input type="radio"/> 21.25		
Total	1.00 / 1.00	

Question 6

Consider again the `mtcars` data set and a linear regression model with mpg as predicted by weight (in 1,000 lbs). A “short” ton is defined as 2,000 lbs. Construct a 95% confidence interval for the expected change in mpg per 1 short ton increase in weight. Give the lower endpoint.

Your Answer	Score	Explanation
<input type="radio"/> 4.2026		
<input type="radio"/> -6.486		
<input type="radio"/> -9.000		
<input checked="" type="radio"/> -12.973	✓ 1.00	
Total	1.00 / 1.00	

Question 7

If my X from a linear regression is measured in centimeters and I convert it to meters what would happen to the slope coefficient?

Your Answer	Score	Explanation
<input type="radio"/> It would get divided by 10		
<input checked="" type="radio"/> It would get multiplied by 100.	✓ 1.00	
<input type="radio"/> It would get divided by 100		
<input type="radio"/> It would get multiplied by 10		
Total	1.00 / 1.00	

Question 8

I have an outcome, Y , and a predictor, X and fit a linear regression model with $Y = \beta_0 + \beta_1 X + \epsilon$ to obtain $\hat{\beta}_0$ and $\hat{\beta}_1$. What would be the consequence to the subsequent slope and intercept if I were to refit the model with a new regressor, $X + c$ for some constant, c ?

Your Answer	Score	Explanation
<input type="radio"/> The new slope would be $c\hat{\beta}_1$		
<input type="radio"/> The new slope would be $\hat{\beta}_1 + c$		
<input checked="" type="radio"/> The new intercept would be $\hat{\beta}_0 - c\hat{\beta}_1$	✓ 1.00	
<input type="radio"/> The new intercept would be $\hat{\beta}_0 + c\hat{\beta}_1$		
Total	1.00 / 1.00	

Question 9

Refer back to the mtcars data set with mpg as an outcome and weight (wt) as the predictor. About what is the ratio of the the sum of the squared errors, $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2$ when

comparing a model with just an intercept (denominator) to the model with the intercept and slope (numerator)?

Your Answer		Score	Explanation
<input checked="" type="radio"/> 0.25	✓	1.00	
<input type="radio"/> 0.75			
<input type="radio"/> 0.50			
<input type="radio"/> 4.00			
Total		1.00 / 1.00	

Question 10

Do the residuals always have to sum to 0 in linear regression?

Your Answer		Score	Explanation
<input type="radio"/> The residuals never sum to zero.			
<input type="radio"/> The residuals must always sum to zero.			
<input type="radio"/> If an intercept is included, the residuals most likely won't sum to zero.			
<input checked="" type="radio"/> If an intercept is included, then they will sum to 0.	✓	1.00	
Total		1.00 / 1.00	