

Coulomb + Newton: Towards a Cosmology

Claes Johnson¹

¹KTH Royal Institute of Technology, Stockholm, Sweden , claesjohnson@gmail.com

August 13, 2026

Abstract

RealQM/RealNucleus models nuclei, atoms and molecules as *non-overlapping unit charge densities* — protons carrying the mass m_p and electrons the mass m_e — interacting by Coulomb potentials. Being continuum mechanics, the model has the same mathematical form as the models of the macroscopic materials those atoms build, so a hierarchy of continuum models over increasing scales follows from it. Adding **Newtonian gravitation**, which acts on the energy of that matter and matters only in the large, carries the hierarchy to galactic scales: one form of model from nuclei to the cosmic web. Cosmology is its natural arena, being the subject in which the nuclear scale meets the astronomical.

The chain is computed throughout. A small perturbation of the electric potential φ_E has its curvature read as charge, quantized by indivisibility into units of ± 1 in two spatial extents — the compact heavy proton, m_p , and the diffuse light electron, m_e . Coulomb builds nuclei, atoms and molecules from them. Newtonian gravitation then acts on the resulting energy density through its departure from the mean, $\nabla^2 \varphi_g = 4\pi G(\rho - \bar{\rho})$, a source of both signs although the matter is positive throughout, so the large scale separates into the overdense **cosmic web** and the underdense **voids**. The background source vanishes and the expansion **coasts**, $a \propto t$, $q = 0$ — falsifiable, and separated from Λ CDM by ≈ 0.2 mag of shape on the supernova Hubble diagram. Only at the end is it recorded that the two forces are one two-valued Poisson form read with opposite signs; that parallel is used nowhere in the construction. Neither dark component is accounted for, and the large-scale inhomogeneity is an input.

Keywords: RealQM; RealNucleus; two-valued Poisson; Coulomb and Newton; charge quantization; density contrast; gravitation.

1 Introduction

RealQM formulates quantum mechanics as multiphase 3D continuum mechanics — unit-charge densities on non-overlapping domains separated by free boundaries, minimising the Coulomb energy [2], developed in the treatise *Real Quantum Mechanics* [1]. Its nuclear extension RealNucleus revives the pre-1932 proton–electron picture (neutron = proton + electron) and reproduces nuclear binding from pure Coulomb interaction, with no strong-force postulate [3]. Because these are continuum-mechanical models — densities on domains, forces from potentials, free boundaries where phases meet — they have the same mathematical form as the models of bulk matter, and a hierarchy over increasing scales follows with no change of mathematical character on the way up. Standard quantum mechanics, formulated in $3N$ -dimensional configuration space, has no such continuum limit handing directly to the mechanics of materials.

This paper carries that hierarchy to its top. **Newtonian gravitation** acts on the energy of the matter RealQM builds and matters only in the large, so adding it extends the same model to galactic scales — a seamless extension, in that nothing about the mathematics changes: one more Poisson equation, one more force term in the same continuum mechanics. Where the

extension leads is cosmology, and cosmology is its natural arena, being the subject in which the nuclear scale meets the astronomical. Stars shine by nuclear fusion: a process at 10^{-15} m sets the luminosity and chemical output of objects at 10^9 m and thereby the evolution of galaxies at 10^{21} m. Light-element abundances are fixed by nuclear reaction rates yet read off the sky as cosmological data. And the dark sector, though *posed* at galactic scales, is *answered* at the smallest — dark matter sought as a new particle in particle colliders and underground detectors — so the instrument aimed at the largest missing mass in the universe is a machine that collides protons. A model that spans from nuclei to galaxies is therefore not an ambition imposed on the subject but what the subject already requires.

The main idea, and the plan. *One small perturbation of one potential, followed upward by computation in one form of model, yields quantized charges and masses, then nuclei, atoms and molecules, and finally a web-like cosmos — and every step is computed, not assumed.* The continuity of that chain is the point, more than any one of its links, and the paper follows it in the order the model builds it. Section 2 states the exact model, both tiers. Sections 3–4 start the chain with the perturbation of φ_E ; Sections 5–7 hand the resulting protons and electrons to RealQM and RealNucleus; Section 8 discretises both tiers; Section 9 adds gravitation on the energy density of the matter built; Sections 10–12 ask what more a cosmology would require — more than is delivered here. Only at the end (Section 13) is the parallel between the two forces recorded, as an observation about the finished construction. Security decreases along the chain and is marked as it does: the seed is a posit, the large-scale inhomogeneity an input, the cosmological claims weakest of all. Nothing along the way is static: the equilibria are balances maintained by motion, and the large-scale sector coarsens without ever settling.

2 The exact model

The macroscale model is **small-viscosity** (or **vanishing-viscosity**) compressible self-gravitating flow. With $\rho \geq 0$ the mass/energy density of matter, $\mathbf{m} = \rho \mathbf{u}$ the momentum, e the internal energy and p the pressure,

$$\begin{aligned}\dot{\rho} + \nabla \cdot \mathbf{m} &= 0, \\ \dot{\mathbf{m}} + \nabla \cdot (\mathbf{m} \mathbf{u}) + \nabla p + \rho \nabla \varphi_g &= 0, \\ \dot{e} + \nabla \cdot (e \mathbf{u}) + p \nabla \cdot \mathbf{u} &= 0,\end{aligned}\tag{1}$$

closed by $\mathbf{u} = \mathbf{m}/\rho$, $p = \gamma_s e$, $e = \rho T$, and coupled to gravitation through the Poisson equation with the **mean-subtracted** source,

$$\nabla^2 \varphi_g = 4\pi G (\rho - \bar{\rho}), \quad \bar{\rho} = \langle \rho \rangle.\tag{2}$$

Equation (1) is the Euler system with a gravitational body force. Its right-hand sides are zero because no dissipation is put in as a modelling choice; solutions are not smooth — the flow is turbulent and collapse forms shocks — so what is computed is a small-viscosity solution, the viscosity being that of the discretisation, mesh-dependent rather than physical. The meaningful output is therefore what is *stable* as that viscosity is decreased, which also settles a question of physics: collapse is arrested by the adiabatic index and not by dissipation, since the arrest must survive the limit while an artificial-viscosity arrest does not (Section 9). Because ρ stays positive, $\mathbf{u} = \mathbf{m}/\rho$ is regular and no density floor or clamp is needed anywhere — unlike a genuinely signed density, for which $\mathbf{u}$ is singular at every zero crossing, precisely at the interfaces where the structure lives.

The $\bar{\rho}$ in (2) makes the gravitational source take both signs although the matter is positive throughout. It is not an added assumption: φ_g **exists only if its source has zero mean**, so some constant must be subtracted, and $\bar{\rho} = \langle \rho \rangle$ is the simplest such constant (Section 9).

The small-scale model is the Coulomb energy of RealQM: one non-negative density ψ_i per electron, each on its own domain Ω_i , the domains non-overlapping, $\Omega_i \cap \Omega_j = \emptyset$, separated by *free* boundaries. The state minimises

$$E[\{\psi_i\}, \{\Omega_i\}] = \sum_i \int_{\Omega_i} \left(\frac{1}{2} |\nabla \psi_i|^2 - \sum_a \frac{Z_a}{|x - x_a|} \psi_i^2 \right) dx + \frac{1}{2} \sum_{i \neq j} \iint \frac{\psi_i^2(x) \psi_j^2(y)}{|x - y|} dx dy, \quad (3)$$

subject to $\int_{\Omega_i} \psi_i^2 = 1$, the domains free to move: the interfaces settle where the energy says. The sum $i \neq j$ carries **no self-interaction** term — its role is taken by the localisation energy $\frac{1}{2} \int |\nabla \psi_i|^2$ — and the problem lives in **three** dimensions however many electrons there are, one field each. Both (1)–(2) and (3) are discretised on the same kind of grid in Section 8.

3 The seed: charge and mass from a perturbation

In the standard account the origin of matter is an initial hot dense state, with inflation supplying the fluctuation spectrum. Here that role is played by a **small perturbation of the electric potential** φ_E , magnified by the Laplacian $\rho_E = -\varepsilon_0 \nabla^2 \varphi_E$ into $\pm$ charge and hence into protons, electrons and all Coulombic matter. That is all the seed is asked to do. It creates the *matter*; it is not asked to lay out the large-scale *distribution* of that matter, and the gravitational sector makes no use of it.

The seed’s amplitude at the creation scale is fixed rather than chosen. If φ_E oscillates there with wavelength λ and amplitude A , each half-wavelength is a domain of one sign carrying charge $q \sim \varepsilon_0 A \lambda$; demanding $q = \pm e$ exactly — one indivisible quantum per oscillation — gives

$$A \sim \frac{e}{\varepsilon_0 \lambda}. \quad (4)$$

At $\lambda \sim 1$ fm this is $\sim 2 \times 10^7$ V, of the same order as the Coulomb potential $e/4\pi\varepsilon_0\lambda \approx 1.4 \times 10^6$ V of the single charge it creates. The small-scale sector therefore reduces to **one unknown length** λ , the same unknown as the proton size (Section 14).

And with the charges, the masses. A cloud’s energy is fixed by its extent (localisation cost against Coulomb attraction) and its inertia *is* that energy, E/c^2 (Section 7), so the two extents deliver **two inertial masses** and only two: the compact positive cloud is the heavy one, m_p , the diffuse negative one the light one, m_e . Quantization of charge carries quantization of mass with it — exactly two stable species because exactly two extents, so no intermediate mass appears any more than an intermediate charge does. Not settled here is the quantitative map: the observed $m_p/m_e \approx 1836$ is not reproduced by the naive inverse-size relation, which misses the proton’s compactness by about a factor of ten. The *count* of masses is a result; their *values* are open (Section 14).

Two clarifications. The seed replaces the Big Bang as the origin of *matter*, not as a beginning of time: there is no initial singularity, and equally no account of why the seed is there or how it evolves — φ_E has no equation of motion here. And the large-scale distribution of matter is not derived from it: a statistically homogeneous seed leaves ρ uniform up to counting noise ($\delta \sim N^{-1/2} \sim 10^{-34}$, thirty orders of magnitude short of what structure requires), so the large-scale inhomogeneity is an **input** to the gravitational sector, not an output of the electric one. Whether one seed could supply both — which would need its long wavelengths to modulate its short-wavelength power, i.e. local-type non-Gaussianity — is left to Section 14.

4 The arena

The equations are written in a **3D Euclidean coordinate system** — one of many possible, and nothing depends on which. It is a system of coordinates in which to express the fields, not an absolute space and not a physical medium; no ether is postulated and no frame is privileged. The only operator needed is the Laplacian. Charge is not postulated separately but is the curvature of the potential, $\rho_{\text{charge}} = -\varepsilon_0 \nabla^2 \varphi_E$, with two immediate consequences. **Both signs:** ρ_{charge} is positive where φ_E is concave, negative where convex. **Net zero:** $\int \nabla^2 \varphi_E dV$ is a surface term that vanishes, so the universe carries zero net charge automatically — equal numbers of protons and electrons, with no baryogenesis, and once the neutron is read as a proton–electron pair, in the far stronger *local* form of Section 5. Because the Laplacian multiplies each Fourier mode by k^2 , a short-wavelength oscillation becomes a large charge density.

5 The small scale: RealQM and RealNucleus

The seed has delivered charge in both signs, quantized at ± 1 , in two spatial extents: small dense positive and large diffuse negative charge, the **proton** and the **electron**. From here the whole charged world is built by (3), by Coulomb geometry alone.

Energy, not mass. In RealQM a particle is a charge cloud whose size is fixed by localisation cost against Coulomb attraction, and its inertia is its energy content E/c^2 — a *measured* identity (the Penning trap, the nuclear mass defect), not a relativistic postulate [4]. No primitive “mass” is invoked: charged matter carries energy, and energy gravitates. A contracted cloud costs more, so small clouds are energy-dense — the small proton is the heavy one. Whether size fixes energy *quantitatively* is another matter, left open; the cosmology needs only that gravity couples to energy.

Self-repulsion is kinetic energy. RealQM clouds do not overlap and carry no self-interaction. The role self-repulsion would play, resisting collapse to a point, is carried by the localisation energy $\frac{1}{2} \int |\nabla \psi|^2$: compressing a cloud sharpens its gradients and costs kinetic energy, so it settles at finite size. A free electron spreads; a bound one takes the size where localisation balances attraction.

Charge is quantized: ± 1 the only possibility. To exclude an electron’s Coulomb energy with *itself*, the “self” must be well defined: there must be an *indivisible* entity whose interior is left out of the Coulomb sum. A divisible continuum would make “self” ambiguous — is half a cloud a self? do the halves repel? — and the prescription ill-posed. So removing self-repulsion *requires* indivisible units. A unit is identified by its spatial occupation, and for all electrons, and equally all protons, to be the same entity, each domain must carry the same charge. Every electron then carries -1 and every proton $+1$, and no intermediate charge can exist. Granted that the no-self-interaction law is one *universal* law rather than an ad-hoc per-blob subtraction, quantization is a consequence, not a postulate — where the Standard Model must *import* it, anomaly cancellation fixing only the $\pm$ ratio and true quantization needing a Dirac monopole or a compact unifying group, both unobserved. What this does not fix is the *magnitude* of the unit, nor the fractional charges of confined quarks, outside RealQM’s no-substructure scope.

Nuclei and atoms, one mechanism at two scales. With a neutron taken as $p + e$, a neutral atom (A, Z) holds A protons and A electrons: $A - Z$ *compact* inside the nucleus and Z *expanded* in orbitals. The electron occupies two size-phases, its extent adapting to the well: compact in a deep one (nuclear, $\sim \text{fm}$, $\sim \text{MeV}$), expanded in a shallow one (atomic, $\sim \text{\AA}$, $\sim \text{eV}$). Chemistry and nuclear physics are then one Coulomb free-boundary mechanism at two scales: a molecule is

nuclei glued by expanded electrons, a nucleus is protons glued by compact ones. RealNucleus reproduces nuclear binding this way — the He-4/deuteron binding ratio comes out 13.1 against an experimental 12.7, from Coulomb geometry alone [3] — and light-nucleus fusion appears as a rearrangement of protons and electrons, not the action of a separate force.

Neutrality is local. Because a neutron *is* a bound proton–electron pair, every neutron carries its own compensating charge: matter is neutral *pointwise*, not merely on average. Standard cosmology gets global neutrality from an unexplained balance of baryogenesis and lepton number; here it is automatic twice over, globally from the net-zero seed and locally because the neutron cannot unbalance a count of $\pm$ pairs.

6 Complexity, and the two domains

The world is built not on a charge imbalance (charge is exactly balanced) but on a *size* imbalance: the proton has a fixed small size while the electron’s is variable, adapting to its well. The fixed proton supplies anchors; the variable electron *spans scales* — spreading to bind atoms, compacting to bind nuclei — producing the nested hierarchy that *is* complexity. With a single fixed size only one scale exists and nothing complex is built, so the nuclear/atomic separation is a **precondition for complexity**: an anthropic reason why the scales are separated, though not a derivation of how far. Since a neutron is structurally $p + e$ it is heavier than a proton by construction, so proton and hydrogen stability is automatic — the usual fine-tuning bound “ α must not be too large or the proton decays” does not arise.

Why the proton must be compact. A caged electron, held by several protons, sits at zero localisation and takes the proton size R_p ; a free one takes the atomic size a_0 , whence $a_0/R_p \approx 1/\alpha^2$: the two domains barely overlap because the two branches are separated by $\alpha^{-2} \approx 2 \times 10^4$. The picture also states its own validity limit. Since inertia is energy, a compact electron bound by Coulomb at the nuclear scale has binding $\sim e^2/R_p$, and requiring it to stay below the electron’s own energy E_e gives $R_p \gtrsim e^2/E_e$: a floor under the proton size, a plain comparison of two measured energies needing no appeal to α . Nature sits essentially *at* the floor — at the deuteron radius (~ 2.1 fm) the Coulomb binding $e^2/R_p \approx 0.7$ MeV already exceeds $E_e \approx 0.5$ MeV — so the nuclear scale saturates the theory’s own validity limit, which is why β -decay is where the static-cloud description is most stretched.

7 Inertia is energy; the local speed limit

Throughout, inertia is energy content: $\mathbf{p} = (E/c^2)\mathbf{v}$, the measured relation of the Penning trap [4], with $\mathbf{F} = \dot{\mathbf{p}}$ governing the response. In the small the force is electromagnetic and couples to *charge*; in the large it is gravity and couples to *mass* $= E/c^2$. One mechanics, one inertia, two forces reading two properties of the same matter.

The constant c enters *only* the electromagnetic sector, through the field momentum of a moving charge. Hence a **local speed limit**: feeding energy to a charge only increases its inertia E/c^2 without bound as $v \rightarrow c$. But Newtonian gravitation is c -free and imposes no speed limit on the large-scale motion of neutral bodies: two galaxies may recede at relative speed $> c$, a global relative velocity of gravitating matter rather than a local signal, violating no local causality — as in observational cosmology, where distant galaxies recede faster than light without any local law being broken.

8 The computational model

Both tiers are solved by the same numerical primitive: a Poisson solve coupled to a relaxation on a uniform Cartesian grid, ∇^2 the standard 7-point stencil, the matter field then advanced — downhill in energy at the small scale, in time at the large one. No basis sets, no wavefunctions in $3N$ dimensions, no fitted parameters.

	Small scale (Coulomb)	Large scale (Newton)
unknown	densities $\psi_i \geq 0$ on domains Ω_i	contrast ρ , momentum m , energy e
field solve	$-\nabla^2 \varphi = \rho_{\text{charge}}/\varepsilon_0$	$\nabla^2 \varphi_g = 4\pi G(\rho - \bar{\rho})$
solver	multigrid V-cycles, Jacobi smoother	red-black SOR, $\omega = 2/(1 + \sin \pi/N)$
relaxation	steepest descent on the Coulomb energy	explicit compressible flow in time
free surfaces	interfaces $\partial\Omega_i$ move with the energy	the $\rho = 0$ contact wall
typical grid	100^3 per atom	$200^3 = 8 \times 10^6$ cells

At the small scale the cost grows *linearly* in electron number where a wavefunction method grows exponentially, and the Neumann-free interface — located by the flow rather than imposed — lets one code describe an atom, a molecule and, with the electron compacted, a nucleus. These solvers run in the browser (WebGPU).

At the large scale, since the model is small-viscosity flow the scheme supplies no physical viscosity: the computation *is* the regularisation, and a result counts only if it persists as that regularisation is weakened. Two numerical points carry physics. First, the velocity is taken from $|\rho|$, $u = m/(|\rho| + \rho_{\min})$, so **inertia is positive in both phases** while the sign enters only the gravitational coupling — the direct expression of the contrast reading, underdense regions being ordinary matter of which there is less than average. Taking the velocity from signed ρ instead produces the classic runaway: the negative phase cannot self-support and is consumed, the void fraction collapsing from 50% to a few percent. Second, transport must be conservative and upwind; with centred differencing the web survives but is over-diffusive and the filaments never sharpen.

What the computations establish. At the small scale they are quantitative: binding energies, ionisation energies and nuclear binding follow from the same free-boundary minimisation without adjustable parameters. At the large scale they are *morphological only* — dimensionless units, a periodic box, no expansion, no redshift, no calibration to any survey — and nothing in them tests the cosmological claims of Sections 10–12.

9 The large scale: Newtonian gravitation

The mass/energy density of matter — protons, electrons, their binding and radiation — is ρ , positive throughout. It is the source of the second force, but enters its Poisson equation only through its departure from the mean, (2), so the *source* takes both signs — positive where matter is denser than average, negative where sparser — while the matter itself is positive everywhere. There is no negative mass and no second species: the two signs are *excess* and *deficit* of one substance.

Why the subtraction is forced, and what is chosen. The zero mean is not an assumption added to Newton’s law; it is what makes φ_g exist. On an unbounded (or periodic) domain the Poisson equation $\nabla^2 \varphi_g = 4\pi G\sigma$ is solvable only if $\langle \sigma \rangle = 0$: for a uniform positive source the potential diverges, and the net force on a point depends on the order in which the infinite distribution is summed — the Neumann–Seeliger difficulty, which is not a peculiarity of this model but of Newtonian cosmology as such. Some constant must therefore be subtracted, and

no Newtonian cosmology escapes it. Standard practice supplies it as bookkeeping and calls it the Jeans swindle.

What is *chosen* is only which constant, and $\bar{\rho} = \langle \rho \rangle = M_{\text{tot}}/V_{\text{tot}}$ is the simplest — the one that makes the total gravitational charge of the universe vanish, matching the zero net charge of the Coulomb sector. Two consequences follow. First, $\bar{\rho}$ is a boundary condition on the universe as a whole, so the local force law refers to a globally fixed constant: the model is non-local in this one limited sense, and a mean over some finite radius R would instead make the dynamics depend on an arbitrary R . Second, $\bar{\rho}$ is time-dependent — under expansion it dilutes as a^{-3} with the matter, so the cancellation is preserved at every epoch.

The non-standard step is thus not the subtraction but the refusal to treat it as bookkeeping: here the mean genuinely carries no gravitational charge, and the consequences are followed to the end (Section 10). This is where the model parts company with general relativity rather than with Newton, since in general relativity a homogeneous background *does* curve spacetime and the existence argument above does not apply — the solvability difficulty is Newtonian. What that costs is stated in Section 12.

This is *not* modified gravity but a modified source. The force law is untouched — no altered $1/r^2$, no extra field, no acceleration-dependent coupling — and for a galaxy, where the cosmic mean lies thirty orders of magnitude below the local density, subtracting $\bar{\rho}$ changes nothing. What is non-standard is what counts as a source at the largest scale. The accurate description is *standard Newtonian dynamics with a non-standard cosmological boundary condition*.

The force-rule. Under **like sources attract, unlike repel**, all four combinations are realised and each is observed. Two overdense regions attract: clustering. Two underdense regions attract: *voids merge*, as they are seen to. An overdense and an underdense region repel: matter drains from voids into filaments. The large scale therefore separates into overdense and underdense domains — the luminous **cosmic web** of filaments and clusters, and the **voids**, which are not inert leftovers but push their surroundings outward. The asymmetry with the charge sector is worth marking: a positive charge can be isolated, whereas the two signs here are excess and deficit of one substance, meaningless without the mean to measure against. That is why charge comes in indivisible ± 1 units carrying an identity while gravitational “charge” is continuous and reference-dependent; the quantization argument belongs to the Coulomb sector and must not be carried over.

What sets the scale, and what arrests collapse. The web scale is the **Jeans length** $\lambda_J = 2\pi c_s / \sqrt{G\rho}$ with $c_s = \sqrt{\gamma_s T}$: a box below one Jeans length is pressure-supported and inert. What *arrests* collapse is the adiabatic index $\gamma_{\text{ad}} = 1 + \gamma_s$ through $M_J \propto c_s^3 / \sqrt{\rho}$: under adiabatic compression $c_s^2 \propto \rho^{\gamma_{\text{ad}}-1}$, so $M_J \propto \rho^{3(\gamma_{\text{ad}}-1)/2-1/2}$, rising with density when $\gamma_{\text{ad}} > 4/3$ and falling below it. Above $4/3$ collapse shuts itself off at finite scale; below, each collapsing region keeps fragmenting. (An earlier version attributed the arrest to an artificial shock viscosity. Numerical experiment does not support that: the shock-heating term destroys the web in a thermal runaway and worsens rather than converges under timestep refinement.)

Web formation is standard, and illustrated rather than claimed. Equations (1)–(2) are ordinary Newtonian self-gravitating gas dynamics with a mean-subtracted source; nothing in them is new. That such systems develop a filamentary web from a Gaussian seed is classical: by the Zel’dovich approximation [7] collapse proceeds anisotropically, since the tidal-tensor eigenvalues generically differ, so a region contracts first to a sheet, then a filament, then a node — the morphology made quantitative by Hessian-eigenvalue classification [8]. Figure 1 shows the model’s own equations doing this. Two features bear on how such pictures should be judged. The web is a *transient*: in a non-expanding box the filament fraction falls from 33% to 1% within a factor of three in time, and neither snapshot reproduces the volume fractions inferred for the

real web (voids $\sim 60\text{--}80\%$, walls $\sim 20\text{--}30\%$, filaments $\sim 10\text{--}20\%$). What regulates the rate in the observed universe is expansion, omitted here. And visual impression disagrees with volume statistics: the right panel *looks* more web-like yet is classified almost entirely void, thin bright structures dominating the eye while occupying little volume — a standing caution about web-like images generally.




Figure 1: Gravitational instability of the contrast source (2) from a Gaussian seed (128^3 , thin slab; bright = overdense, dark = underdense). *Left*: sheet–filament stage, 33% filament and 46% wall by volume. *Right*: the same run later, collapsed to nodes, 1% filament and 93% void. The morphology is the classical Zel’dovich sequence [7], not particular to this model. Non-expanding box, so the web stage is transient.

The honest debt runs the other way. Standard simulations reproduce the observed web using *cold dark matter* in an *expanding* background, neither available here. Baryons alone are the harder case: before recombination they are coupled to the radiation and cannot grow, and gas pressure suppresses structure below the Jeans scale — which is precisely the structure-formation argument *for* dark matter, and among the strongest. That these equations admit webs is no answer to it.

Mathematically the mechanism is a **phase separation**: ρ is an order parameter with two stable phases separated by an unstable state at $\rho = 0$, segregated not by local diffusion but by the long-range gravitational kernel — a nonlocal Cahn–Hilliard [12] of Ohta–Kawasaki type [13], but with the kernel attractive for like sources, so that where Ohta–Kawasaki’s repulsion *frustrates* coarsening, gravity *drives* it. The web is then a spinodal-decomposition pattern arrested at finite scale by $\gamma_{\text{ad}} > 4/3$. One qualification: the double well is **not** symmetric, since $\delta \geq -1$ (a void can at most be empty) while overdensities grow without bound, so filaments and voids are not statistical mirrors. That symmetry *would* hold for a genuinely signed mass density, and the asymmetry of real voids counts against that reading.

10 Voids, the background, and the spectrum

Void repulsion is real but is not dark energy. A region below the mean is a negative source in (2), so it pushes matter outward and grows: voids are actively driven, not merely evacuated. But this is a local, peculiar effect, present in any mean-subtracted gravity, and must not be confused with dark energy, whose defining role is *accelerated expansion of the background*. There the model is explicit and negative: with the mean not gravitating the Friedmann source vanishes, $\ddot{a} = 0$, and the expansion coasts ($q = 0$) rather than accelerating ($q_0 \approx -0.55$). **The model does not**

reproduce what dark energy was introduced to explain. What it offers instead is that no Λ need be introduced and that the coincidence problem does not arise: the cancellation is exact at every epoch rather than mysteriously near-exact now. Acceleration is *excluded*, not merely absent — both sides of the contrast are the same substance and dilute together as a^{-3} , so the cancellation holds for all time; acceleration requires $w < -1/3$ while exact cancellation forces $w = 0$.

Negative mass as a substance: set aside. An earlier form of this model, and work by others, takes the negative sector to be a substance rather than a deficit: Farnes [6] proposes a negative-mass fluid with continuous creation; the Janus model of Petit and d’Agostini [9] and Hossenfelder’s exchange-symmetric bimetric theory [10] give the two signs their own metrics. We do not follow this route. Such a mass would be a species never detected, whose repulsion conveniently places it where it cannot be seen; it would *lens* with the opposite sign, whereas void lensing is measured and consistent with ordinary underdensity; and a signed density makes $\mathbf{m}/\rho$ singular at every zero crossing, forcing regularizations that dominated the result in our own numerical experiments. The deficit reading needs none of them.

Inflation, and the spectrum. The Laplacian amplifies short wavelengths, $\rho_k \propto k^2 \varphi_k$: an inflation of the density *field*, not the exponential expansion of *space*, which addresses the horizon and flatness problems and is not reproduced here. One might expect the k^2 weighting to leave nothing at large scales. It does not: in power $P_\rho(k) = k^4 P_\varphi(k)$, so a seed with equal fluctuation power per octave ($P_\varphi \propto k^{-3}$, requiring no tuning) gives $P_\rho \propto k$, the standard form of the primordial matter spectrum. We claim no more than that the expected conflict does not arise; the observed departure from this form is not derived, and would have to be compared through a transfer function this model does not possess.

11 The cosmic sequence and the helium fraction

The two sectors exchange energy in both directions. *Upward*, the energy of the charged world is mass and sources gravity. *Downward*, collapse converts gravitational potential energy into heat, driving the electromagnetic processes: gravity supplies the energy, electromagnetism spends it as structure and light. As contracting proton–electron matter heats, deuterons and nuclei form ($\sim$ MeV), the Coulomb binding energy radiating away and appearing as the mass defect; on cooling, nuclei capture expanded electrons into atoms ($\sim$ eV), again radiating. At recombination the matter turns neutral, electromagnetism screens out, radiation streams free, and only then does slow gravitational aggregation build the web. A bound state survives once the bath can no longer ionise it, so the two electron phases switch on at two temperatures set by their binding energies, forcing the order plasma $\rightarrow$ nuclei $\rightarrow$ atoms.

epoch	process	temperature
plasma	free $\pm$ charges	$> 10^{10}$ K
nuclei	compact e captured $\rightarrow$ nuclei	$\sim 10^{10}$ K ($kT \sim$ MeV)
atoms	expanded e captured $\rightarrow$ atoms; radiation	~ 3000 K ($kT \sim$ eV)

Table 1: Two condensation epochs, separated by $\sim 10^6$ in temperature.

The helium fraction: a consistency, not a prediction. The electron’s two size-phases — compact (inside a neutron, since neutron = p + compact e) and expanded (atomic) — are interconverted by the two weak processes $p + e \rightarrow n$ and $n \rightarrow p + e$. Three quantities enter and all three are *inputs*: the energy gap E_* between the phases, which plays the role of the neutron–proton mass difference and is identified with it, $E_* \approx 1.29$ MeV (its absolute size is the open nuclear-scale problem — RealNucleus fixes dimensionless ratios, not the MeV scale); the

transition temperature $T_* \sim E_*/k \sim 10^{10}$ K; and the freeze-out temperature $T_f \approx 0.8$ MeV, set as in standard big-bang nucleosynthesis by the weak-rate-versus-expansion balance. In equilibrium n/p is the Boltzmann factor $e^{-E_*/kT}$, locking at freeze-out; a fraction β -decays back (survival factor d) and essentially all the rest end in He-4:

$$\left(\frac{n}{p}\right)_{\text{nuc}} = \frac{(n/p)_f d}{1 + (n/p)_f(1 - d)}, \quad Y = \frac{2(n/p)_{\text{nuc}}}{1 + (n/p)_{\text{nuc}}}. \quad (5)$$

With $E_* \approx 1.29$ MeV, $T_f \approx 0.8$ MeV and $d \approx 0.74$: $(n/p)_f = 0.199 \rightarrow 0.140 \rightarrow Y = 0.245$, the observed primordial helium. Nothing here is derived; RealNucleus's one genuine computed output, the He-4/deuteron binding ratio, does not enter Y . The content is that the proton–electron reinterpretation is *consistent* with the standard calculation.

12 Towards a cosmology: what would be required

Shared with the standard account. Ordinary matter is Coulombic and gravitation attractive between masses; nuclei and atoms form and later assemble into filamentary structure; $Y \approx 0.25$ is accommodated.

Different. No initial singularity and no hot Big Bang, hence no microwave background as a red-shifted afterglow; no inflationary expansion of space; no dark-matter particle; the neutron composite rather than elementary; nuclear binding Coulombic rather than strong. Above all, the mean density does not gravitate, so the background neither decelerates nor accelerates.

On the cosmological constant: fixed, not absent. In the Newtonian limit of general relativity with Λ the field equation reads $\nabla^2\varphi = 4\pi G\rho - \Lambda c^2$, so Λ enters precisely as a constant subtracted from the source — the same form as (2), with $\Lambda c^2 = 4\pi G\bar{\rho}$. Historically this is no coincidence: Einstein introduced Λ in 1917 with the Neumann–Seeliger difficulty of Newtonian cosmology in view, the same difficulty that forces a prescription for $\bar{\rho}$ here. What distinguishes this model is not the absence of such a term but its *status*: in Λ CDM it is a free parameter whose near-coincidence with today's matter density is unexplained, whereas here it is fixed by vanishing total gravitational charge and tracks the matter as a^{-3} .

A falsifiable prediction: the coasting Hubble diagram. With a vanishing background source $\ddot{a} = 0$ and $a \propto t$, so $H(z) = H_0(1 + z)$ and

$$d_L(z) = \frac{c}{H_0} (1 + z) \ln(1 + z). \quad (6)$$

Since H_0 is degenerate with the supernova absolute magnitude, only the *shape* of $\mu(z)$ is testable. Marginalising the additive constant against Λ CDM ($\Omega_m = 0.315$) leaves a residual tilt of ≈ 0.21 mag peak-to-peak over $0.01 < z < 2.3$ (+0.07 mag at $z = 0.1$, −0.09 mag at $z = 1$), an rms of 0.07 mag. Against a per-supernova scatter of ~ 0.12 mag this is invisible in one object, but modern compilations hold $\sim 10^3$ supernovae and reach a systematic floor of 0.02–0.03 mag, so the prediction sits $5\text{--}8\times$ above the noise and is **already decidable**. Published comparisons of coasting ($R_h = ct$) models have generally favoured Λ CDM, a conclusion contested by Melia [11], and the tilt is concentrated where calibration systematics are largest. If the model is wrong, this is where it should fail.

The obstacles, in order of severity

1. Structure cannot grow in a coasting background — an obstruction, not a gap.

The linear growth equation is $\ddot{\delta} + 2H\dot{\delta} = 4\pi G\bar{\rho}\delta$, the source being the perturbation's own self-gravity, so the mean sets the amplitude even though only the contrast gravitates. In

standard cosmology $\bar{\rho}$ and H are *locked* by Friedmann, $\bar{\rho} = 3H^2/8\pi G$, so source and friction scale alike and $\delta \propto a$: the factor 10^5 that carries 10^{-5} to order unity. Here they are decoupled — $\bar{\rho} \propto a^{-3} \propto t^{-3}$ against $H^2 \propto t^{-2}$ — so the source is subdominant at late times, leaving $\ddot{\delta} + (2/t)\dot{\delta} \approx 0$ with modes $\delta = \text{const}$ and $\delta \propto t^{-1}$: **growth freezes**. Either the mean must gravitate after all, discarding the coasting prediction, or the contrast must be *supplied* at the required amplitude rather than grown — which would make cosmic structure something the seed specifies rather than something gravity builds, and would then have to carry the observed shape of $P(k)$ as well. This is the sharpest obstacle between the present framework and a cosmology.

2. **No account of the dark-matter phenomena.** With no extra gravitating component co-located with galaxies, flat rotation curves and cluster mass-to-light ratios stand where Newtonian dynamics without dark matter leaves them. This is a deficit, not a simplification, and it exceeds the economy gained by dispensing with Λ .
3. **No acceleration:** $q = 0$ (Section 10).
4. **A law of motion for the seed.** $\rho = \nabla^2 \varphi$ is a definition; φ_E has no dynamics here and its spectrum is an input.
5. **Remaining observational fronts:** the CMB acoustic-peak spectrum under any alternative reading, large-scale-structure statistics including the baryon acoustic feature, and light-element abundances beyond helium.

None is addressed here, and item 1 may not be addressable without giving up something the framework asserts. That is why this paper claims an extension to gravitation and not a cosmology.

An eternal universe as the less brutal alternative. If the seed is not an initial condition but a **permanent statistical property** of φ_E , there is no first moment and no origin to account for. Such a universe is *statistically stationary* rather than repeating: laws and ensemble properties fixed, contents forever different, structures forming and dissolving without a preferred epoch. It is not the steady state of Hoyle, Bondi and Gold, which requires continuous creation of matter to hold the density constant. This changes the standing of item 1: with infinite time available, unbounded growth of structure would be inadmissible, so frozen growth becomes closer to a requirement than a defect — though it still does not explain how the observed structure arose. Eternity is not free, however. Tolman’s objection — entropy accumulating across an infinite past — must be met by some recycling mechanism, and none is offered; Olbers’ paradox is eased by expansion and redshift but not dismissed; the observed evolution of source populations with redshift, which told against the steady-state model, must be accommodated. And a coasting expansion has a *finite* age $1/H_0 \approx 14.5$ Gyr, so eternity requires a bounce, cycles, or different early behaviour, not supplied here.

13 Conclusion: two tiers, one two-valued Poisson form

The construction is complete — a perturbation of φ_E quantized into charges, Coulomb building nuclei, atoms and molecules from them, Newtonian gravitation acting on the energy of what is built — and an observation can now be recorded that was used as a premise nowhere above. Each tier was set up on its own terms; that is what makes the observation worth recording.

The two tiers, arrived at independently, turn out to have *one* structure and to differ in a single sign. Each force is the gradient of a potential obeying Poisson’s equation with a two-valued source,

$$\text{electric: } \rho_c = -\varepsilon_0 \nabla^2 \varphi_E, \quad \text{gravitational: } \rho_m = \frac{1}{4\pi G} \nabla^2 \varphi_g, \quad (7)$$

the force being $-(\text{charge}) \nabla \varphi$ in each case. The one structural difference is the relative sign, and with it the force law:

	like sources	unlike sources
electric (charge)	repel	attract
gravitational (mass)	attract	repel

There are thus two independent binary signs — the source $\pm \nabla^2 \varphi$ and the force $\mathbf{F} = \pm \nabla \varphi$ — whose four combinations are exactly the four “charges” the two forces couple to: the repulsive-for-like sign is electromagnetism, its $\pm$ source the $\pm$ charge of the small scale; the attractive-for-like sign is gravitation, its $\pm$ source the $\pm$ contrast of the large. One operator and two signs yield $\{\pm \text{charge}, \pm \text{mass}\}$.

That single sign difference generates the two tiers. Because like charges repel, matter neutralises into $\pm$ pairs and electromagnetism *screens*, staying local and small-scale. Because like masses attract, mass *accumulates* into large-scale structure, and because unlike sources repel, overdense and underdense regions separate into web and voids. The scale separation is thus a *consequence* of the force-sign, not a separate assumption. A second asymmetry — the *direction* in which the Poisson relation is used — fixes the scale of each force. For electromagnetism the potential is primary and charge is obtained by *differentiating*: the Laplacian is local and magnifies short wavelengths ($\rho_k \propto k^2 \varphi_k$), so a small-scale perturbation is inflated into sharp charges. For gravitation the source is primary and the potential is obtained by *integrating*: the inverse Laplacian is global and smooths ($\varphi_k \propto \rho_k / k^2$), building a broad potential. The same relation run forward makes small-scale electromagnetism and run backward large-scale gravitation — and that is why charge is small and gravity large.

This is a *structural parallel*, observed after the fact, and we resist promoting it into a premise. It does not say gravitation is derived from electromagnetism, or that one seed generates both sectors: the gravitational source here is the matter’s own energy density and owes nothing to φ_E beyond the matter it made. Nor does the shared form make the two constants one — whether α and α_G reduce is the open question below, and the parallel is a reason to ask it, not an answer. What can fairly be said is that two forces spanning forty orders of magnitude in strength, introduced for entirely separate reasons — one to make matter, one to gather it — were found to be the same equation read with two signs.

14 Open problems

1. **The proton–electron size difference**, in three parts of unequal status. (i) *The ratio* is read here as geometric: a caged, zero-localisation electron sits at R_p and a free one at a_0 , so $a_0/R_p \approx 1/\alpha^2$, needing no relativity. This the model addresses. (ii) *The absolute scale* of R_p — the same datum as the mass ratio and the nuclear energy scale: one open number, not derived here, nor in the Standard Model. (iii) *The sign asymmetry*: why $+$ is the compact sign at all. Pure Coulomb with the Laplacian is charge-conjugation symmetric, so the asymmetry cannot arise from these ingredients; a natural candidate is a *non-Gaussian (skewed)* seed, whose super- and sub-level sets are not mirror images — one sign forming sharp compact pits, the other broad swells — a geometric route from the *statistics* of the seed rather than a new force.
2. **Could the two potentials be one field?** Split a single potential into scales, $\varphi = \varphi_{\text{short}} + \varphi_{\text{long}}$, reading charge as the short-wavelength Laplacian and mass contrast as the long-wavelength one. The objection that a neutral hydrogen cloud has vanishing charge but non-vanishing mass does not apply, the two roles never being evaluated at the same scale. Two conditions would have to be met: a filter scale fixed by something physical, presumably where matter condenses; and the abundance of small-scale structures tracking $\nabla^2 \varphi_{\text{long}}$, which

for a Gaussian seed fails identically and requires local-type non-Gaussianity — the same skewness wanted in item 1(iii), so one assumption might serve both. This would also be what it takes for the seed to supply the large-scale inhomogeneity that is currently an input. Whether a skewed seed’s small-scale power correlates with its own long-wavelength Laplacian, and with which sign, is a well-posed calculation, not attempted here.

3. **The fluctuation spectrum and a law of motion.** $\rho = \nabla^2 \varphi$ is a definition, not a dynamics: φ has no equation of motion and the seed spectrum is an input. Section 10 shows only that the natural choice is consistent with the observed matter spectrum — a consistency check, not a prediction. Supplying φ with a law of motion is the substantial piece of theory still missing.
4. **The mass ratio $m_p/m_e = 1836$, and why it is not the nuclear scale.** The mass ratio implies a nuclear size $a_0/(m_p/m_e) \approx 29$ fm, whereas the actual scale is $\sim 2\text{--}3$ fm; equivalently the size ratio $a_0/R_p \approx 2.5 \times 10^4 \approx 1/\alpha^2$ exceeds 1836 by a factor ~ 14 . The nucleus is that much more compact than the naive size–mass law allows, because the caged electron is flat, held collectively by several protons. Deriving that factor would make m_p/m_e fix the nuclear scale, hence α , and collapse two numbers into one; asserting it incidental would be numerology.
5. **The neutrino.** β -decay’s continuous spectrum — the 1930 objection that sank the proton–electron neutron [5]. The “electron expands” step needs a home for that energy and spin; an attempt to have a field carry it came out too weak, so the neutrino is neither derived nor eliminated here.
6. **Precision and scope.** Abundances beyond helium, fluctuation-to-structure statistics, and the extreme regimes (strong-field gravity, the highest energies) lie beyond the Coulomb+Newton scope kept here.

The count of constants. Only dimensionless numbers are physical. Stripped to essentials the model carries **three**:

$\alpha = 1/137$	Coulomb strength	sets the size hierarchy, $a_0/R_p \approx \alpha^{-2}$
$m_p/m_e = 1836$	the two charge species	sets the mass hierarchy
$\alpha_G = Gm_p^2/\hbar c \approx 5.9 \times 10^{-39}$	gravitational strength	sets where Newton takes over

The first two are both small-scale: atoms, molecules, chemistry and nuclear binding ratios follow from them parameter-free, gravity playing no part. The third governs the transition between the tiers, and does so at a definite scale: a body can be held up electromagnetically only up to about $\alpha_G^{-3/2} \approx 2 \times 10^{57}$ protons $\approx 1.8 M_\odot$, the stellar mass scale, beyond which gravity wins. The smallness of α_G is therefore not an embarrassment but the very thing that creates the two-tier structure: gravity is so weak that matter can be chemical over an enormous range before Newton asserts itself. Against this, the Standard Model carries some nineteen free parameters and standard cosmology adds several more. Three dimensionless numbers over the domain covered here is a striking reduction — with the caveat that α_G is not derived (it is the hierarchy problem, and no one has derived it), that m_p/m_e and $1/\alpha^2$ differ by the unexplained factor ~ 14 of item 4, and that whether α and G themselves reduce to one is the deeper question this deliberately parallel construction does not settle.

References

- [1] C. Johnson, *Real Quantum Mechanics* (Body&Soul, Vol. VI), ~ 340 pp — the comprehensive treatise, of which RealQM, RealNucleus, the inertia-as-energy analysis and the present cosmology are chapters. Free download from the GitHub gallery: <https://claes542.github.io>.

[io/RealMolecule/realquantum3.pdf](https://claes542.github.io/RealMolecule/realquantum3.pdf) (interactive gallery: <https://claes542.github.io/RealMolecule/>).

- [2] C. Johnson, *RealQM: Quantum Mechanics as Multiphase 3D Continuum Mechanics*, in [1].
- [3] C. Johnson, *RealNucleus: nuclear binding from a pure-Coulomb proton–electron model*, in [1].
- [4] C. Johnson, *Energy Without Mass: RealQM, Inertia, and the Penning Trap*, in [1].
- [5] W. Pauli, open letter to the Tübingen group (1930), proposing the neutrino to save energy conservation in β -decay.
- [6] J. S. Farnes, *A unifying theory of dark energy and dark matter: Negative masses and matter creation within a modified Λ CDM framework*, *Astronomy & Astrophysics* **620**, A92 (2018); arXiv:1712.07962.
- [7] Ya. B. Zel’dovich, *Gravitational instability: an approximate theory for large density perturbations*, *Astronomy & Astrophysics* **5**, 84 (1970); see also S. F. Shandarin and Ya. B. Zel’dovich, *Reviews of Modern Physics* **61**, 185 (1989).
- [8] O. Hahn, C. M. Porciani, C. M. Carollo and A. Dekel, *Properties of dark matter haloes in clusters, filaments, sheets and voids*, *Monthly Notices of the Royal Astronomical Society* **375**, 489 (2007); J. E. Forero-Romero et al., *MNRAS* **396**, 1815 (2009) — the tidal-tensor classification.
- [9] J.-P. Petit and G. d’Agostini, *Negative mass hypothesis in cosmology and the nature of dark energy*, *Astrophysics and Space Science* **354**, 611 (2014) — the Janus bimetric model.
- [10] S. Hossenfelder, *A bi-metric theory with exchange symmetry*, *Physical Review D* **78**, 044015 (2008) — negative gravitational mass without the classical runaway.
- [11] F. Melia, *The $R_h = ct$ universe*, *Monthly Notices of the Royal Astronomical Society* **419**, 2579 (2012); see also the subsequent literature contesting its fit to the supernova Hubble diagram.
- [12] J. W. Cahn and J. E. Hilliard, *Free energy of a nonuniform system. I. Interfacial free energy*, *J. Chem. Phys.* **28**, 258 (1958); S. M. Allen and J. W. Cahn, *Acta Metall.* **27**, 1085 (1979).
- [13] T. Ohta and K. Kawasaki, *Equilibrium morphology of block copolymer melts*, *Macromolecules* **19**, 2621 (1986) — the Cahn–Hilliard functional with a long-range interaction.