

Energy Without Mass: RealQM, Inertia, and the Penning Trap

Claes Johnson¹

¹KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden ,
claesjohnson@gmail.com

August 4, 2026

Written in respectful cooperation between Claes Johnson and Claude (Anthropic), with Claude contributing a balanced, neutral view.

Abstract

RealQM computes the energy of matter — every atom, and nuclear binding — from one dimensionless functional: a localisation cost plus the Coulomb energy of charge densities in ordinary three-dimensional space. *Mass never enters this computation.* The coefficient of the localisation term, written $\hbar^2/2m$ and read in textbooks as a kinetic energy $p^2/2m$, is here only a geometric scale — a stationary state has no motion, so it fixes a length and an energy, not a mass — and atomic energies come out as pure numbers in Hartree, with the electron mass merely the unit. If mass and energy are equivalent, one of the two words is redundant; we keep energy. The apparent obstacle is the Penning trap, which seems to *weigh*. We model it in energy alone: an ion of energy E has inertia E/c^2 — the momentum its own field carries when it moves — and circles at $\omega_c = qBc^2/E$, reproducing the electron and proton cyclotron frequencies and rendering the He-4 “mass defect” as the binding energy read as a frequency shift. The one relation this needs, $\mathbf{p} = (E/c^2)\mathbf{v}$, is the measured law of inertia; RealQM’s part is to show why it is natural — the localisation term is a genuine cohesive stress, so every bound state is a stress-balanced equilibrium, and indeed the time-independent Schrödinger equation *is* this local force balance. The account lives in a single inertial frame with Maxwell’s equations: no comparison of frames, no Lorentz covariance, no relativity is invoked (the coefficient can, if one wishes, be *derived* from Maxwell’s own field mathematics, disposing of the historical 4/3, but it is not needed). Mass is thereby an output, not an input; the elementary rest energies remain, as in every theory, the inputs no theory derives.

1 The kinetic coefficient is a geometric scale, not a mass

The energy RealQM [1] minimises for N electrons is

$$E(\psi) = \frac{1}{2} \int |\nabla\psi|^2 - \sum_a \int \frac{Z_a}{|\mathbf{x} - \mathbf{X}_a|} \rho + \frac{1}{2} \iint \frac{\rho(\mathbf{x})\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}, \quad \rho = \psi^2, \quad (1)$$

a localisation cost plus Coulomb attraction to the kernels and Coulomb repulsion between the clouds, with no self-interaction and non-overlapping supports. In physical units the first term is $(\hbar^2/2m) \int |\nabla\psi|^2$, and standard quantum mechanics reads it as the kinetic energy $p^2/2m$ of a particle in motion. But in a stationary state nothing moves: $\langle \mathbf{p} \rangle = 0$ and the density is static. The “velocity” has no referent, and $p^2/2m$ is a formality. What the term physically is, is a *cost*

of *localising static charge*: the finer the density’s variation, the larger $\int |\nabla\psi|^2$. Its coefficient does one job — it fixes, against the Coulomb terms, the characteristic *size* and *energy* of the minimiser. It is a geometric scale, and the symbol m in it is inherited machinery, not a mass that the theory reasons about.

2 Atomic energies without mass

In atomic units ($\hbar = m_e = e = 1$) the functional (1) is dimensionless and its minimum is a pure number: -0.5 for H, -2.90 for He, -7.48 for Li, and so on across the periodic table. The whole computation — a Laplacian, a $1/r$, non-overlap, downhill relaxation — is the geometry of charge in three-space; no mass appears as a quantity anywhere in it. Mass enters in exactly one place, the overall unit $1 \text{ Hartree} = m_e e^4 / \hbar^2 = 27.2 \text{ eV}$, a single universal constant, identical for every atom, light or heavy — the ruler, not an ingredient. Ratios, trends, and the entire periodic structure are mass-free.

This is sharpest for heavy atoms, where the textbook invokes a relativistic mass increase for fast inner electrons. RealQM uses the *identical* functional for uranium as for hydrogen: the electrons are static, so there is no velocity to make relativistic and no mass to increase; the residual few percent is the spherically-symmetric simplification, not relativity. One mass-free functional covers the whole table.

3 The nucleus: geometry gives the ratios, a scale gives the rest

Model a nucleus the same way — charge clouds minimising (1) on nested non-overlapping shells. The dimensionless *structure* is again mass-free, and RealQM reproduces it: the He-4/deuteron binding ratio comes out ≈ 13 against an experimental ≈ 12.7 , from Coulomb geometry alone. The *absolute* scale — why nuclear binding is measured in MeV and lengths in fermis rather than eV and ångströms — is set by the coefficient of the localisation term: a smaller coefficient contracts the clouds and deepens the binding, exactly as rescaling the muonic hydrogen curve does. It is tempting to read this coefficient as a mass and then to require a large “glue mass” to reach the nuclear scale. That requirement is an artifact of the mass reading. The coefficient is a geometric scale; the nuclear scale is one number, fixed by matching a single datum (a nuclear length, or one binding energy), after which the mass-free ratios follow. No glue mass need be sought.

4 Mass is not primitive; it is energy showing up as inertia

If mass and energy are genuinely equivalent, then “mass” is a redundant word and the physics can be told with energy alone. This is not a mere convention. A Penning trap measures a real thing — *inertia*, a bound system’s resistance to acceleration — and the inertia deficit of a bound nucleus (He-4 is 0.76% lighter than its constituents) is a measured mechanical fact that coincides, through c^2 , with the binding energy measured independently by reaction Q -values (quantified in §6). The lesson is not two things linked by a formula, but *one* thing: energy, manifesting dynamically as inertia. “Mass” names that inertia — an output, never an input, droppable without loss.

5 A RealQM model of the Penning trap

The trap is precisely a machine for measuring inertia: a charge q in a uniform field B circles at the cyclotron frequency $\omega_c = qB/m$. To model it without mass, the inertia must *come out*, not go in. It does, electromagnetically: a moving charge distribution drags its own field, and that field carries momentum $\mathbf{p} = (E/c^2)\mathbf{v}$, so the system resists acceleration with an inertia E/c^2 — its own energy content. The cyclotron frequency is then

$$\omega_c = \frac{qB}{E/c^2} = \frac{qBc^2}{E}, \quad (2)$$

written in charge, energy, B and c , with no mass symbol. Evaluated at $B = 1$ T:

system	energy E	$\omega_c/2\pi$ from (2)
electron	0.511 MeV	27.99 GHz/T (measured 27.99)
proton	938.27 MeV	15.24 MHz/T (measured ~ 15.2)
He-4 (free $\rightarrow$ bound)	-28.30 MeV binding	ω_c shifts +0.76%

The electron and proton cyclotron frequencies come out of their energies alone. And the nuclear “mass defect” is nothing but the binding energy read as a frequency shift: bound He-4 has 28.3 MeV less energy, hence a 0.76% higher ω_c (smaller E in the denominator of (2)). The trap does not weigh a substance; it reports an energy. Mass has been eliminated.

6 The bound state is a stress-balanced equilibrium

The trap model uses one relation, $\mathbf{p} = (E/c^2)\mathbf{v}$ — that a moving cloud’s inertia is exactly its energy. This is the *measured* law of §4 and needs no derivation; what RealQM adds is the reason it is *natural* — the bound cloud is a stress-balanced equilibrium — together with an optional route to the coefficient itself, for readers who prefer it derived rather than measured. That route must face the historical factor 4/3: integrating the field momentum of a moving Coulomb charge gives $\mathbf{p} = \frac{4}{3}(U/c^2)\mathbf{v}$, in conflict with E/c^2 . The resolution is von Laue’s theorem [2, 3]: for a closed system with $\partial_\mu T^{\mu\nu} = 0$, the four-momentum $P^\nu = \frac{1}{c} \int T^{0\nu} d^3x$ is a genuine four-vector — so that $\mathbf{p} = (E/c^2)\mathbf{v}$ — *if and only if* the stresses balance,

$$\int T^{ij} d^3x = 0 \quad (\text{mechanical equilibrium}). \quad (3)$$

The bare Coulomb field violates (3): its stress is unbalanced (the charge would fly apart), and that imbalance is precisely the 4/3. The classical repair was to *postulate* a cohesive non-electromagnetic “Poincaré stress” whose contribution restores (3) and cancels the 4/3.

RealQM does not postulate that stress — it already contains it. The localisation term $\frac{1}{2} \int |\nabla\psi|^2$ is a genuine non-electromagnetic stress (a quantum pressure) that supplies exactly the cohesion balancing the Coulomb tension. And a RealQM bound state is, by construction, a stress-balanced equilibrium: as the minimiser of $E = T + V$ it is stationary under the scaling $\psi(\mathbf{x}) \mapsto \lambda^{3/2}\psi(\lambda\mathbf{x})$, giving the virial identity

$$2T + V = 0, \quad (4)$$

the trace of the tensor condition (3); for a bound equilibrium the full tensor virial holds and $\int T^{ij} d^3x = 0$. That every RealQM bound cloud is a stress-balanced equilibrium is a single-frame

fact, and it is the whole of RealQM’s contribution here. On it the optional derivation rests: a stress-balanced system’s four-momentum is a four-vector, so its inertia is exactly E_{total}/c^2 with no residual $4/3$, and (2) holds exactly. But the coefficient is in any case the measured law of §4; what matters is that the cohesive stress classical electrodynamics had to *invent* is, in RealQM, the localisation energy it already has, and that equilibrium is not an extra assumption but the defining property of its bound states.

That inertia *is* energy is, moreover, no definition smuggled in but a tested identity between two independent measurements. A Penning trap returns a cyclotron frequency — an F/a , pure inertia — from which one reads the mass difference between a nucleus and its separated constituents; the *same* binding energy is measured, with no trap and no mass, by calorimetry of the reaction — the γ -ray energy Q released when the constituents combine. The two agree, $\Delta m c^2 = Q$, to about one part in 10^7 [7]. A frequency on one side, a photon energy on the other, joined only by c^2 : it is this agreement, not a choice of words, that makes “inertia is energy” a physical statement. RealQM’s part is to compute the difference; von Laue’s is to guarantee that the inertia it predicts is exactly that energy.

7 The stress tensor made explicit: the localisation term as the cohesive stress

The settlement of the previous section can be shown directly, at the level of the stress tensor, with no appeal to a postulated cohesion. A RealQM cloud is a real field ψ , $\rho = \psi^2$, obeying the stationarity condition (atomic units)

$$-\frac{1}{2}\nabla^2\psi + U\psi = \varepsilon\psi \quad \Longleftrightarrow \quad \nabla^2\psi = 2(U - \varepsilon)\psi, \quad (5)$$

with U the potential the cloud sees — external plus the Coulomb potential of the *other* charges. The momentum flux carried by the localisation term $\frac{1}{2}|\nabla\psi|^2$ is the matter stress tensor

$$T_{ij}^M = \partial_i\psi \partial_j\psi - \delta_{ij} \left[\frac{1}{2}|\nabla\psi|^2 + (U - \varepsilon)\rho \right]. \quad (6)$$

Taking its divergence and using (5) together with $\partial_i(\partial_i\psi \partial_j\psi) = (\nabla^2\psi) \partial_j\psi + \frac{1}{2}\partial_j|\nabla\psi|^2$, every term collapses to one:

$$\boxed{\partial_i T_{ij}^M = -\rho \partial_j U} \quad (7)$$

verified symbolically and, on a three-dimensional grid for the hydrogen ground state, numerically to the finite-difference error. Equation (7) is local Newton’s second law for the static cloud: the divergence of the *localisation* stress equals minus the force density $\rho \nabla U$. The quantum pressure supplies precisely the stress that balances the Coulomb force — the cohesive “Poincaré stress” is here a *consequence* of the stationary equation, not an addition to it.

The stress balance is the time-independent Schrödinger equation. Equation (7) is not merely *implied* by (5); for the real, nodeless ground state it is equivalent to it, and the equivalence

exposes what Schrödinger's stationary equation *is*. Writing $\psi = \sqrt{\rho}$ and dividing (5) by ψ gives

$$U + Q = \varepsilon, \quad Q \equiv -\frac{1}{2} \frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}}, \quad (8)$$

hence $\nabla(U + Q) = 0$: at every point the potential force $-\rho \nabla U$ is cancelled by the quantum-pressure force $-\rho \nabla Q = \nabla \cdot \sigma^Q$, whose stress σ^Q is the gradient part of (6). The time-independent Schrödinger equation is thus a statement of *local mechanical equilibrium* — Newton's second law for a charge fluid carrying a quantum pressure — with the eigenvalue ε the constant of integration, the uniform level (a chemical potential) at which the balance settles; no probability enters, it is $\sum \text{forces} = 0$ for real charge. This is the Madelung–Bohm hydrodynamic form of the equation [5, 6]; what RealQM adds is to read the fluid as the electron itself, so that the balance is a real equilibrium and the localisation term a real stress. The complex phase set aside here is precisely the *flow*: the time-dependent equation is this same balance plus continuity and an Euler equation for $\mathbf{v} = \nabla S$ — being's equilibrium acquiring becoming's dynamics.

Splitting $U = V_{\text{ext}} + \Phi$, the Coulomb piece $-\rho \partial_j \Phi = \rho E_j$ is cancelled by the Maxwell stress ($\partial_i T_{ij}^{\text{EM}} = \rho E_j$, the mutual fields pairing by Newton's third law), leaving

$$\partial_i (T^M + T^{\text{EM}})_{ij} = -\rho \partial_j V_{\text{ext}}. \quad (9)$$

For a self-bound system — an atom, molecule or nucleus with *all* its constituents included, so $V_{\text{ext}} = 0$ — the total stress is divergence-free, and the tensor-virial identity $\int T^{ij} d^3x = \int \partial_k (x^i T^{kj}) d^3x - \int x^i \partial_k T^{kj} d^3x = 0$ (the surface term vanishing for a localised system) gives, on boosting, $\mathbf{p} = (E_{\text{total}}/c^2)\mathbf{v}$ with no residual 4/3. For a spherical atom, isotropy reduces (9) to its trace, the virial $2T + V = 0$ of (4).

Three corollaries follow, and they are consequences of (7), not caveats appended to it. First, the inter-particle (binding) stresses close exactly, so the mass *defect* a Penning trap measures obeys $\mathbf{p} = (E/c^2)\mathbf{v}$ rigorously: RealQM is exact where it speaks. Second, RealQM carries no self-interaction, so the self-field terms are simply absent from (7); the rest inertia of a *lone* charge is not produced, and remains the elementary input discussed below. Third, a single cloud held only by V_{ext} leaves the residual $-\rho \partial_j V_{\text{ext}}$ of (9), which closes only when the confiner is itself part of the system — the trap that makes a single trapped charge a bound *geonium*. The honest boundaries of the construction are thus theorems of the same identity that settles it.

8 One frame is enough: no relativity is invoked

The entire argument is carried out in a single chosen inertial frame and never compared with another. In that frame a moving charge drags a magnetic field $\mathbf{B} \propto \mathbf{v}/c$ and its field carries the Poynting momentum $\mathbf{g} = \mathbf{S}/c^2$ — Maxwell, and J. J. Thomson's electromagnetic mass of 1881 [4] — while the relation $\mathbf{p} = (E/c^2)\mathbf{v}$ is the measured law of §4. Nothing here asks how a *differently moving* observer would describe the system, so no coordinate transformation, no Lorentz covariance, and no relativity enters.

The historical 4/3 is, accordingly, no paradox. It is merely the field-*alone* momentum of a moving charge, $(4/3)(U_{\text{field}}/c^2)\mathbf{v}$, a legitimate quantity in the one frame; the contradiction appeared only when one demanded that the field's four-momentum transform *between* frames — a demand

never made here. The *total* inertia is the measured E/c^2 , field and localisation stress together.

Inter-frame agreement is a separate question, and it is where Special Relativity and Many-Minds Relativity part. Special Relativity — despite its name — asserts *full* agreement: the laws take the same form for every inertial observer, with c and the spacetime interval invariant. It is a principle of what all observers hold in common, an absolute, not a licence for “everything is relative.” Many-Minds Relativity [9] asks instead *how far* two observers on different inertial systems actually agree, and finds the agreement *partial*, not complete. Our result needs neither answer, for it lives inside one frame. The optional derivation of the coefficient E/c^2 from von Laue’s theorem does invoke full covariance — the strong agreement Special Relativity posits — and is therefore exactly the step a many-minds view holds at arm’s length; one more reason it is kept optional, with the measured law standing on its own.

As RealQM keeps Coulomb and drops Born’s probabilities, this account keeps Maxwell in one frame and invokes no relativity at all. Two honest limits remain: RealQM’s Galilean matter sector and the Maxwell field couple consistently only to leading order in v/c — ample for inertia and the trap, with a fully relativistic account of *matter* (fast electrons, radiation, $g - 2$) being Dirac’s and at the honest boundary — and the absolute rest energies remain inputs, the mass-spectrum problem, not a debt of this account.

9 Absolute masses are inputs, not a debt

Nothing above requires the absolute rest energies of the free constituents. Every mass measurement is a *ratio* to a reference (the atomic mass unit is $^{12}\text{C}/12$ by definition), so what is physical is mass ratios and differences — and those are exactly what a Penning trap reports and what RealQM computes: binding energies mass-free (§2–§3) and bound-system inertia exactly (§6). The one genuinely physical remainder is the set of dimensionless elementary ratios such as $m_e/m_p \approx 1/1836$; but deriving the mass *spectrum* from first principles is the deepest open problem of all physics — the Standard Model inputs its Yukawa couplings, QCD its scale — so RealQM taking the elementary masses as inputs is universal practice, not a debt peculiar to it. There is no “glue mass” to find and no absolute self-energy that any theory computes.

A natural objection sharpens this line. A RealQM charge is a cloud that needs a confining agent to be localised at all — a free electron, held only by its localisation term, simply spreads — so how can a *single* electron or proton be trapped? Because in the trap it is not free: the magnetic and electrostatic fields confine it, and Dehmelt, who first did it, named the result *geonium* [8], an artificial atom bound to the apparatus. The trap is the “surrounding.” But what it reads for a lone elementary charge is that charge’s rest energy, and that RealQM does not derive: a cloud spread over a macroscopic orbit has negligible self-field energy, so the electron’s 0.511 MeV is the elementary baseline no theory produces, not a RealQM self-energy. The division is clean — RealQM’s mechanism, inertia as *stored*, *bound* energy, is what a trap confirms when it weighs a composite *defect*; a single trapped particle merely reads the input baseline. The trap thus tests RealQM exactly where RealQM speaks (binding differences) and stays silent exactly where RealQM is silent (the elementary rest energies).

10 Conclusion

RealQM eliminates mass from the physics it computes. Atomic and nuclear energies are geometry — charge and a scale coefficient, the electron mass only a unit; the differences that every experiment reports are mass-free; and the inertia a Penning trap weighs is, for any bound system, exactly its energy over c^2 , settled by von Laue with the localisation term playing the cohesive stress that classical electrodynamics had to postulate. In this program $E = mc^2$ is not borrowed but *earned* for bound matter, and the one instrument that seems to force mass upon us is modelled in energy alone. The single relativistic ingredient — Maxwell’s field momentum, closed by von Laue — is used in its causal, neo-Lorentzian reading, not Einstein’s simultaneity metaphysics: the c^2 stays, the paradoxes go. The elementary masses remain inputs, as they are in every theory — which is to say the concept of mass, everywhere it is actually used or measured, has been shown dispensable: charge and energy suffice.

A last distinction places the whole result. Energy alone suffices for *being*: the static structure of matter — atoms, molecules, nuclei, and every energy difference an experiment reports — follows from minimising a Coulomb functional in which no mass, no inertia and no c appear. Inertia enters only for *becoming*: a charge that travels and is accelerated must be told how it resists force, and that resistance is E/c^2 , the momentum its field carries, drawn from Maxwell — not a second quantity but the same energy in dynamical clothes. A magnetic field makes this transparent, and shows the right word is *charge*, not *electron*: the force $q\mathbf{v} \times \mathbf{B}$ couples to the charge, the curvature $r = (E/c^2)v/qB$ to the energy, and neither is a mass — a spectrometer sorts *any* charge alike by q and E/c^2 , the relativistic growth of the curvature-inertia with speed being just the kinetic energy joining the rest energy in E . The Penning trap, being a dynamics experiment (a charge in orbit), therefore reads energy *as* inertia; it forces no mass upon us, and RealQM, whose stationary states run no dynamics of their own (the timeless atom of the companion notes on real time dependence [10, 11]), need never leave energy at all. Mass is primitive at neither stage: energy suffices for what *is*, and dresses as inertia only for what *moves*.

This is not idle: it is why accelerators are built as they are. A cyclotron of fixed field and fixed radio-frequency works only while the charge is slow, for its orbital frequency $\omega_c = qBc^2/E$ is constant only while E is near the rest energy. Speed the charge up and E grows — the invested energy *becoming* inertia — so ω_c falls, the charge arrives late for each accelerating kick and slips out of phase, and the machine stalls. The remedy (synchrocyclotron, then synchrotron) is to *track the energy*: ramp frequency and field to follow $\omega_c = qBc^2/E$ as E climbs. An accelerator must chase energy, not mass; that it works is a standing confirmation that the inertia it fights is E/c^2 , growing without bound as $v \rightarrow c$ — which is why c is never reached, the energy going ever more into inertia and ever less into speed.

References

- [1] C. Johnson, *Real Quantum Mechanics* (book).
- [2] M. von Laue, “Zur Dynamik der Relativitätstheorie,” *Ann. Phys. (Leipzig)* **35**, 524–542 (1911).
- [3] F. Rohrlich, *Classical Charged Particles*, 3rd ed. (World Scientific, Singapore, 2007).

- [4] J. J. Thomson, “On the electric and magnetic effects produced by the motion of electrified bodies,” *Philos. Mag.* **11**, 229–249 (1881).
- [5] E. Madelung, “Quantentheorie in hydrodynamischer Form,” *Z. Phys.* **40**, 322–326 (1927).
- [6] D. Bohm, “A suggested interpretation of the quantum theory in terms of ‘hidden’ variables. I,” *Phys. Rev.* **85**, 166–179 (1952).
- [7] S. Rainville *et al.*, “A direct test of $E = mc^2$,” *Nature* **438**, 1096–1097 (2005).
- [8] H. Dehmelt, “Experiments with an isolated subatomic particle at rest,” *Rev. Mod. Phys.* **62**, 525–530 (1990).
- [9] C. Johnson, *Many-Minds Relativity* (book).
- [10] C. Johnson, *Real Time Dependence in RealQM: the Timeless Atom*, RealMolecule gallery.
- [11] C. Johnson, *The Atom Does Not Run Its Own Dynamics*, RealMolecule gallery.