

Constructive Solution of the Clay Navier–Stokes Problem

Claes Johnson

KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden

`claesjohnson@gmail.com`

October 2, 2026

written up with AI assistance (Claude, Anthropic)

Abstract

The solution of alternative (C) of the Clay Navier–Stokes problem announced in September 2026 represents, beyond its result, a new form of mathematics: one performed by AI as a digital optimisation process, fitting a large set of network parameters to given data by forward and backward propagation. Mathematics done this way constructs an object rather than asserting that one exists, and AI in this sense may be seen as a form of automated thinking. We describe a far more restricted version of the same thing, developed over a long period and implemented in the FEniCS software: the automated solution of differential equations, ASDE, a self-learning computational algorithm with dual-based output error control directed at minimal computational effort, whose backward pass is the adjoint and is the same computation as backpropagation. We then present what ASDE delivers when directed at alternative (A): constructive existence of turbulent solutions of the incompressible Navier–Stokes equations at high Reynolds number, for data given in advance, together with qualitative uniqueness of their mean-value outputs, drawn quantitatively by computed stability factors so that a functional can fail the test. We do not claim a discharge of (A) as that alternative is formulated, since (A) asks for smoothness and the solutions that occur are not smooth: their gradients grow like $\nu^{-1/2}$ while their velocities stay bounded, a Hölder-1/3 field neither differentiable nor unbounded. We argue that this third case is what the equations produce, and close with a scaling test which shows that the singular solution constructed for (C) vanishes in the limit of small viscosity in which fluid mechanics lies: held to a fixed admissible forcing class its amplitude, support and lifetime all tend to zero, and held at fixed amplitude its forcing diverges like ν^{-1} and leaves the class.

1 A new form of mathematics, and a restricted version of it

On 8 September 2026 OpenAI presented a solution of alternative (C) of the Navier–Stokes Millennium problem [32]: a proof that a singular solution exists, obtained with ten thousand agents and of the order of 3×10^{11} output tokens [33].

An AI is a digital optimisation process: a very large set of network parameters is fitted to given data by forward and backward propagation, the backward pass carrying the sensitivity of an output to everything that fed it. Mathematics done this way does not establish that an object exists and leave it there; it *constructs* one, by fitting. AI in this sense is a form of *automated thinking*, and the (C) solution is an instance: the singular velocity and pressure fields are built first, to have the required growth, and the force is then defined as what is left over — the residual of the construction, arranged to be smooth.

Over a long period, together with students and colleagues, we have developed a far more restricted version of the same thing: the *automated solution of differential equations*, which we will call ASDE. The project was founded in 2003 with the aim of automated computational mathematical modelling [24]; the programme was set out in Anders Logg’s thesis of the following year, *Automation of Computational Mathematical Modeling* [25], and the phrase is his [26]. It is realised in the FEniCS software [27, 28, 20, 21, 22, 23], whose standard reference is titled for it — *Automated Solution of Differential Equations by the Finite Element Method* [27] — and which is now widely used. ASDE fits a *digital* solution to a differential equation rather than a network to data, and it does so by the same forward and backward passes. It is a small, exactly specified corner of automated thinking, and it has one advantage over the general case: what it constructs comes with a bound on the error in whatever is asked of it.

What follows is a constructive solution directed to alternative (A). We do not claim it is a solution of (A) as that alternative is formulated, and the difference should be stated at the outset rather than discovered later. (A) asks, for *all* admissible data and with $f \equiv 0$, for a solution that exists and is *smooth* for all time. What ASDE delivers, for data given in advance, is *constructive existence* and *qualitative uniqueness* of *turbulent* solutions in the case of high Reynolds number — that is, of small viscosity ν , with the velocity scale and the size of the domain normalised to unity so that ν is the one parameter left and the limit $\nu \rightarrow 0$ can be taken at all — and in *bounded* domains of space, which is where drag, lift and the other outputs of interest are produced. It does not deliver smoothness, because the solutions that occur do not have it — their gradients grow like $\nu^{-1/2}$ while their velocities stay bounded (Section 4).

So this is offered as an answer to what (A) is *about* — whether the equations determine a flow, and what of it they determine — and not as a discharge of (A) as written. Whether the two coincide is for the reader to judge, and it is the one judgement we would ask for.

The Clay Navier–Stokes problem was first addressed on these lines in Chapter 14 of *Computational Turbulent Incompressible Flow*, a book by Johan Hoffman and Claes Johnson published by Springer in 2007 as volume 4 of *Applied Mathematics: Body and Soul* [38]. What follows recalls that chapter and brings it up to date. What is new is the occasion.

2 Basics of ASDE

ASDE is a self-learning computational algorithm with dual-based output error control, directed at minimal computational effort. Its three parts are the following.

A discretisation that fits the equation. Given a differential equation, a domain and data, a finite element method produces a digital solution U whose *residual* $R(U)$ measures by how much it fails to satisfy the equation. Here the method is General Galerkin, G2: continuous piecewise linear in space and time, cG(1)cG(1), with least-squares control of the residual [15]. Nothing is supplied by the user beyond equation, domain and data — no mesh chosen by hand, no turbulence model, no closure, no constant fitted to measurement. This matters for what follows: a computation carrying a fitted turbulence model could not be used to say what the Navier–Stokes equations contain, because the turbulence would have been put in by hand.

A backward pass that converts residual into error. The error in an output $M(u) = \int_Q u \cdot \psi$ is, to leading order, the residual tested against a *dual weight function* φ ,

$$M(u) - M(U) \approx \int_0^T \int_{\Omega} R(U) \cdot \varphi \, dx \, dt, \quad |M(u) - M(U)| \leq S \|R(U)\|_{-1}, \quad (1)$$

with S the stability factor. And φ is obtained by solving the equation linearised about the computed solution *backward in time* from the output functional as data at $t = T$. This is the same object, and the same computation, as the backward pass in the training of a neural network: the adjoint of a forward evolution, propagating sensitivity from an output to everything that fed it. ASDE and AI are doing the same linear algebra in opposite directions through the same trajectory, and that is the sense in which ASDE is the restricted case.

Self-learning: adaptivity towards minimal effort. The dual-based form of this is set out for differential equations generally in [30] and for parabolic problems in [29]. The mesh is neither fixed in advance nor refined uniformly. Refinement goes where the product of residual and dual weight is largest, and stops when the resulting bound meets the tolerance asked. Adaptivity and error control are one mechanism, not two: the computation that says how large the error is also says where to refine to reduce it, and the result is the least computational work that achieves the stated accuracy. The algorithm learns the mesh its own output requires.

What ASDE therefore produces. Not an assertion that a solution exists, but an object: the mesh with its cell count and smallest cell, the discretisation carrying the solution, the residual measured in the norm the bound uses, a computed stability factor for each output, and for each output a value with a bound on its error. That is the solution *de facto*. The record of the method and its computations is in [38], with the residual-based dissipation in [18], the parameter-free formulation in [19], the earlier statement in [31], and two results the same computations gave on the way: the resolution of d'Alembert's paradox [39] and a theory of flight [40]. Further material is collected under the label *clay problem* [45].

Where it now stands. The development since 2007 has gone to compressible flow and to aeronautical Reynolds numbers, carried substantially by Johan Jansson, who has a leading role in the AIAA CFD High Lift Prediction Workshop (HLPW-5 and HLPW-6), where computational predictions of high-lift configurations are submitted blind and compared against wind-tunnel measurement. That is the hardest ordinary test there is for a turbulence prediction: separated flow at aeronautical Reynolds number, outputs that are exactly the mean-value functionals treated below, and a comparison made before the measurement is known. The same method had earlier been submitted to the high-order CFD benchmark workshops, on the tandem-sphere case and on workshop-provided meshes [43]. ASDE is tested in the ordinary way, against experiment, by people with no stake in this article.

The method has also left the laboratory. It is in industrial use for the design of aircraft, wind turbines and ships under the name Adaptive Euler predictive simulation, with FEniCS as a core component. Simulations of three cases were presented at IFASD 2026 [44]: a validation against the benchmarking data of NASA's High Lift Prediction Workshop 4, and — both described there as preliminary — tail buffet on the Saab Open Aircraft Research 2 configuration and a fluid–

structure interaction computation of transonic buffet on the NASA AePW8/DPW4 full-aircraft benchmark.

These are worth more here than a demonstration of software. They are the computation this article offers as a solution, run on geometries nobody chose for being tractable, and judged against measurement by people who did not make the prediction — which is the only test of qualitative uniqueness that matters. A mean-value output either comes out right or it does not, and a benchmark workshop settles it without reference to any argument made above.

3 ASDE applied to Navier–Stokes

The problem is the incompressible Navier–Stokes equations

$$\dot{u} + (u \cdot \nabla)u - \nu \Delta u + \nabla p = f, \quad \nabla \cdot u = 0, \quad (2)$$

on a domain $\Omega \subset \mathbb{R}^3$ with initial data u_0 , forcing f and boundary conditions, at small viscosity, that is at high Reynolds number. The official statement, as set out by Charles Fefferman [36], poses them instead on $\mathbb{R}^3$ or a periodic box with the viscosity fixed, and asks whether solutions remain smooth for all time. Applied to it, ASDE delivers the following two things.

Constructive existence of turbulent solutions. A G2 computation of (2) started from smooth potential data becomes turbulent and stays bounded. The stabilisation supplies dissipation in proportion to the residual, so the dissipation is generated by the computation rather than prescribed by a model; in the vanishing-viscosity limit the same scheme is an Euler solver whose turbulent dissipation appears automatically. Existence here is not an assertion about a solution set being non-empty: a solution is exhibited, for data given in advance, with the error in each output bounded by (1). Ask a non-constructive existence theorem for the drag and there is nothing there to ask; ask a computed solution and it answers with a number and a bound.

There is more to this than error control. A computed solution is an elaborate explicit construction of a velocity–pressure field — millions of numbers on a spacetime mesh, every one available — and such an object can be taken apart: where vorticity concentrates and in what shape, how the velocity gradient decomposes at a point, where energy passes from large scales to small, where the dissipation actually sits as against where a model would put it. A solution concept whose objects can be inspected generates conjectures; one whose objects cannot be exhibited does not. The triple decomposition of the velocity gradient [16], and the energy stability analysis built on it [17], came out of looking closely at resolved velocity fields, and at measured ones [8].

Qualitative uniqueness. Classical uniqueness asks that the data determine the field pointwise. At high Reynolds number that is false, and not marginally: two computations of the same flow on independently generated meshes give velocity fields differing by order one at any chosen point, and continuing to differ under refinement. What the data *do* determine is the qualitative content, and this is what the two computations are found to share: the drag agrees to a few per cent [41] and the lift likewise; the separation line sits in the same place; the wake is organised the same way and sheds at the same frequency; the total dissipation rate is the same.

Qualitative uniqueness is the statement that this is so, with the line between the two lists drawn *quantitatively* rather than by assertion — and that last clause is what keeps it from being a soft version of uniqueness. The dual solution attaches to each functional a computed stability factor: where it is moderate the output is determined by the data to a stated precision, and refinement where the dual weight is large will improve it; where the factor is large the output is not determined, and no refinement will make it so. The claim is falsifiable one functional at a time, and a functional can fail it. That last case is not a failure of the method but an answer: the equations do not determine that quantity, and no computation should be believed which claims otherwise.

Turbulence is a necessity, not a possibility. High Reynolds number flow starting out laminar develops shear instabilities with sharp velocity gradients, which are eventually damped by viscosity in turbulent flow. All high Reynolds number laminar flow is thus unstable and cannot continue to exist over time; it turns turbulent in order to continue to exist. What ASDE constructs is therefore not one possibility among several but what the equations must do.

4 Why smoothness is not what the solutions have

Since (A) as written asks for smoothness, it is worth being precise about what the computed solutions are instead, because they are neither smooth nor singular. Dissipation at a rate independent of the viscosity gives gradients $|\nabla u| \sim \nu^{-1/2}$, and the scales at which that gradient is realised follow from the same constant:

$$\eta \sim \left(\frac{\nu^3}{\varepsilon}\right)^{1/4} \sim \nu^{3/4}, \quad \delta u(\eta) \sim (\varepsilon \eta)^{1/3} \sim \nu^{1/4}, \quad \frac{\delta u(\eta)}{\eta} \sim \nu^{-1/2}, \quad (3)$$

the first two being the Kolmogorov–Obukhov scaling [3, 4], refined by the intermittency corrections of [5, 6] and the multifractal picture of [7], and the middle relation being Hölder continuity with exponent 1/3, the Onsager exponent. As $\nu \rightarrow 0$ the structures refine without limit while the velocity stays bounded at every scale: the gradient diverges, so the solution is not smooth, and the velocity does not, so it is not singular. A Hölder-1/3 field is exactly the object that is neither differentiable nor unbounded, which is why no amount of regularity theory reaches it.

This category is not merely hypothetical. On Onsager’s account [35], developed in [11, 12, 13], non-smoothness is the *mechanism* by which a turbulent flow dissipates energy at a rate that does not vanish with the viscosity — the dissipation anomaly, observed since Taylor [10] and still under active examination [9] — and a local energy equation can be written in which the non-smoothness is the direct cause of the dissipation [14]. Existence of weak solutions has been available since Leray [37, 42], and for small data, the low Reynolds number case for suitably scaled problems, global existence of a smooth solution has been known for sixty years [1, 2]. The difficulty lies at the other end, and that is the end ASDE addresses.

5 The (C) solution under the viscosity test

Two observations about the announced solution, offered without disputing it.

(C) is not the negation of (A). Fefferman’s four statements do not form two pairs of negations. (A) and (B) are posed with $f \equiv 0$ and quantify over *all* admissible data; (C) and (D) ask for the *existence* of admissible data *together with an admissible smooth forcing* f for which smoothness fails. So a proof of (C) leaves (A) untouched: it does not show that a fluid tears itself into a singularity, only that something can be done to it that does. The paper states its own scope in just these terms — it establishes (C), with the periodic case giving (D) — and does not claim (A) or (B). So (A) remains open after it, which is why a solution of (A) is worth presenting.

The viscosity test. The test is general and applies to any construction exhibiting a singularity at prescribed forcing. Navier–Stokes rescales: if v solves the equations at viscosity one with forcing g , then

$$u(x, t) = a v(bx, ct), \quad a = \nu b, \quad c = ab, \quad f = a^2 b g, \quad (4)$$

solves them at viscosity ν , and the map is onto: *one* unit-viscosity blowup generates the entire family. A theorem asserting blowup “for every $\nu > 0$ ” is therefore not a family of results. It is one result, rescaled, containing exactly what its $\nu = 1$ member contains.

Everything then turns on what (4) does to the constants, and since it is the forcing class that the official statement fixes, two normalisations bear on it. Hold the solution at size one, $a = 1$: then the forcing amplitude is $a^2 b = \nu^{-1}$, so the force required *blows up* as the viscosity falls and leaves any fixed admissible class. Or hold the forcing in a fixed class, $a^2 b = 1$: then $b = \nu^{-2/3}$ and

$$\text{amplitude } a = \nu^{1/3} \rightarrow 0, \quad \text{length } b^{-1} = \nu^{2/3} \rightarrow 0, \quad \text{time } c^{-1} = \nu^{1/3} \rightarrow 0. \quad (5)$$

The singularity shrinks to nothing: at $\nu = 10^{-6}$ the velocity reaches 10^{-2} on a scale of 10^{-4} before blowing up. Neither branch survives the limit — one exits the admissible data, the other converges to the zero solution — and there is no third option. Set this beside the flow the problem is about: turbulence at small viscosity has amplitude *fixed* and structure refining, by (3). The constructed family has amplitude vanishing like $\nu^{1/3}$, which is the opposite behaviour, or else forcing diverging like ν^{-1} , which is not a fluid at all. The independent exclusion of the mechanism under analytic forcing [34] points the same way from a different direction.

6 In closing

Two constructions, then, of the same formal kind and opposite in direction. The (C) solution fixes the singularity wanted and solves backwards for a force that produces it, the force being whatever the residual requires. ASDE fixes the data a flow actually has and solves forwards, the residual becoming the bound on what may be asked of the result. Both are automated, both exhibit, and only the second exhibits a fluid.

We do not ask for the problem to be restated, and we do not claim to have discharged (A) as it is formulated. What we present is what ASDE delivers when directed at it — constructive existence of turbulent solutions and qualitative uniqueness of their mean-value outputs, for specific data, with error bounds the computation itself produces — and we leave the reader to judge how near that comes to the question the problem was meant to ask. The material has been in print for some twenty years, from [31] onward and set out in Chapter 14 of [38];

the programme has continued since into compressible flow and aeronautical Reynolds numbers. What has changed is that a solution of (C) now exists, by methods that are themselves a new form of mathematics, and that makes it worth asking what the same kind of mathematics delivers on (A).

We end on the terms the problem set for itself. Fefferman is explicit about why four alternatives are offered rather than one: “To give reasonable leeway to solvers while retaining the heart of the problem, we ask for a proof of one of the following four statements” [36]. The Institute’s description says where that heart lies — waves follow the boat across the lake, turbulent air currents follow the aircraft, and the secrets are said to be hidden in the equations.

We have taken the leeway and used it to reach the heart. The flows constructed here are turbulent, at high Reynolds number, in bounded domains, for data a fluid actually has; they exist by exhibition and their mean-value outputs are determined to a stated precision. That is what ASDE delivers when directed at alternative (A), and we offer it for what it is worth.

Declaration of interest

The Adaptive Euler simulations cited above are the work of Icarus Digital Math, of which the author is a co-founder; Johan Jansson, whose results are cited, is a co-founder and its chief executive. The evidence those simulations provide is therefore not independent of the author, and should be weighed accordingly.

What is independent is the comparison. The High Lift Prediction Workshop and the high-order CFD workshops set the cases, supply the meshes and hold the measurements, and a prediction submitted before the measurement is disclosed is judged by people with no interest in this article. That is the part of the argument we would ask a sceptical reader to rely on, and it does not require taking anything here on trust.

References

- [1] H. Fujita and T. Kato, *On the Navier–Stokes initial value problem I*, Arch. Rational Mech. Anal. **16** (1964) 269–315.
- [2] T. Kato, *Strong L^p -solutions of the Navier–Stokes equation in $\mathbb{R}^m$, with applications to weak solutions*, Math. Z. **187** (1984) 471–480.
- [3] A. N. Kolmogorov, *The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers*, Dokl. Akad. Nauk SSSR **30** (1941) 301–305.
- [4] A. M. Obukhov, *On the distribution of energy in the spectrum of turbulent flow*, Bull. Acad. Sci. USSR, Geog. Geophys. Ser. **5** (1941) 453–466.
- [5] A. N. Kolmogorov, *A refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at high Reynolds number*, J. Fluid Mech. **13** (1962) 82–85.
- [6] A. M. Obukhov, *Some specific features of atmospheric turbulence*, J. Geophys. Res. **67** (1962) 3011–3014.

- [7] U. Frisch and G. Parisi, *On the singularity structure of fully developed turbulence*, in M. Ghil, R. Benzi and G. Parisi (eds.), *Turbulence and Predictability in Geophysical Fluid Dynamics and Climate Dynamics*, North-Holland, 1985.
- [8] B. Musci, B. Dubrulle, J. LeBris, D. Geneste, P. Braganca, J.-M. Foucaut, C. Cuvier and A. Cheminet, *Experimental measurement of the vorticity–strain alignment around extreme energy transfer events*, J. Fluid Mech. **1016** (2025) A53.
- [9] K. P. Iyer, T. D. Drivas, G. L. Eyink and K. R. Sreenivasan, *Turbulence without walls: whither the zeroth law of turbulence?*, Phys. Rev. Lett. **135** (2025) 134001.
- [10] G. I. Taylor, *Statistical theory of turbulence*, Proc. R. Soc. Lond. A **151** (1935) 421–444.
- [11] G. L. Eyink, *Energy dissipation without viscosity in ideal hydrodynamics I. Fourier analysis and local energy transfer*, Physica D **78** (1994) 222–240.
- [12] C. De Lellis and L. Székelyhidi Jr., *Dissipative Euler flows and Onsager’s conjecture*, J. Eur. Math. Soc. **16** (2014) 1467–1505.
- [13] T. Buckmaster, C. De Lellis, P. Isett and L. Székelyhidi Jr., *Anomalous dissipation for $1/5$ -Hölder Euler flows*, Ann. of Math. **182** (2015) 127–172.
- [14] J. Duchon and R. Robert, *Inertial energy dissipation for weak solutions of incompressible Euler and Navier–Stokes equations*, Nonlinearity **13** (2000) 249–255.
- [15] N. Fehn, M. Kronbichler and G. Lube, *From anomalous dissipation through Euler singularities to stabilized finite element methods for turbulent flows*, Flow Turbul. Combust. **115** (2025) 347–388.
- [16] J. Kronborg and J. Hoffman, *The triple decomposition of the velocity gradient tensor as a standardized real Schur form*, Phys. Fluids **35** (2023) 031703.
- [17] J. Hoffman, *Energy stability analysis of turbulent incompressible flow based on the triple decomposition of the velocity gradient tensor*, Phys. Fluids **33** (2021) 081707.
- [18] J. Hoffman, J. Jansson and R. Vilela de Abreu, *Adaptive modeling of turbulent flow with residual based turbulent kinetic energy dissipation*, Comput. Methods Appl. Mech. Engrg. **200** (2011) 2758–2767.
- [19] J. Hoffman, J. Jansson, N. Jansson and R. Vilela de Abreu, *Towards a parameter-free method for high Reynolds number turbulent flow simulation based on adaptive finite element approximation*, Comput. Methods Appl. Mech. Engrg. **288** (2015) 60–74.
- [20] J. Hoffman, J. Jansson et al., *Unicorn: Parallel adaptive finite element simulation of turbulent flow and fluid–structure interaction*, Comput. Fluids, 2013.
- [21] N. Jansson, J. Hoffman and J. Jansson, *Framework for massively parallel adaptive finite element computational fluid dynamics*, SIAM J. Sci. Comput., 2012.
- [22] J. Hoffman, J. Jansson and N. Jansson, *FEniCS-HPC: Automated predictive high-performance finite element computing with aerodynamics applications*, Lecture Notes in Computer Science, Springer, 2016.

- [23] J. Hoffman, J. Jansson *et al.*, *FEniCS-HPC: Coupled multiphysics in computational fluid dynamics*, Lecture Notes in Computer Science, Springer, 2017.
- [24] A. Logg, T. Dupont, J. Hoffman, C. Johnson, R. C. Kirby, M. G. Larson and L. R. Scott, *The FEniCS project*, Technical Report 2003–21, Chalmers Finite Element Center Preprint Series, 2003.
- [25] A. Logg, *Automation of Computational Mathematical Modeling*, Ph.D. thesis, Chalmers University of Technology, 2004.
- [26] A. Logg, *Automated solution of differential equations*, Sixth International Congress on Industrial and Applied Mathematics (ICIAM07) and GAMM Annual Meeting, Zürich, 2007.
- [27] A. Logg, K.-A. Mardal and G. N. Wells (eds.), *Automated Solution of Differential Equations by the Finite Element Method: The FEniCS Book*, Springer, 2012.
- [28] M. S. Alnæs, J. Blechta, J. Hake, A. Logg *et al.*, *The FEniCS Project Version 1.5*, Archive of Numerical Software **3** (100) (2015) 9–23.
- [29] K. Eriksson, A. Logg and C. Johnson, *Adaptive computational methods for parabolic problems*, in *Encyclopedia of Computational Mechanics*, 2017, pp. 1–28.
- [30] C. Johnson, J. Hoffman and A. Logg, *Topics in adaptive computational methods for differential equations*, CEDYA 2001: Congreso de Ecuaciones Diferenciales y Aplicaciones, 2001.
- [31] J. Hoffman and C. Johnson, *A new approach to computational turbulence modeling*, Comput. Methods Appl. Mech. Engrg. **195** (2006) 2865–2880.
- [32] OpenAI, *Finite time blowup for Navier–Stokes*, preprint, 2026. <https://cdn.openai.com/pdf/32d9f210-8b73-45e0-91bc-82a30aef8a9a/navier-stokes.pdf>
- [33] T. Buckmaster, *Statement*, 2026. <https://cims.nyu.edu/~tristanb/statement.pdf>
- [34] P. Constantin, M. Ignatova and V. Vicol, *Regularity of asymptotically axisymmetric solutions to the 3D Navier–Stokes equations with analytic forcing*, arXiv:2609.20803, 2026.
- [35] L. Onsager, *Statistical hydrodynamics*, Nuovo Cimento **6**, Suppl. 2 (1949) 279–287.
- [36] C. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, Official Problem Description, Clay Mathematics Institute, 2000.
- [37] J. Leray, *Sur le mouvement d’un liquide visqueux emplissant l’espace*, Acta Math. **63** (1934) 193–248.
- [38] J. Hoffman and C. Johnson, *Computational Turbulent Incompressible Flow*, Applied Mathematics: Body and Soul, vol. 4, Springer, 2007.
- [39] J. Hoffman and C. Johnson, *Resolution of d’Alembert’s paradox*, J. Math. Fluid Mech. **12** (2010) 321–334.
- [40] J. Hoffman, J. Jansson and C. Johnson, *New theory of flight*, J. Math. Fluid Mech. **18** (2016) 219–241.

- [41] J. Hoffman, *Efficient computation of mean drag for the subcritical flow past a circular cylinder using general Galerkin G2*, Int. J. Numer. Meth. Fluids **59** (2009) 1241–1258.
- [42] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [43] J. Jansson, *Adaptive Direct FEM Simulation with Unicorn/FEniCS-HPC for CS1*, 5th International Workshop on High Order CFD Methods; Zenodo, 2022. <https://doi.org/10.5281/zenodo.6942599>.
- [44] J. Jansson, *Adaptive Euler: validation against NASA High Lift Prediction Workshop 4, and preliminary tail-buffet and transonic-buffet simulations*, presented at IFASD 2026; video, https://www.linkedin.com/posts/johan-jansson-56aa0a41_fenics-activity-7472908903618338817--spa.
- [45] C. Johnson, posts under the label *clay problem*, <https://claesjohnson.blogspot.com/search/label/clay%20problem>.