

The Spinor Residue: $g = 2$ and Stern–Gerlach without Relativity

Claes Johnson¹

¹KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden ,
claesjohnson@gmail.com

July 19, 2026

Argument by the author; drafting prepared with AI assistance (Claude, Anthropic), reviewed and verified by the author.

Abstract

The charges-as-configurations reading of RealQM removes relativity and gives the electron’s magnetic moment as a quantised current-winding: an orbital winding $e^{im\phi}$ carries $\mu_z = -m\mu_B$ with gyromagnetic factor $g = 1$. This leaves exactly one residue — the electron’s *spin* moment has $g = 2$, twice the orbital value per unit angular momentum — and one experiment that RealQM in its present form cannot reproduce: *Stern–Gerlach*, whose two-spot splitting is the signature of two-valued spin $\frac{1}{2}$. We make the residue precise, and argue that it is *not* a demand for relativity. By Lévy-Leblond’s theorem $g = 2$ follows from a first-order (linearised), *non-relativistic* spinor wave equation; the “2” is the SU(2) double cover of the rotation group, a geometric fact, not a consequence of Lorentz invariance. Concretely, writing the kinetic energy with $\boldsymbol{\sigma} \cdot \mathbf{p}$ rather than p^2 produces, in a field, the term $-q\boldsymbol{\sigma} \cdot \mathbf{B}$ — the $g = 2$ Zeeman coupling — automatically. The target for RealQM is therefore identified without importing relativity: equip the non-overlapping domains with the $\pm\frac{1}{2}$ spinor label the base theory already suggests, implemented as a $\boldsymbol{\sigma} \cdot \mathbf{p}$ structure, so that the moment becomes two-valued ($\mu_z = \pm\mu_B$) and Stern–Gerlach’s two spots and $g = 2$ follow together. We are explicit that this is a *target*, not a completed derivation: what is established is that the residue is geometric and reachable, not relativistic.

1 The one residue

In the current-bearing form of RealQM the magnetic moment is that of the charge current, $\boldsymbol{\mu} = \frac{1}{2} \int \mathbf{r} \times \mathbf{J}$, and an orbital phase winding $e^{im\phi}$ gives the quantised value $\mu_z = -m\mu_B$ [1, 2]. This is gyromagnetic factor $g = 1$: one unit of orbital angular momentum, $L_z = m\hbar$, yields m Bohr magnetons. The charges-as-configurations picture [3] builds on this to remove any need for relativity: sizes and moments are properties of Coulomb configurations, a “free” electron is only a weakly-confined one, and the winding it can afford gives μ_B with no rest-energy scale invoked.

Exactly one thing is left unaccounted. The electron’s *spin* moment is not $g = 1$ but $g = 2$: a spin of only $\hbar/2$ produces a *full* Bohr magneton, twice what an orbital current of the same angular momentum would give. An orbital winding cannot supply this factor of two; it is the signature of spin, and RealQM has, so far, no spin- $\frac{1}{2}$ moment. This is the residue.

2 Its experimental face: Stern–Gerlach

The residue is not academic; it is the whole content of the Stern–Gerlach experiment. A beam of silver atoms — each with a single valence electron in an $L = 0$ (orbitally spherical) ground state — passes through an inhomogeneous magnetic field and splits into *two* discrete spots. Two spots is the signature of a two-valued moment, $\mu_z = \pm\mu_B$, i.e. spin $\frac{1}{2}$ with $g = 2$; an orbital moment would split into an *odd* number $2l + 1$, and for the $L = 0$ silver electron would not split at all ($m = 0$, $\mu_z = 0$).

RealQM in its present form therefore *fails* Stern–Gerlach: with only orbital windings the $L = 0$ valence electron carries no moment and the beam would pass undeflected. Reproducing the two-spot pattern requires precisely the two-valued spin- $\frac{1}{2}$ structure that is missing. Stern–Gerlach and $g = 2$ are one problem: the two spots *are* $g = 2$ made visible.

3 The residue is geometric, not relativistic

It is tempting to conclude that spin, $g = 2$, and Stern–Gerlach force relativity into the theory, since in the textbook account $g = 2$ falls out of the Dirac equation. This conclusion is false, and the point is decisive for the present programme.

Lévy-Leblond. In 1967 Lévy-Leblond showed that $g = 2$ follows from a first-order (linearised) *non-relativistic*, Galilean spinor wave equation [4]. Relativity is sufficient to produce $g = 2$ but not necessary; the factor of two survives the non-relativistic limit because it does not come from Lorentz invariance in the first place.

The mechanism. Write the kinetic energy not as $p^2/2m$ but as $(\boldsymbol{\sigma} \cdot \mathbf{p})^2/2m$ — legal for free motion since $(\boldsymbol{\sigma} \cdot \mathbf{p})^2 = p^2$. In a magnetic field, minimal coupling $\mathbf{p} \rightarrow \mathbf{p} - q\mathbf{A}$ gives

$$\frac{(\boldsymbol{\sigma} \cdot (\mathbf{p} - q\mathbf{A}))^2}{2m} = \frac{(\mathbf{p} - q\mathbf{A})^2}{2m} - \frac{q}{2m} \boldsymbol{\sigma} \cdot \mathbf{B}, \quad (1)$$

and the last term is exactly the $g = 2$ spin–Zeeman coupling, appearing *automatically* the moment the dynamics is linearised in $\boldsymbol{\sigma} \cdot \mathbf{p}$. No relativity is used in (1); it is the ordinary Pauli Hamiltonian.

The geometry. The “2” is the **SU(2) double cover of the rotation group**: a spinor returns to itself only after 720° , rotating at half angle. That half is reciprocally the double in the moment. It is the same species of fact as the winding-number topology that fixes $\mu_z = -m\mu_B$ in the orbital case — a statement about how spin- $\frac{1}{2}$ objects transform, geometric and scale-free, owing nothing to rest energy.

4 The target for RealQM

The residue thus lands on the reachable side of the ledger. What RealQM needs is not relativity but the right *spinor structure* on its domains, and the base theory already points at it: the ψ_i on

Ω_i acquiring an internal $\pm\frac{1}{2}$ label. The programme is to implement that label not as a bolt-on but as a $\boldsymbol{\sigma} \cdot \mathbf{p}$ linearisation of the one-electron dynamics on each domain, so that:

- the spin–Zeeman term $-\frac{q}{2m}\boldsymbol{\sigma} \cdot \mathbf{B}$ of (1) appears, giving $g = 2$;
- the moment becomes *two-valued*, $\mu_z = \pm\mu_B$, so that an $L = 0$ electron in an inhomogeneous field splits into *two* beams — Stern–Gerlach — rather than none;
- the non-overlap/free-boundary structure, and the current $\mathbf{J}$ of the complex form, are retained unchanged, since (1) is a rewriting of the kinetic term, not a new interaction.

We state plainly that this is a *target*, not a finished calculation: we have not carried the $\boldsymbol{\sigma} \cdot \mathbf{p}$ structure through the multiphase free-boundary problem, and doing so — in particular checking that the two-valued moment coexists with the geometric realisation of Pauli exclusion by non-overlap without double-counting the spin — is the open foundational step. What is established is the map: $g = 2$ and Stern–Gerlach are one residue, that residue is the $SU(2)$ spinor factor of two, and it is obtainable non-relativistically, so it belongs inside RealQM rather than outside it.

5 Conclusion

The charges-as-configurations picture closes the free-charge and relativity problems and leaves a single, sharp residue: the electron’s spin factor $g = 2$, whose experimental embodiment is the two-spot Stern–Gerlach splitting. That residue is not a relativistic quantity smuggled in through the Dirac equation; by Lévy-Leblond it is a non-relativistic, geometric consequence of the $SU(2)$ spinor structure, delivered by a $\boldsymbol{\sigma} \cdot \mathbf{p}$ linearisation. RealQM therefore has a concrete, relativity-free target for spin: give its domains the $\pm\frac{1}{2}$ label as a $\boldsymbol{\sigma} \cdot \mathbf{p}$ structure, and $g = 2$ and Stern–Gerlach’s two beams should follow together. Until that is carried through the multiphase problem the point is a target, not a result — but it is the right size of target, and it is inside the theory, not beyond it.

References

- [1] C. Johnson, *Complex Time-Dependent RealQM: Currents, Radiation, and Magnetism from the Real-Space Continuum*, 2026.
- [2] C. Johnson, *Magnetism as Circulating Charge: Why the Nuclear Electron Carries No Magnetic Moment*, 2026.
- [3] C. Johnson, *Elementary Charges as Configurations, not Objects: Why RealQM Needs No Relativity*, 2026.
- [4] J.-M. Lévy-Leblond, *Nonrelativistic particles and wave equations*, Commun. Math. Phys. **6**, 286 (1967).