Shell Occupancy in RealQM: the Two Constructions Bracket the Exact Energy, and the Variational Principle Picks the Wrong End

Claes Johnson[1]

[1]Royal Institute of Technology, Stockholm (emeritus)

Abstract

RealQM represents N electrons by N unit charge densities on non-overlapping domains of ordinary space, with the energy a localisation term plus the Coulomb energy of the configuration, pairs counted once. For a shell holding more than one electron the construction admits two readings, and both are in use: the electrons may be *split* into disjoint angular sectors, or *homogenised* into one shared density with the self-repulsion reduced by $(N_s-1)/N_s$. The first is what the stated non-overlap condition prescribes; the second is what the accurate atomic calculations use. On helium they give -3.018 and -2.840 Hartree against an exact -2.9037, so the choice is worth about 6% and nothing in the theory fixes it.

We connect the two constructions by a one-parameter family and locate the exact answer inside it. The connection rests on an identity: at an interface which is a plane of symmetry, the kinetic energy of the *sum* $\psi=\psi_1+\psi_2$ equals the sum of the two domain kinetic energies computed with the natural condition on each domain — to machine precision. Both ends of the family therefore carry the *same* kinetic energy, and the entire difference between them is the pair repulsion, which is a bounded functional. Parameterising the repulsion rather than the densities removes an obstruction: a family built on the densities sends the localisation cost to infinity at the split end, because a jump in $\sqrt{\rho}$ is not in H^1, and the split end is then unreachable.

With the kernel softening calibrated on the *one-electron* exact energy $-Z^2/2$, so that no two-electron information enters, the family reproduces both published endpoints of helium to 0.2–0.3%, and the exact two-electron energy lies strictly inside the bracket for $Z=2,3,4$ at a nearly constant fraction 0.36, 0.32, 0.30 of the way from overlap towards split.

The energy is monotone along the family. Minimising over the parameter therefore drives it to the split end, which over-binds by 4.2%. Two consequences follow. The occupancy parameter is not determined by the variational principle and must be supplied, which places it alongside the core radius rather than among the model's outputs. And the homogenised shell, which is the more accurate of the two, is not a variational competitor at all but a higher-energy configuration that happens to sit nearer measurement — so its accuracy is cancellation rather than correctness. We state the mesh limitation of the numbers and what would settle them.

1 The ambiguity

The RealQM energy for N electrons on domains Ω_i is a localisation term plus the Coulomb energy with pairs counted once and no self-interaction. For one electron per domain the prescription is unambiguous. For a shell holding $N_s>1$ electrons it is not, and two readings are in use.

The *split* reading follows the stated non-overlap condition: the N_s electrons occupy disjoint angular sectors of the shell, two of them as hemispheres, four as a tetrahedron, eight as two interleaved tetrahedra. The *homogenised* reading gives all N_s electrons one shared spherically symmetric density and reduces the self-repulsion by $(N_s-1)/N_s$, which is pair counting restated. The accurate atomic calculations use the second; the stated condition prescribes the first.

On helium the difference is not small. The two constructions give -3.018 and -2.840 Ha against an exact -2.9037, so the choice is worth about 6% of the correlation-sized part of the energy and straddles the answer. Nothing in the formulation selects between them.

2 The two ends share their kinetic energy

The obstruction to connecting them is apparent rather than real, and removing it is the technical content of this note.

Write $\psi = \psi_1 + \psi_2$ for the sum of the two one-electron functions. In the split construction each ψ_i is supported on its own half and is discontinuous across the interface, so $\psi_i \notin H^1$ and its localisation energy is not defined as a distribution. But the sum is smooth, and the quantity actually computed in a split calculation — the localisation energy assembled over each domain with the natural condition on its own boundary — is exactly the localisation energy of that sum. We verified this numerically on an axisymmetric grid: for a state symmetric in z with the interface on a cell face,

$$T[\psi] = 1.40322755, \qquad T_1^{\mathrm{dom}} + T_2^{\mathrm{dom}} = 1.40322755,$$

agreeing to 2.2×10^{-16}, and likewise at a second spacing. For a state deliberately skewed so that $\partial\psi/\partial z \neq 0$ at the interface the two differ by 1–2%, which is the straddling-face term that the symmetry kills. So at a plane of symmetry the split construction's kinetic energy is not an approximation to anything: it is the kinetic energy of one smooth function, attributed by region.

Two consequences. First, no state drops to zero anywhere and no H^1 condition is violated, because ψ_1 and ψ_2 are never formed separately. Second, and this is what makes the connection possible, both ends of the construction carry the *same* kinetic energy for a given total density:

$$\text{homogenised: } \rho_i = \psi^2 \;\Rightarrow\; T = \int |\nabla\psi|^2, \qquad \text{split: } \rho_i = 2\psi^2 \mathbf{1}_{\mathrm{half}} \;\Rightarrow\; T = \int |\nabla\psi|^2.$$

The entire difference between the constructions is the pair repulsion.

3 The family, and why it must be built on the repulsion

Let w be a weight with $w(z) + w(-z) = 1$ and set $\rho_1 = 2w\psi^2$, $\rho_2 = 2(1-w)\psi^2$ with $\int \psi^2 = 1$. Taking $w = \frac{1}{2}$ gives the homogenised shell and w a step gives the split shell. Holding the localisation and nuclear terms at their common value,

$$E(\lambda) = \int |\nabla\psi|^2 - 2Z\int \frac{\psi^2}{r} + U(\lambda), \qquad U(\lambda) = \int P[\rho_1]\,\rho_2, \qquad w = \left(1 + e^{-z/\lambda}\right)^{-1}, \quad (1)$$

minimised over ψ at each λ. Both ends are finite: $U \to J[\psi^2]$ as $\lambda \to \infty$ and $U \to$ the cross-term between disjoint halves as $\lambda \to 0$.

It matters that the family is built on U and not on the densities. A family built by partitioning $\sqrt{\rho}$ carries a localisation cost $\frac{1}{2}\int\psi^2 Q$ with $Q = \sum_i |\nabla\sqrt{2w_i}|^2$, which diverges as w approaches a step — a jump in $\sqrt{\rho}$ is not in H^1. We computed that family as well; its energy rises monotonically away from full overlap, the split end is unreachable, and the exact answer lies outside it altogether. The apparent bracketing of the exact energy is recovered only when the parameterisation is placed on the bounded functional.

4 The exact energy lies inside, at a nearly constant position

The kernel is softened, and the softening is calibrated *per element on the one-electron exact energy* $-Z^2/2$, so that no two-electron quantity enters the calibration and the two-electron result is a prediction. On helium the family then returns -2.840 at the overlap end against the published homogenised -2.845, and -3.018 at the split end against the published -3.026: both ends reproduced to 0.2–0.3% by a calibration that used neither.

Z	overlap	exact	split	inside?	λ^*	fraction along
2	-2.840	-2.904	-3.018	yes	0.275	0.36
3	-7.198	-7.280	-7.454	yes	0.210	0.32
4	-13.555	-13.656	-13.892	yes	0.170	0.30

The exact energy lies strictly inside the bracket for all three ions. The parameter itself is not a clean law: $\lambda^* \sim Z^{-0.7}$, so $Z\lambda^*$ drifts by 24% across the series. What is nearly constant is the position *along* the family, 0.36, 0.32, 0.30 — the exact answer sits about one third of the way from overlap towards split, to within $\pm 9\%$, drifting slowly with Z.

The scaling is physically sensible. λ is the length over which the two territories blend, so it is an avoidance length; the atom shrinks as $1/Z$ while λ^* shrinks more slowly, so the blending region grows relative to the atom as Z rises — which is what one expects when the nuclear attraction increasingly dominates the mutual repulsion and the electrons have less reason to avoid one another.

5 The variational principle picks the wrong end

$E(\lambda)$ is monotone along the family: for helium, -2.845, -2.852, -2.868, -2.893, -2.929, -2.962, -2.990, -3.006 as λ decreases. Minimising over λ therefore drives it to zero, and the minimiser is the fully split configuration, which over-binds by 4.2%.

Two conclusions follow, and they are the point of this note.

The occupancy parameter is not an output. It cannot be obtained by minimising, because minimising selects an end rather than the interior. It therefore has to be supplied, which places it alongside the core radius rather than among the model's results. A construction with one electron per domain and nothing else specified does not determine a shell's occupancy.

The accurate construction is not the variational one. Strict non-overlap is admissible and has the lowest energy in the family, so it is what "minimise the energy" selects — and it over-binds. The homogenised shell, the more accurate of the two and the one behind the

published atomic totals, is a *higher*-energy configuration lying nearer measurement. Its accuracy is therefore cancellation rather than correctness, and should not be read as evidence that it is the right reading of the construction.

6 What these numbers do not establish

One mesh per element, with no mesh ladder. The endpoint agreement at 0.2–0.3% is reassuring but is not a convergence check, and the fractions in the table are differences of differences, hence the most mesh-sensitive quantity reported. We would want 0.36, 0.32, 0.30 reproduced at a second spacing before the near-constancy is treated as established rather than suggestive.

The kernel is softened rather than bare. The softening is calibrated on a one-electron quantity, which is the cleanest available choice and makes the two-electron numbers predictions, but it is a regularisation and the absolute energies carry its error.

Only two-electron shells are treated. Whether the same position along the family describes a shell of four or eight — where the split is a tetrahedron rather than a plane, and where the capacities 2 and 8 of the pair-and-octet construction are at stake — is untested here, and is the obvious continuation.