

clinical investigations in critical care

Interest of a Therapeutic Optimization Strategy in Severe ARDS*

Nathalie Guinard, MD; Sadek Beloucif, MD, PhD; Claire Gatecel, MD; Joaquim Mateo, MD; and Didier Payen, MD, PhD

Study objective: Evaluate the interest of the response to a therapeutic optimization as a predictor of prognosis in ARDS.

Design: Prospective study.

Setting: ICU of a University Hospital.

Patients: Thirty-six consecutive patients with severe ARDS addressed for extracorporeal carbon dioxide removal (ECCO₂R).

Interventions: We studied the response during the first 2 days after arrival to the therapeutic optimization strategy consisting in a combination of the following: (1) decrease in extravascular lung water (diuretics or hemofiltration); (2) selection of the best ventilatory mode; (3) permissive hypercarbia; and (4) correction of hypoxemia by alveolar recruitment, additional continuous oxygen insufflation, body position changes (prone position), inhaled nitric oxide, enhancement of hypoxic pulmonary vasoconstriction with almitrine, and drainage of pleural or mediastinal effusions. In patients remaining severely hypoxemic despite these modalities, ECCO₂R was then proposed.

Measurements and results: Thirty-six patients were addressed after 8.3 ± 5.5 days of mechanical ventilation. On arrival, mean simplified acute physiologic score was 46.8 ± 14.2 , multiple system organ failure score was 1.8 ± 1.6 , Murray score was 3.4 ± 0.4 , PaO₂ was 75.3 ± 31.3 (fraction of inspired oxygen [FIO₂]=1) for a positive end-expiratory pressure level of 12.3 ± 3.4 cm H₂O. Nineteen of 36 patients improved their gas exchange within 2 days and their mortality was 21%. The seventeen remaining patients did not improve PaO₂/FIO₂; PaCO₂ and airway pressures remained high and their mortality was 88%. This different response to therapeutic optimization appeared using stepwise logistic regression as the most predictive factor for mortality ($p < 0.05$). **Conclusions:** In patients with severe ARDS, the response to an early performed therapeutic optimization used to improve hypoxemia appeared to be a highly discriminant factor distinguishing deceased from surviving patients. (CHEST 1997; 111:1000-07)

Key words: adult respiratory distress syndrome (ARDS); extracorporeal CO₂ removal (ECCO₂R); hypoxia; prognosis factor; severity scores; therapy

Abbreviations: ECCO₂R=extracorporeal carbon dioxide removal; ECMO=extracorporeal membrane oxygenation; FIO₂=fraction of inspired oxygen; LFPPV=low-frequency positive pressure ventilation; MSOF=multiple system organ failure; NO=nitric oxide; PEEP=positive end-expiratory pressure; PIP=peak inspiratory pressure; SAPS II=simplified acute physiologic score II; TO=therapeutic optimization

*From the Department of Anesthesiology and Critical Care Medicine, Hôpital Universitaire Lariboisière, Paris, France. Partially supported by Institutional Grants from University Paris VII, and the Direction de la Recherche et de l'Enseignement Doctoral (DRED).

Manuscript received March 22, 1996; revision accepted October 15.

Reprint requests: Didier Payen, MD, PhD, Department of Anesthesiology and Critical Care Medicine, Hôpital Universitaire Lariboisière, 2 Rue Ambroise Paré, 75475 Paris Cedex 10, France

More than 25 years after its initial description,¹ ARDS remains a challenging organ failure to assist, with a high mortality rate. Despite efforts to clinically define ARDS,² a multifactorial disease, a

For editorial comment see page 845

disparity between mortality rates is noted according

to the different diagnosis criteria used.³ The last decade has consisted in development of several techniques or therapies to support lung failure, but none of them has been evaluated in terms of mortality and morbidity. The recent introduction of permissive hypercapnia,⁴⁻⁷ ultrafiltration for lung water control,⁸⁻¹⁰ low insufflation pressures to prevent barotrauma,^{5,6,11,12} and nitric oxide (NO) inhalation,¹³⁻¹⁶ associated or not with almitrine,^{17,18} might modify the therapeutic management of such patients. Finally, if extracorporeal carbon dioxide removal (ECCO₂R) has been proposed for many years,¹⁹ its indications remain difficult, especially if one considers that the recent therapeutic advances improve the parameters originally used to indicate such an invasive support. Recently, it has been shown that the number of extracorporeal membrane oxygenation (ECMO) procedures performed in pediatric patients has been reduced with the introduction of NO inhalation therapy.²⁰

The impact on mortality or morbidity of most of these recent therapies has not been yet evaluated by multicentric studies (to our knowledge). However, they might help to stratify the patients as responders or nonresponders to supportive therapies for hypoxemia. We hypothesized that the response to a therapeutic optimization strategy, prospectively tested in patients with severe ARDS, could predict the outcome of such patients. Using stepwise logistic regression, this response was the strongest predictive factor for outcome. Finally, such an approach selected a subgroup of patients with extremely severe disease for whom future trials could be proposed.

MATERIALS AND METHODS

Patients

Thirty-six consecutive patients suffering from primary or secondary ARDS, without any history of respiratory failure and addressed for ECCO₂R with low-frequency positive pressure ventilation (LFPPV) from April 1991 to November 1995 were studied if the following criteria were present: (1) a Lung Injury Score (Murray Score)²¹ higher than 2.5; and (2) entry criteria used in the ECMO study,²² *ie*, either rapid entry criteria (PaO₂ <50 mm Hg for >2 h when measured at an inspired oxygen fraction [FIO₂] of 1.0 and a positive end-expiratory pressure [PEEP] ≥5 cm H₂O) or slow entry criteria (PaO₂ <50 mm Hg for >12 h when measured at FIO₂ ≥0.6 and a PEEP ≥5 cm H₂O after 48 h of maximal medical therapy). On arrival, simplified acute physiologic score II (SAPS II)²³ and multiple system organ failure (MSOF) scoring were calculated according to criteria of Gattinoni et al.¹⁹ Patients were sedated, paralyzed, and mechanically ventilated (with a Servo 900 C Ventilator; Siemens Elema; Lund, Sweden). Quasistatic lung compliance, mean peak inspiratory, and end-expiratory airway pressures were measured using a ventilatory module (Sirecust 1280; Siemens). Presence of barotraumatic lesions²⁴ with evidence of extra-alveolar gas (pneumo-

thorax, pneumomediastinum, pneumoperitoneum, subcutaneous emphysema, subpleural air cysts) was analyzed on chest radiograph and thoracic CT scan.

Decision-Making Strategy

Thoracic, abdominal, and cerebral CT scans were obtained in all patients. The initial thoracic CT scan allowed the evaluation of the extent of pulmonary edema, pleural effusion, localization of condensation, existence of barotrauma lesions, and bronchial alterations. These CT scan observations and the recently acquired advances in ARDS pathophysiology allowed us to define the following therapeutic optimization (TO) strategy, in addition to classical therapeutic support: (1) decrease in extravascular lung water (with diuretics or continuous hemofiltration) when water balance was positive; significant interstitial edema was present on thoracic CT scan and/or chest radiograph; (2) drainage of bilateral pleural effusion, especially when diffuse lung edema was present;⁸⁻¹⁰ (3) tentative action to limit pulmonary blood flow when a hyperkinetic syndrome was present, since in addition to the relationship between intrapulmonary shunt and pulmonary blood flow, passive pulmonary hypertension may potentially worsen lung edema;^{25,26} (4) selection of the best ventilatory mode to obtain the lowest possible tidal volume, peak inspiratory pressure (PIP), and mean airway pressure ensuring an adequate alveolar recruitment;¹² a goal of a maximum of 35 cm H₂O for PIP was considered acceptable, and permissive hypercarbia⁴⁻⁷ was allowed; and (5) correction of arterial hypoxia by ventilation/perfusion ratio manipulation: best alveolar recruitment ("best PEEP" level, inspiration to expiration (I/E) ratio modification); additional continuous tracheal oxygen insufflation;^{27,28} reduction of blood flow to shunting areas with body position changes based on CT scan imaging, vasodilation of nonshunting zones (NO inhalation <15 ppm monitored with chemiluminescence technique), and/or reinforcement of hypoxic pulmonary vasoconstriction with almitrine bismesylate (Servier; Suresnes, France).^{13-18,29-31}

All these modalities were used together or separately according to the repetitive evaluation of lung function and status based on gas exchange, radiologic patterns, and lung mechanics. The results of the TO allowed us to stratify the population of patients as shown in Figure 1. A patient was considered as a responder, when PaO₂ on FIO₂ of 1 was improved and stabilized over 100 mm Hg during at least 6 h. A patient was considered as a nonresponder when (1) PaO₂ on FIO₂ of 1 was lower than 100 mm Hg, or did not improve during at least 6 h; (2) control of PaCO₂ was difficult (even considering permissive hypercapnia); or (3) lung mechanics deteriorated. In these patients, even after several days of evaluation, ECCO₂R-LFPPV was discussed as a "last chance therapy." This invasive technique was performed as described by Gattinoni et al¹⁹ only after careful consideration of its contraindications:^{19,32} potential hemorrhagic lesions of the CNS with contraindications to anticoagulation; hemorrhagic disease; refractory shock; persistent organ failures exceeding four; major burns (total body surface >40%); chronic systemic disease that limits survival (*eg*, cancer, hematologic malignancy, severe inflammatory disease). When a contraindication to ECCO₂R-LFPPV was present, the patient was maintained under a regimen of conventional ventilation.

When ECCO₂R-LFPPV was performed, the extracorporeal circuit used was a venovenous bypass through two cannulas inserted percutaneously into the two femoral veins (the drainage cannula being located in the lower part of the inferior vena cava, and the return cannula just below the right atrium). The continuously rewarmed and heparinized blood was driven by an occlusive pump (COBE; Lakewood, Calif) through two silicone artificial membrane lungs [7 m² total membrane surface area

CLINICAL DECISION MAKING




FIGURE 1. Clinical decision-making strategy proposed.

(Ultrox I; Sci Med; Minneapolis)] used in series. A bypass flow of 20 to 30% of cardiac output allowed us to remove CO₂ since membranes were ventilated with a heated humidified mixture of air and oxygen. Lungs were continuously maintained inflated by a permanent positive pressure associated with three to four mechanical cycles per minute with peak airway pressure limited to 40 to 45 cm H₂O by pressure control. Mean airway pressure was kept similar to pre-ECCO₂R value (*ie*, close to the PEEP level). A continuous 1 to 2 L/min oxygen flow was added through a small catheter positioned in the tracheal tube, just above the carina, to cover the needs of whole body oxygen consumption. This average flow rate was increased in case of pleural leak by bronchopleural fistula.

Statistical Analysis

Comparisons between responders and nonresponders (including those treated or not with ECCO₂R) were performed with the nonparametric Mann-Whitney test, and evolution of a given parameter within a group with the nonparametric Wilcoxon test. Results are expressed as mean±SD. A *p* value <0.05 was considered significant.

Potential univariate correlates of mortality were identified by χ^2 analysis for qualitative parameters and Student's *t* test for quantitative parameters. Multivariate analysis was performed using stepwise logistic regression (Biomedical Data Processing Package; UCLA; Los Angeles). Many authors have stressed the difficulty in knowing precisely the exact value of

alpha in multiple regression procedures,^{33,34} which may lead to an overestimation of the number of predictive variables. In the present study, we used conservative criteria to select predictive variables: (1) limits to enter or remove variables in the regression equation must have had a 5% probability value; (2) for each candidate as a predictive factor, the ratio between the corresponding regression coefficient and its SE must have been greater than 2;³⁵ and (3) results were verified using two different numerical procedures, an asymptomatic covariance estimate and the maximum-likelihood method; *p* values corresponding to the improved χ^2 are given in the tables.

RESULTS

Patients' Characteristics on Arrival

Table 1 summarizes the clinical characteristics of the 36 patients on arrival. Mean SAPS II was 46.8±14.2, MSOF was 1.8±1.6, and Murray score was 3.4±0.4 (range, 2.6 to 4). Only nine patients suffered from isolated respiratory failure, without other system organ failure. The etiologies of the ARDS were both primitive (25 patients) and secondary (11 patients). Duration of mechanical ventilation before arrival was 8.2±5.5 days (range, 1 to 20 days). All patients had the rapid or slow ECMO criteria.

Table 1—Clinical Characteristics of Patients on Arrival

Patient No./Age, yr/Sex	Diagnosis or Risk Factor for ARDS	SAPS II	MSOF*	Murray Score	Duration of Mechanical Ventilation,† d
1/17/M	Fat embolism	42	3	3.7	1
2/22/F	Pneumonia	41	1	3.5	11
3/25/M	Pneumonia	37	2	3.7	1
4/22/M	Multiple trauma	46	0	3	11
5/36/M	Pneumonia	24	1	2.7	13
6/22/M	Multiple trauma	29	0	3.3	19
7/36/F	Gastric aspiration	26	1	3.5	7
8/44/M	Pneumonia	43	4	3	3
9/41/F	Gastric aspiration	41	0	2.6	2
10/51/M	Pneumonia	50	2	4	5
11/52/F	Vasculitis	52	4	3.66	7
12/48/M	Sepsis	47	1	3.75	7
13/25/F	Lung contusion	23	0	2.7	6
14/42/F	Pneumonia	31	1	3	2
15/26/F	Eclampsia	40	0	3.5	9
16/20/F	Sepsis	85	3	3.75	10
17/20/F	Multiple trauma	50	0	3.7	18
18/47/M	Pneumonia	58	4	3.5	9
19/55/F	Sepsis	70	3	3.75	11
20/28/F	Gastric aspiration	56	2	3.25	21
21/39/F	Gastric aspiration	38	0	3	11
22/31/F	Pneumonia	51	2	3.5	9
23/33/M	Pneumonia	60	3	3.25	10
24/33/F	Pneumonia	62	4	3	20
25/60/M	Sepsis	49	2	3	10
26/47/M	Pneumonia	56	5	3.75	6
27/16/F	Pneumonia	35	1	4	1
28/21/M	Multiple trauma	36	1	4	4
29/43/F	Pneumonia	48	4	3.33	5
30/29/F	Gastric aspiration	81	4	3.66	10
31/45/M	Pneumonia	38	0	3.33	7
32/39/M	Gastric aspiration	55	1	3.33	13
33/58/M	Pneumonia	41	0	3.33	10
34/38/F	Sepsis	54	4	4	3
35/21/F	Pneumonia	35	1	3.75	2
36/39/F	Eclampsia+aspiration	54	1	3.25	1
Mean 35.3, F=20	Primary=23	46.8	1.8	3.4	8.2
SD 12.4, M=16	Secondary=11	14.2	1.6	0.4	5.5

*Calculated with the same definitions as Gattinoni et al.¹⁹

†Duration of mechanical ventilation reflects total duration before admission.

Mean PaO₂ was 75.3±31.3 mm Hg at an FIO₂ of 1 for a mean PEEP level of 12.3±3.4 cm H₂O. Mean PaCO₂ was 50.7±13.7 mm Hg. PIP and mean airway pressures were elevated (39.7±8 and 22.1±4.9 cm H₂O, respectively). Quasistatic pulmonary compliance was 26.9±10.4 mL/cm H₂O. Fifty-five percent of the patients had barotraumatic lesions.

Response to TO

According to their response to TO, the population was separated in two groups named “responders” and “nonresponders” (see “Materials and Methods” section). Considering the severity criteria on arrival in the two groups, the only differences concerned MSOF score and PaCO₂ values, which were initially

higher in nonresponders (Table 2). It should be noted that Murray score, lung mechanic parameters, PaO₂, PEEP level, and duration of mechanical ventilation on arrival did not differ.

Responders Group (n=19): Nineteen of 36 patients were considered as responders according to the defined criteria and were maintained under conventional ventilation. All these patients improved their gas exchange within 2 days. In addition to PaO₂/FIO₂ increase, PaCO₂, PEEP, and mean airway pressures did not change and PIP decreased (Fig 2). This improvement seen in the first 2 days did not correspond to the best values observed, since the conditions of some patients continued to improve in the following days. Conversely, despite an initial

Table 2—Initial Severity Criteria on Arrival in Responders and Nonresponders to TO*

	Responders (n=19)	Nonresponders (n=17)
Age, yr	35.5±12.4	35.1±12.7
SAPS II	43±14	51±14
MSOF	1.3±1.3	2.4±1.7 [†]
Murray score	3.4±0.5	3.5±0.3
Duration of mechanical ventilation, d	6.7±4.3	9.9±6.2
PaO ₂ (FIO ₂ =1)	76.8±34.9	73.6±27.7
PaCO ₂	47.1±13.6	54.8±13.1 [†]
PIP	38.4±6.1	41.1±9.8
Mean Paw	22.3±5	22±5
PEEP	13.1±3.6	11.4±3.1
Cqs	29.6±12.3	23.7±6.9
Barotrauma, %	53	59

*Paw=airway pressure; Cqs=quasi-static lung compliance.

[†]p<0.05 (mean±SD).

improvement, the conditions of two patients of this group deteriorated so severely at day 5 and day 6 that an ECCO₂R-LFPPV was decided, with good success. In this group, four patients died (21%) after the third week because of MOSF (n=2), septic shock (n=1), and hemoptysis (n=1).

Nonresponders Group (n=17): Seventeen patients of the initial population (47%) did not improve their PaO₂/FIO₂ during TO during the first 2 days, as shown in Figure 2 (nonresponders). In addition, PaCO₂, PIP, airway pressure, and PEEP levels remained high despite maximum TO, leading us to discuss an indication of ECCO₂R-LFPPV. Applying the contraindication criteria described in the "Materials and Methods" section, ECCO₂R-LFPPV was not possible for nine patients (53% of nonresponders), all who died from different causes (78% with MSOF). For the remaining eight patients, ECCO₂R-LFPPV was performed, and a high mortality rate (six of eight patients) was observed during the procedure. One patient died of persistent hypoxemia and barotraumatic lesions; other causes of deaths were three hemorrhagic complications (one intracerebral hemorrhage and two pulmonary/pleural bleedings), and evolution of an initially severe MSOF in two patients presenting on arrival with a score of 4 and 5, respectively. For the two surviving patients, ECCO₂R-LFPPV technique was maintained during 3 days. Therefore, overall mortality in nonresponders was 88%, compared with a 21% mortality rate in responders (p<0.0001).

Univariate and Multivariate Analysis

Table 3 summarizes the variables analyzed in relation to outcome. Considering the initial severity criteria on arrival in responders and nonresponders

to TO listed in Table 2, univariate analysis revealed that, in addition to a favorable response to TO, SAPS II, MSOF, previous duration of mechanical ventilation on arrival, and PaCO₂ were associated with an adverse outcome. Age, Murray score, PaO₂, airway pressures, lung compliance, and existence of barotrauma lesions were not univariate correlates of adverse outcome.

Stepwise logistic regression analysis was then performed, and the response to TO appeared as the most predictive factor for survival. Then SAPS II was selected as the second independent predictor. None of the other parameters tested fulfilled the selection criteria of the multivariate procedure.

DISCUSSION

ARDS remains one of the most challenging organ failures to assist, even using sophisticated modes of mechanical ventilation.^{5,12,36} Since the initial definition by Ashbaugh et al¹ in 1967, the improvement in outcome remains debated, even if mortality is probably lower than previously reported.³ The absence of a clear amelioration in the outcome may result from different causes. First, the definition is large (although a new definition has been recently attempted²) and concerns different abnormalities and mechanisms; second, the underlying conditions of the patients play a crucial role; third there is no widely accepted consensus on the therapeutic strategy. During the past decade, the concept of MOSF has become increasingly crucial in the prognosis of severely ill patients, including those with ARDS. In addition, the better understanding of the pathophysiologic condition had led us to take into account different aspects of the lung lesion mechanisms, especially those induced by mechanical ventilation itself.^{37,38} Moreover, to our knowledge, no data have clearly demonstrated that a symptomatic correction of hypoxemia results in improved outcome of such patients.

Using the approach of the response to different therapeutic strategies aimed at correcting hypoxemia while limiting iatrogenic lung lesions, one could expect to obtain a useful classification of the patients in terms of (1) outcome, (2) pathophysiology, and (3) selection of subgroups of severely ill patients for whom future controlled trials or sophisticated assistance techniques could be tested. In the present study, despite a relatively small number of patients, a combination of techniques was used to improve hypoxemia, and the response to such a strategy discriminated responders from nonresponders. The absence of any response to TO was frequently




FIGURE 2. Evolution of gasometric and ventilatory parameters (mean \pm SE) in "responders" and "nonresponders" on arrival and during the first 48 h of TO. Mean Paw = mean airway pressure. Asterisk indicates $p < 0.05$ compared with value on arrival in the same group; dagger, $p < 0.05$ between responders and nonresponders.

associated with a more severe MOSF, reflecting the severity of the pulmonary and systemic inflammation.

The population selected in the present study was similar, considering Murray score,²¹ to recently published series,^{32,39} with a comparable global mortality rate.^{3,32,40} Mortality was mainly due to MOSF, sepsis, and ECCO₂R adverse effects, but rarely to respiratory failure *per se*, and, except for SAPS II, it was impossible to predict outcome based on initial data upon arrival in the ICU. In the present study, the TO used included various individual elements such as inhaled NO, prone position, or permissive hypercarbia. Considering the difficulty in conducting prospective controlled studies on the benefit of each of these individual therapies on outcome, we observed that the response to TO was the strongest discriminant criterion for outcome.

Using this TO strategy, we observed that half of the patients improved in terms of PaO₂/FIO₂. The follow-up of this group showed a significantly better prognosis than in nonresponders. The discussion of these results can only be speculative. First, comparison of the two groups suggests that nonresponders were more severe in terms of inflammatory reaction, since the incidence of organ failure was higher. In addition, their lung disease might also have been worse since PaCO₂ was higher for a similar ventilation. Although gas exchange function is not the main parameter to assess severity in ARDS, its improvement can be viewed as a dynamic means to evaluate the severity of the underlying inflammatory disease. Finally, although it has not been clearly demonstrated that hypoxia might precipitate organ failure, these consequences might be limited by hypoxemia correction. In the present population, severe hypoxia

Table 3—Univariate and Multivariate Analysis of Outcome

	Deceased Patients (n=19)	Surviving Patients (n=17)	p Value Univariate Analysis	p Value Multivariate Analysis
Responders to TO, %	21.1	88.2	<0.05	<0.05
Age, yr	37.4±13	32.9±11.6	NS	NS
SAPS II	54.1±13.6	38.6±9.8	<0.05	<0.05
MSOF	2.5±1.6	1±1.1	<0.05	NS
Murray score	3.5±0.3	3.3±0.5	NS	NS
Duration of mechanical ventilation, d	10.5±5.3	5.6±4.5	<0.05	NS
PaO ₂ (FIO ₂ =1)	74.5±24.8	76.2±38	NS	NS
PaCO ₂	57.4±15.6	43.2±5.1	<0.05	NS
PIP	41.4±8.9	37.8±6.7	NS	NS
Mean Paw	23.6±4.8	20.5±4.6	NS	NS
PEEP	12.7±3.3	11.8±3.6	NS	NS
Cqs	26.1±10.2	27.7±11	NS	NS
Barotrauma, %	52.6	58.8	NS	NS

*See Table 2 for explanation of abbreviations. NS=not significant.

was the cause of death for only two patients. Such a proportion is low, but not negligible, and promotes the idea of a low threshold of hypoxia tolerable by the organism. Above this level (around 60 mm Hg), the benefit for tissues of a PaO₂ increase is still debated. Experimental data suggest, nevertheless, that in addition to the metabolic aspects, hypoxia worsens the immunologic response and stimulates cytokines release.^{41,42} Conversely, the reoxygenation might also induce some tissue damage.^{43,44} It is then difficult to conclude about the role of the correction of a hypoxia as a prognosis factor rather than a marker of severity. Accordingly MSOF, rather than acute respiratory failure, remains the main cause of mortality in ARDS despite efficiency of supportive therapy.⁴⁵⁻⁴⁷

Following the above discussion, the use of ECCO₂R as a technique to improve PaO₂ and to limit induced-lung lesions is questionable. The present data suggest again that the benefit of such technique is influenced by the origin group, *ie*, responders or nonresponders. Two patients in the responders group were treated by ECCO₂R because of a secondary deterioration and survived. Conversely, in nonresponders, survival rate when ECCO₂R was performed was only 25%. Despite these very small numbers of patients, it is important to note that all patients with a contraindication to ECCO₂R died. This might suggest that ECCO₂R could be helpful as a last therapeutic support in selected patients failing to respond to TO, although the efficiency of this technique is still limited by its intrinsic complications.

The ability of the lung to recover structurally and functionally after a severe insult seems an important prognosis factor in ARDS, but assessment of the potential reversibility of the pulmonary injury is difficult. Gasometric or pulmonary mechanics pa-

rameters are poor predictive factors. In contrast, the response to a TO strategy appeared, using stepwise logistic regression, as the most predictive factor for mortality. Such an approach could also be used in the future to select subgroups of patients for investigational trials.

ACKNOWLEDGMENTS: The authors wish to thank Dr. Eric Vicaut for his statistical advice.

REFERENCES

- 1 Ashbaugh DG, Bigelow DB, Petty TL, et al. Acute respiratory distress in adults. *Lancet* 1967; 2:319-23
- 2 Bernard G, Artigas A, Brigham K, et al. The American-European Consensus Conference on ARDS: definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994; 149:818-24
- 3 Schuster D: What is the mortality of ARDS? In: Vincent J, ed. *Yearbook of intensive care and emergency medicine*. Berlin: Springer-Verlag, 1995; 59-68
- 4 Feihl F, Perret C. Permissive hypercapnia: how permissive should we be? *Am J Respir Crit Care Med* 1994; 150:1722-37
- 5 Hickling K. Ventilatory management of ARDS: can it affect the outcome? *Intensive Care Med* 1990; 16:219-26
- 6 Hickling K, Walsh J, Henderson S, et al. Low mortality rate in adult respiratory distress syndrome using low-volume, pressure-limited ventilation with permissive hypercapnia: a prospective study. *Crit Care Med* 1994; 22:1568-78
- 7 Tuxen DV. Permissive hypercapnic ventilation. *Am J Respir Crit Care Med* 1994; 150:870-74
- 8 Hudson L. Fluid management strategy in acute lung injury. *Am Rev Respir Dis* 1992; 145:988-89
- 9 Simmons R, Berdine G, Seidenfeld J, et al. Fluid balance and the adult respiratory distress syndrome. *Am Rev Respir Dis* 1987; 135:924-29
- 10 Garzia F, Todor R, Salea T. Continuous arteriovenous hemofiltration countercurrent dialysis (CAVH-D) in acute respiratory failure (ARDS). *J Trauma* 1991; 31:1277-85
- 11 Marini J, Kelsen S. Re-targeting ventilatory objectives in adult respiratory distress syndrome: new treatment prospects—persistent questions. *Am Rev Respir Dis* 1992; 146:2-3
- 12 Amato M, Barb s C, Medeiros D, et al. Beneficial effects of the 'open lung approach' with low distending pressures in

- acute respiratory distress syndrome: a prospective randomized study on mechanical ventilation. *Am J Respir Crit Care Med* 1995; 152:1835-46
- 13 Rossaint R, Falke K, Lopez F, et al. Inhaled nitric oxide for the adult respiratory distress syndrome. *N Engl J Med* 1993; 328:399-405
- 14 Rossaint R, Gerlach H, Schmidt-Ruhnke H, et al. Efficacy of inhaled nitric oxide in patients with severe ARDS. *Chest* 1995; 107:1107-15
- 15 Zapol W, Rimar S, Gillis N, et al. Nitric oxide and the lung. *Am J Respir Crit Care Med* 1994; 149:1375-80
- 16 Levy B, Bollaert P, Bauer P, et al. Therapeutic optimization including inhaled nitric oxide in adult respiratory distress syndrome in a polyvalent intensive care unit. *J Trauma* 1995; 38:370-74
- 17 Payen D, Gatecel C, Plaisance P. Almitrine effect on nitric oxide inhalation in adult respiratory distress syndrome [letter]. *Lancet* 1993; 341:1664
- 18 Wysocki M, Delclaux C, Roupie E, et al. Additive effect on gas exchange of inhaled nitric oxide and intravenous almitrine bismesylate in the adult respiratory distress syndrome. *Intensive Care Med* 1994; 20:254-59
- 19 Gattinoni L, Pesenti A, Mascheroni D, et al. Low-frequency positive-pressure ventilation with extracorporeal CO₂ removal in severe acute respiratory failure. *JAMA* 1986; 256: 881-86
- 20 Kinsella J, Abman S. Inhalational nitric oxide therapy for persistent pulmonary hypertension of the newborn. *Pediatrics* 1993; 91:997-98
- 21 Murray J, Matthay M, Luce J, et al. An expanded definition of the adult respiratory distress syndrome. *Am Rev Respir Dis* 1988; 138:720-23
- 22 Zapol W, Snider M, Hill D, et al. Extracorporeal membrane oxygenation in severe acute respiratory failure: a randomized prospective study. *JAMA* 1979; 242:2193-96
- 23 Le Gall J, Lemeshow S, Saulnier F. A new simplified acute physiology score (SAPS II) based on a European/North American multicenter study. *JAMA* 1993; 270:2957-63
- 24 Marcy T. Barotrauma: detection, recognition and management. *Chest* 1993; 104:578-84
- 25 Lynch J, Mhyre J, Dantzker D. Influence of cardiac output on intrapulmonary shunt. *J Appl Physiol* 1979; 46:315-21
- 26 Dantzker D, Lynch J, Weg J. Depression of cardiac output is a mechanism of shunt reduction in the therapy of acute respiratory failure. *Chest* 1980; 77:636-42
- 27 Ravenscraft S, Burke W, Nahum A, et al. Tracheal gas insufflation augments CO₂ clearance during mechanical ventilation. *Am Rev Respir Dis* 1993; 148:345-51
- 28 Nakos G, Zakinthinos S, Kotanidou A, et al. Tracheal gas insufflation reduces the tidal volume while PaCO₂ is maintained constant. *Intensive Care Med* 1994; 20:407-13
- 29 Marshall B, Hanson C, Frasch F, et al. Role of hypoxic vasoconstriction in pulmonary gas exchange and blood flow distribution. *Intensive Care Med* 1994; 20:379-89
- 30 Gottschall E, Feryak S, Wuertemberger G, et al. Almitrine mimics hypoxic vasoconstriction in isolated rat lungs. *Am J Physiol* 1992; 263:H383-91
- 31 Reyes A, Roca J, Rodriguez-Roisin R, et al. Effects of almitrine on ventilation-perfusion distribution in adult respiratory distress syndrome. *Am Rev Respir Dis* 1988; 137: 1062-67
- 32 Morris A, Wallace C, Menlove R, et al. Randomized clinical trials of pressure-controlled inverse ratio ventilation and extracorporeal CO₂ removal for adult respiratory distress syndrome. *Am J Respir Crit Care Med* 1994; 149:295-305
- 33 Dixon W. BMDP statistical software. Berkeley, Calif: University of California Press, 1985
- 34 Draper N, Smith H. Regression analysis. New York: John Wiley, 1967
- 35 Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 1959; 22:719-48
- 36 Kollef M, Schuster D. The acute respiratory distress syndrome. *N Engl J Med* 1995; 332:27-37
- 37 Tsuno K, Miura K, Takeya M, et al. Histopathologic pulmonary changes from mechanical ventilation at high peak airway pressures. *Am Rev Respir Dis* 1991; 143:1115-20
- 38 Dreyfuss D, Saumon G. Role of tidal volume, FRC, and end-inspiratory volume in the development of pulmonary edema following mechanical ventilation. *Am Rev Respir Dis* 1993; 148:1194-1203
- 39 Brunet F, Mira J, Belghith M, et al. Extracorporeal dioxide removal technique improves oxygenation without causing overinflation. *Am J Respir Crit Care Med* 1994; 149:1557-62
- 40 Lewandowski K, Metz J, Deutschmann C, et al. Incidence, severity, and mortality of acute respiratory failure in Berlin, Germany. *Am J Respir Crit Care Med* 1995; 151:1121-25
- 41 Ertel W, Morrison M, Ayala A, et al. Hypoxemia in the absence of blood loss or significant hypotension causes inflammatory cytokine release. *Am J Physiol* 1995; 269:160-66
- 42 Karakurum M, Shreeniwas R, Chen J, et al. Hypoxic induction of interleukin-8 gene expression in human endothelial cells. *J Clin Invest* 1994; 93:1564-70
- 43 Kennedy T, Rao N, Hopkins C, et al. Role of reactive oxygen species in reperfusion injury of the rabbit lung. *J Clin Invest* 1989; 83:1326-35
- 44 Murata T, Nakazawa H, Mori I, et al. Reperfusion after a 2-hour period of pulmonary artery occlusion causes pulmonary necrosis. *Am Rev Respir Dis* 1992; 146:1048-53
- 45 Bell R, Coalson J, Smith J, et al. Multiple organ system failure and infection in adult respiratory distress syndrome. *Ann Intern Med* 1983; 99:293-98
- 46 Bone R, Balk R, Slotman G, et al. Adult respiratory distress syndrome: sequence and importance of development of multiple organ failure. *Chest* 1992; 101:320-26
- 47 Gattinoni L, Brazzi L, Pesenti A. Extracorporeal carbon dioxide removal in ARDS. In: Marini J, Roussos C, eds. Ventilatory failure. Berlin: Springer-Verlag, 1991; 308-17