

ADDITIVE SUMSET SIZES WITH TETRAHEDRAL DIFFERENCES

MELVYN B. NATHANSON

ABSTRACT. Experimental calculations suggest that the h -fold sumset sizes of 4-element sets of integers are concentrated at h numbers that are differences of tetrahedral numbers. In this paper it is proved that these “popular” sumset sizes exist and explicit h -adically defined sets are constructed for each of these numbers.

1. THE SUMSET SIZE PROBLEM

The h -fold sum of a set A of integers is the set of all sums of h not necessarily distinct elements of A . The core problem of additive number theory is to understand h -fold sumsets.

If A is a finite set of k integers, then hA is a finite set and

$$(1) \quad h(k-1) + 1 \leq |hA| \leq \binom{h+k-1}{k-1}.$$

We have $|hA| = h(k-1) + 1$ if and only if A is an arithmetic progression of length k , and $|hA| = \binom{h+k-1}{k-1}$ if and only if A is a B_h -set, that is, a set such that every integer in the sumset hA has a unique representation (up to permutation of the summands) as a sum of h not necessarily distinct elements of A .

The *integer interval* defined by real numbers u and v is the set

$$[u, v] = \{n \in \mathbf{Z} : u \leq n \leq v\}.$$

The *integer part* of the real number u is denoted $[u]$.

Let $\mathcal{R}_{\mathbf{Z}}(h, k)$ be the set of h -fold sumset sizes of sets of size k , that is,

$$\mathcal{R}_{\mathbf{Z}}(h, k) = \{|hA| : A \subseteq \mathbf{Z} \text{ and } |A| = k\}.$$

Inequality (1) implies

$$(2) \quad \mathcal{R}_{\mathbf{Z}}(h, k) \subseteq \left[h(k-1) + 1, \binom{h+k-1}{k-1} \right].$$

Not every possible sumset size is actually the size of a sumset. For example, relation (2) gives

$$\mathcal{R}_{\mathbf{Z}}(3, 3) \subseteq [7, 10].$$

We have

$$3\{0, 1, 2\} = \{0, 1, 2, 3, 4, 5, 6\} \quad \text{and} \quad |3\{0, 1, 2\}| = 7$$

Date: July 16, 2025.

2000 Mathematics Subject Classification. 11B13, 11B05, 11B75, 11P70, 11Y16, 11Y55.

Key words and phrases. Sumset, sumset size, popular sumset size, tetrahedral number, additive number theory, combinatorial number theory.

Supported in part by PSC-CUNY Research Award Program grant 66197-00 54.

$$\begin{aligned} 3\{0, 1, 3\} &= \{0, 1, 2, 3, 4, 5, 6, 7, 9\} & \text{and} & & |3\{0, 1, 3\}| &= 9 \\ 3\{0, 1, 4\} &= \{0, 1, 2, 3, 4, 5, 6, 8, 9, 12\} & \text{and} & & |3\{0, 1, 4\}| &= 10 \end{aligned}$$

and so

$$\{7, 9, 10\} \subseteq \mathcal{R}_{\mathbf{Z}}(3, 3).$$

However, there exists no set A of integers with $|A| = 3$ and $|3A| = 8$ (Nathanson [?]). Thus,

$$\mathcal{R}_{\mathbf{Z}}(3, 3) = \{7, 9, 10\}.$$

This example motivates the following problem: For all positive integers h and k , determine the full range of sumset sizes of h -fold sums of sets of k integers, that is, compute the set $\mathcal{R}_{\mathbf{Z}}(h, k)$.¹ For all h and k , we have

$$\mathcal{R}_{\mathbf{Z}}(h, 1) = \{1\} \quad \text{and} \quad \mathcal{R}_{\mathbf{Z}}(1, k) = \{k\}.$$

Sets A and B are *affinely equivalent* if there exist numbers $\lambda \neq 0$ and μ such that $\lambda * A + B = \{\lambda a + \mu : a \in A\}$. If A and B are affinely equivalent, then $|hA| = |hB|$ for all positive integers h . Every finite set A of integers is affinely equivalent to a set B with $\min B = 0$ and $\gcd(B) = 1$. In particular, every set of size 2 is affinely equivalent to the set $\{0, 1\}$. It follows that

$$\mathcal{R}_{\mathbf{Z}}(h, 2) = \{h + 1\}.$$

Erdős and Szemerédi [?] stated that

$$\mathcal{R}_{\mathbf{Z}}(2, k) = \left[2k - 1, \binom{k+1}{2} \right].$$

(A proof is in [?].) Thus, the unsolved problem is to determine $\mathcal{R}_{\mathbf{Z}}(h, k)$ for $h \geq 3$ and $k \geq 3$.

A first step is to fix a positive integer k and find the possible sizes of h -fold sums of sets of size k . Recall that the j th *triangular number* $f_2^j = \binom{j+1}{2}$ is the sum of the first j positive integers. The j th *tetrahedral number* $f_3^j = \binom{j+2}{3}$ is the sum of the first j triangular numbers (Dickson [?]). For $k = 3$, Nathanson [?] proved that

$$\mathcal{R}_{\mathbf{Z}}(h, 3) = \left\{ \binom{h+2}{2} - \binom{i_0+1}{2} : i_0 \in [0, h-1] \right\}.$$

Thus, every sumset size of a 3-element set is of the form $f_2^{h+1} - f_2^{i_0}$, that is, a difference of triangular numbers. For $k \geq 4$, the problem is still open. Numerical experiments (Nathanson [?] and O'Bryant [?]) suggest that, for $k = 4$, the “most popular” sumset sizes are the integers

$$f_3^{h+1} - f_3^{i_0} = \binom{h+3}{3} - \binom{i_0+2}{3}$$

for $i_0 \in [0, h-1]$. These are the differences between the tetrahedral number $f_3^{h+1} = \binom{h+3}{3}$, which is also the size of a 4-element B_h -set, and the h consecutive tetrahedral numbers $f_3^0, f_3^1, \dots, f_3^{h-1}$. It had been an open problem to decide if the integers $f_3^{h+1} - f_3^{i_0}$ are, indeed, sumset sizes for all $h \geq 2$ and $i_0 \in [0, h-1]$. The goal of this paper is to prove that these sumset sizes do exist for all h and i_0 , that is,

$$\left\{ \binom{h+3}{3} - \binom{i_0+2}{3} : i_0 \in [0, h-1] \right\} \subseteq \mathcal{R}_{\mathbf{Z}}(h, 4)$$

¹There is the analogous problem in every additive abelian group or semigroup G : Determine the set $\mathcal{R}_G(h, k)$ of the sizes of h -fold sums of k -element subsets of G .

and to construct explicit h -adically defined sets with exactly these sumset sizes.

For related work on sumset size problems in additive number theory, see [?]-[?].

2. A FAMILY OF h -ADIC SETS

Theorem 1. *For all $h \geq 2$ and $i_0 \in [0, h-1]$, let*

$$c = (h+1-i_0)(h+1).$$

The set

$$A = \{0, 1, h+1, c\}$$

satisfies $|A| = 4$ and

$$|hA| = \binom{h+3}{3} - \binom{i_0+2}{3}.$$

Proof. The set

$$B = \{0, 1, h+1\}$$

is a B_h -set and so B is a B_{h-i} -set for all $i \in [0, h-1]$ and

$$|(h-i)B| = \binom{h-i+2}{2}.$$

Let $0B = \{0\}$.

We have

$$A = B \cup \{c\} = \{0, 1, h+1, (h+1-i_0)(h+1)\}.$$

The inequality $h+1-i_0 \geq 2$ implies $|A| = 4$.

If $i_0 = 0$, then $c = (h+1)^2$ and

$$A = \{0, 1, h+1, (h+1)^2\}.$$

The uniqueness of $(h+1)$ -adic representations implies that A is a B_4 -set and so

$$|hA| = \binom{h+3}{3} = \binom{h+3}{3} - \binom{i_0+2}{3}.$$

Let $i_0 \in [1, h-1]$. We decompose the sumset hA as follows:

$$hA = \bigcup_{i=0}^h ((h-i)B + ic) = \bigcup_{i=0}^h L_i$$

where

$$\begin{aligned} L_i &= (h-i)B + ic \\ &= \bigcup_{j=0}^{h-i} ((h-i-j)(h+1) + [0, j]) + ic \\ (3) \quad &= \bigcup_{j=0}^{h-i} ((h + (h-i_0)i - j)(h+1) + [0, j]). \end{aligned}$$

We have

$$(4) \quad |L_i| = |(h-i)B + ic| = |(h-i)B| = \binom{h-i+2}{2}.$$

The set L_i is the union of $h-i+1$ pairwise disjoint integer intervals whose smallest elements are multiples of $h+1$ and whose lengths are at most h . It follows

that if $n \in L_i$ and $n = q(h+1) + r$ with $r \in [0, h]$, then $q = h + (h - i_0)i - j$ for some $j \in [0, h - i]$ and $r \in [0, j]$. Then L_i contains the integer interval $q(h+1) + [0, j]$.

For all $i \in [0, h - 1]$, we have

$$\min(L_i) = ic < (i+1)c = \min(L_{i+1})$$

and

$$\begin{aligned} \max(L_i) &= (h-i)(h+1) + ic \\ &< (h-i-1)(h+1) + (i+1)c \\ &= \max(L_{i+1}) \end{aligned}$$

and so the sets L_i “move to the right” as i increases from 0 to h . Moreover,

$$\max(L_i) < \min(L_{i+1})$$

if and only if

$$(h-i)(h+1) + ic < (i+1)c$$

if and only if

$$i > h - \frac{c}{h+1}$$

if and only if

$$i \geq 1 + \left\lceil h - \frac{c}{h+1} \right\rceil = i_0.$$

Thus, the sets L_i and L_j are disjoint if $i_0 \leq i < j \leq h$ and

$$\left| \bigcup_{i=i_0+1}^h L_i \right| = \sum_{i=i_0+1}^h |L_i| = \sum_{i=i_0+1}^h \binom{h-i+2}{2}.$$

Because the sets L_i move to the right, we have

$$\left(\bigcup_{i=0}^{i_0} L_i \right) \cap \left(\bigcup_{i=i_0+1}^h L_i \right) = \emptyset$$

and

$$\begin{aligned} |hA| &= \left| \bigcup_{i=0}^h L_i \right| = \left| \bigcup_{i=0}^{i_0} L_i \right| + \left| \bigcup_{i=i_0+1}^h L_i \right| \\ (5) \quad &= \left| \bigcup_{i=0}^{i_0} L_i \right| + \sum_{i=i_0+1}^h \binom{h-i+2}{2}. \end{aligned}$$

We shall compute $L_i \cap L_{i+t}$ for all $i \in [1, h-1]$ and $t \in [1, h-i]$. Relation (??) implies

$$L_{i+t} = \bigcup_{j=0}^{h-i-t} ((h + (h - i_0)(i+t) - j)(h+1) + [0, j]).$$

We have $q(h+1) \in L_i \cap L_{i+t}$ if and only if there exist $j_0 \in [0, h-i]$ and $j_t \in [0, h-i-t]$ such that

$$q = h + (h - i_0)i - j_0 = h + (h - i_0)(i+t) - j_t$$

if and only if

$$\begin{aligned} j_0 &= j_t - (h - i_0)t \\ &\in [0, h - i] \cap [-(h - i_0)t, h - i - t - (h - i_0)t] \\ &= [0, h - i - t - (h - i_0)t]. \end{aligned}$$

It follows that

$$(6) \quad L_i \cap L_{i+t} = \bigcup_{j=0}^{h-i-t-(h-i_0)t} ((h + (h - i_0)i - j)(h + 1) + [0, j])$$

and so

$$\begin{aligned} |L_i \cap L_{i+t}| &= \left| \bigcup_{j=0}^{h-i-t-(h-i_0)t} ((h + (h - i_0)i - j)(h + 1) + [0, j]) \right| \\ &= \sum_{j=0}^{h-i-t-(h-i_0)t} |(h + (h - i_0)i - j)(h + 1) + [0, j]| \\ &= \sum_{j=0}^{h-i-t-(h-i_0)t} (j + 1) \\ &= \binom{h - i - t - (h - i_0)t + 2}{2}. \end{aligned}$$

In particular,

$$(7) \quad |L_i \cap L_{i+1}| = \binom{i_0 + 1 - i}{2}.$$

Relation (??) also implies that, for $t \in [1, h - i]$,

$$L_i \setminus L_{i+t} = \bigcup_{j=h-i-t-(h-i_0)t+1}^{h-i} ((h + (h - i_0)i - j)(h + 1) + [0, j])$$

and so

$$L_i \setminus L_{i+1} \subseteq L_i \setminus L_{i+2} \subseteq \cdots \subseteq L_i \setminus L_h.$$

Therefore,

$$\begin{aligned} L_i \setminus \left(\bigcup_{t=1}^{i_0-i} L_{i+t} \right) &= L_i \cap \left(\bigcup_{t=1}^{i_0-i} L_{i+t} \right)^c = L_i \cap \left(\bigcap_{t=1}^{i_0-i} L_{i+t}^c \right) \\ &= \bigcap_{t=1}^{i_0-i} (L_i \cap L_{i+t}^c) = \bigcap_{t=1}^{i_0-i} (L_i \setminus L_{i+t}) \\ &= L_i \setminus L_{i+1}. \end{aligned}$$

The sets

$$L_i \setminus \left(\bigcup_{t=1}^{i_0-i} L_{i+t} \right)$$

are pairwise disjoint for $i \in [0, i_0]$ and

$$\bigcup_{i=0}^{i_0} L_i = L_{i_0} \cup \bigcup_{i=0}^{i_0-1} \left(L_i \setminus \bigcup_{t=1}^{i_0-i} L_{i+t} \right).$$

From (??) and (??), we obtain

$$\begin{aligned}
\left| \bigcup_{i=0}^{i_0} L_i \right| &= |L_{i_0}| + \sum_{i=0}^{i_0-1} \left| L_i \setminus \bigcup_{t=1}^{i_0-i} L_{i+t} \right| \\
&= |L_{i_0}| + \sum_{i=0}^{i_0-1} |L_i \setminus L_{i+1}| \\
&= |L_{i_0}| + \sum_{i=0}^{i_0-1} (|L_i| - |L_i \cap L_{i+1}|) \\
&= \sum_{i=0}^{i_0} |L_i| - \sum_{i=0}^{i_0-1} |L_i \cap L_{i+1}| \\
&= \sum_{i=0}^{i_0} \binom{h-i+2}{2} - \sum_{i=0}^{i_0-1} \binom{i_0+1-i}{2}.
\end{aligned}$$

Relation (??) gives

$$\begin{aligned}
|hA| &= \left| \bigcup_{i=0}^{i_0} L_i \right| + \sum_{i=i_0+1}^h \binom{h-i+2}{2} \\
&= \sum_{i=0}^h \binom{h-i+2}{2} - \sum_{i=0}^{i_0-1} \binom{i_0+1-i}{2} \\
&= \sum_{i=0}^h \binom{i+2}{2} - \sum_{i=0}^{i_0-1} \binom{i+2}{2} \\
&= \binom{h+3}{3} - \binom{i_0+2}{3}.
\end{aligned}$$

This completes the proof. \square

3. OPEN PROBLEMS

Problem 1. *This paper considers the special class of h -adically defined 4-element sets*

$$A = \{0, 1, h+1, h^2+h+1-p\}$$

with

$$p = 1 + (i_0 - 1)(h + 1)$$

and $i_0 \in [0, h-1]$. It is of interest to compute, for all $h \geq 3$ and all $p \in [0, h^2-1]$, the h -fold sumset sizes of the sets

$$A = \{0, 1, h+1, h^2+h+1-p\}.$$

Problem 2. *For all $h \geq 3$, compute the set of h -fold sumset sizes of the sets*

$$A = \{0, 1, a, b\}$$

for $2 \leq a \leq h$ and $a+1 \leq b \leq ha+1$.

Problem 3. *A next step is to determine the popular sumset sizes of 5-element sets of integers. The fundamental problem is to obtain a complete description of the sumset size set $\mathcal{R}_{\mathbf{Z}}(h, k)$ for all positive integers h and k , to explain the distribution*

of sumset sizes for fixed h and k , and to understand why some numbers cannot be sumset sizes. A solution to this problem might be called the Second Fundamental Theorem of Additive Number Theory.

REFERENCES

- [1] Leonard Eugene Dickson, *History of the Theory of Numbers*, Volume II, Carnegie Institute of Washington, 1920. Reprinted by Chelsea Publishing Company, New York, 1971.
- [2] P. Erdős and E. Szemerédi, On sums and products of integers, in: *Studies in Pure Mathematics*, Birkhäuser, Basel-Boston, 1983, pp. 213–218.
- [3] J. Fox, N. Kravitz, and S. Zhang, Finer control on relative sizes of iterated sumsets, preprint, 2025.
- [4] N. Hegyvári, On representation problems in the additive number theory, *Acta Math. Hungar.* 72 (1996), 35–44.
- [5] N. Kravitz, Relative sizes of iterated sumsets, *J. Number Theory* 272 (2025), 113–128.
- [6] M. B. Nathanson, Inverse problems for sumset sizes of finite sets of integers, *Fibonacci Quarterly* (2025), to appear. arXiv:2411.02365.
- [7] M. B. Nathanson, Problems in additive number theory, VI: Sizes of sumsets of finite sets, *Acta Math. Hungar.*, to appear. arXiv:2411.02365, 2024.
- [8] M. B. Nathanson, Explicit sumset sizes in additive number theory, arXiv: 2505.05329, 2025.
- [9] M. B. Nathanson, Compression and complexity for sumset sizes in additive number theory, arXiv:2505.20998.
- [10] M. B. Nathanson, Triangular and tetrahedral number differences of sumset sizes in additive number theory, arXiv:2506.15015.
- [11] K. O’Bryant, On Nathanson’s triangular number phenomenon, arXiv:2506.20836.
- [12] P. Péringuey and A. de Roton, A note on iterated sumsets races, arXiv:2505.11233.
- [13] V. Schinina, On the sumset of sets of size k , arXiv:2505.07679.

DEPARTMENT OF MATHEMATICS, LEHMAN COLLEGE (CUNY), BRONX, NY 10468
Email address: melvyn.nathanson@lehman.cuny.edu