

Spatiotemporal characteristics of solar resource and photovoltaic productivity over the Euro-Mediterranean area

A climate perspective

Claudia Gutiérrez

24th may 2019

¹Facultad de Ciencias Ambientales y Bioquímica
Universidad de Castilla-La Mancha, Toledo, Spain

Context and introduction

Energy transition

VRE

Objectives and methodology


Results

ΔPV by country


Conclusions

Context and introduction


Energy transition



Energy transition




Energy transition




Photovoltaic

- Increase in **photovoltaic (PV) capacity**
- Continuous growth in projected **trends**.
- Global increase led by China

Solar PV Global Capacity, by Country/Region, 2005–2015




Electricity systems features:




Electricity systems features:

- Demand and supply need to be **balanced**.




Electricity systems features:

- Demand and supply need to be **balanced**.
- Electricity systems are designed for centralized **conventional power plants**.




Electricity systems features:

- Demand and supply need to be **balanced**.
- Electricity systems are designed for centralized **conventional power plants**.
- **VRE**: variable renewable energy.



VRE: variable renewable energy


Balancing may occur at different time scales



VRE: variable renewable energy

- **Intermittency:** not synchronized with the demand


Balancing may occur at different time scales



VRE: variable renewable energy

- **Intermittency:** not synchronized with the demand
- Need of forecasting, planification and/or storage


Balancing may occurs at different time scales




VRE: variable renewable energy

- **Intermittency**: not synchronized with the demand
- Need of forecasting, planning and/or storage
- Variations **from short to long scales** (weather to climate)
 - **short**: operation
 - **medium**: planning, maintenance
 - **long**: resource assessment, financing, planning

Balancing may occur at different time scales



Long scales: links between climate and renewables



Variability sources on PV

- Solar resource is **variable** from short to long-term time scales.
- Main drivers of solar resource variability are **cloudiness** and **aerosols**.



Figure 1: Image from July 16th, 2003. MODIS. Credit:NASA

Need for long-term projections of **resource** **and PV potential**

Previous studies show a **discrepancy** between **GCMs** and **RCMs** surface solar radiation (SSR) over Europe:

- **Increase** projected by GCMs (*Wild et al. 2015, Solar Energy*).
- **Decrease** projected by RCMs (*Jerez et al. 2015, Nature Communications*).

Objectives and methodology

- **1** To illustrate the **inconsistency** between **GCM** and **RCM** projections and to attribute it to missing aerosols forcing.
- **2** To **deliver future projections of PV** potential production over Europe.

METHODS

- **1** To illustrate the **inconsistency** between **GCM** and **RCM** projections and to attribute it to missing aerosols forcing.
 - Use of well-chosen groups of GCM-RCM within the **Euro-CORDEX** ensemble.
 - **2021-2050** summer change in surface solar radiation, **SSR**, with respect of a reference period: **1971-2000**. Use of **RCP8.5** scenario.

GCM	RCM	Aerosols
CNRM-CM5	CCLM4-8-17	-
	ALADIN53	Szopa et al.
	RCA4	-
	CCLM4-8-17	-

- **2** To **deliver future projections of PV** potential production over Europe.

solaR

Parametric PV model. **SSR** from **RCM** as input -> POA and electrical performance. Implemented in R (O. Perpiñán, 2013).




Figure 2: Sscheme of a PV generator.

Results

GCM

RCM

ssr_gimp4.png

Figure 3: SSR change (1971/2000)-(2021/2050) (W/m^2)

GCM

RCM

ssr_gimp3.png

Figure 4: SSR change (1971/2000)-(2021/2050) (W/m^2)

- Only RCMs with evolving aerosols show the increase in SSR as GCMs.

GCM

RCM

clt_gimp3.png

Figure 5: CLT change (1971/2000)-(2021/2050) (%)

GCM

RCM

clt_gimp3.png

Figure 6: CLT change (1971/2000)-(2021/2050) (%)

- RCMs without evolving aerosols: CLT spatial pattern can explain SSR spatial pattern.
- CLT spatial pattern cannot explain SSR spatial pattern in models with evolving aerosols.

Mean changes 2021-2050

GCM	RCM	$\Delta SSR [W/m^2]$	$\Delta CLT [\%]$
CNRM-CM5		9.9	0.5
	CCLM4-8-17	-2.4	-0.8
	ALADIN53	12.6	0.3
	RCA4	-2.6	0.2
EC-EARTH		5.6	-0.3
	CCLM4-8-17	-2.7	-0.9
	RACMO22E	4.8	0.5
	RCA4	-2.1	0.1

Table 2: Spatial changes in SSR and CLT

AOD mean changes 2021-2050




Figure 7: AOD change (1971/2000)-(2021/2050)

- Spatial pattern of ΔAOD similar to ΔSSR when evolving aerosols considered.
- Higher correlation of SSR with AOD than with CLT.

GCM	RCM	ΔAOD	$P_{SSR,CLT}$	$P_{SSR,AOD}$
CNRM-CM5	CCLM4-8-17	-	-0.7	-
	ALADIN53	-0.2	-0.2	-0.9
	RCA4	-	-0.8	-
EC-EARTH	CCLM4-8-17	-	-0.8	-
	RACMO22E	-0.1	-0.3	-0.6
	RCA4	-	-0.8	-

Δ PV relative JJA mean by country

Figure 8: Relative change in PV potential [%]

Figure 9: Relative change in PV potential [%]

- Decrease for models with no-evolving aerosols.

Figure 10: Relative change in PV potential [%]

- Decrease for models with no-evolving aerosols.
- Increase for models with evolving aerosols.
- Central-Europe is the most impacted area.

Conclusions

Conclusions

- For the mid century, an **increase** in **photovoltaic potential** is projected over Europe when the **evolution** of **aerosols** over the area is considered.
- The **magnitude** depends on the country and the models.
- The most impacted areas are in Central-Europe, with an important potential increase of more than **10%** but large uncertainty between models.

Perspectives

- A **robust answer** is needed in order to deliver key messages for the solar industry.
- The FPS-aerosols could help to understand uncertainties and develop better projections for energy purposes.

Thank you for your attention.

claudia.gutierrez@uclm.es