

# Numerical Methods

## PDEs

### 1 Non Matlab exercises

1. From Lecture notes we know that the second derivative is well approximated by the following finite difference:

$$\frac{d^2 u}{dx^2}(0) \simeq \frac{u(-h) - 2u(0) + u(h)}{h^2}.$$

How would you derive this formula if you did not know it?

2. Consider the following PDE:

$$u_t = u_x$$

where  $x \in [0, 1]$ ,  $t \geq 0$ . The boundary condition is periodic:

$$u|_{x=0} = u|_{x=1}.$$

The initial condition is  $u|_{t=0} = \sin(\pi x) + 10 \sin(3\pi x)$ .

- (a) Find the exact solution.
- (b) Design the forward Euler scheme.
- (c) Analyse the stability of the Euler scheme.
- (d) Analyse the stability of the leap-frog scheme:

$$U_j^{n+1} = U_j^{n-1} + \frac{\Delta t}{\Delta x} (U_{j+1}^n - U_{j-1}^n).$$

### 2 Matlab exercises

1. Consider the following PDE:

$$u_t = u_{xx}$$

where  $x \in [0, 1]$ ,  $t \geq 0$ . The boundary condition is

$$u|_{x=0} = u|_{x=1} = 0.$$

The initial condition is  $u|_{t=0} = \sin(\pi x) + 10 \sin(3\pi x)$ .

- (a) Find the exact solution of this problem. Hint: use separation variables technique. (You do not need MATLAB for this.)
- (b) The Euler scheme for this equation is

$$U_j^{n+1} = U_j^n + \frac{\Delta t}{\Delta x^2}(U_{j+1}^n - 2U_j^n + U_{j-1}^n)$$

for  $0 < j < J$  and

$$U_0^{n+1} = U_J^{n+1} = 0.$$

Here  $J + 1$  is the number of grid points in  $x$  direction.

The theory guaranties that the approximate solution  $U$  converges to the exact solution when  $\Delta x, \Delta t \rightarrow 0$ , provided the stability condition holds:

$$\frac{\Delta t}{\Delta x^2} < \frac{1}{2}.$$

Implement the Euler scheme and numerically find an approximate solution for  $t \in [0, 0.1]$ . Plot a graph of the solution.

- (c) Investigate what happens when the stability condition is violated.
- (d) Implement the Implicit Euler scheme.
- (e) Implement the Crank-Nicolson scheme.
- (f) Investigate how the error converges to zero when  $\Delta x, \Delta t \rightarrow 0$  for the all three schemes.

2. Consider the following PDE from the Non matlab section:

$$u_t = u_x$$

where  $x \in [0, 1]$ ,  $t \geq 0$ . The boundary condition:

$$u|_{x=0} = u|_{x=1}.$$

The initial condition is  $u|_{t=0} = \sin(\pi x) + 10 \sin(3\pi x)$ .

- (a) Implement the forward Euler scheme.
- (b) Implement the leap-frog scheme:

$$U_j^{n+1} = U_j^{n-1} + \frac{\Delta t}{\Delta x}(U_{j+1}^n - U_{j-1}^n).$$