

CPSC 340 Tutorial

Week 1

September 13, 2021

Notation and Gradients

- Define: \mathbb{R}^n , $x^\top Ax$, $\|x\|$, $\nabla f(x)$

Notation and Gradients

- Define: \mathbb{R}^n , $x^\top Ax$, $\|x\|$, $\nabla f(x)$
- If $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$, $A \in \mathbb{R}^{n \times m}$, find the dimensions of:

$$x^\top A, \quad x^\top Ay, \quad xx^\top$$

Notation and Gradients

- Define: \mathbb{R}^n , $x^\top Ax$, $\|x\|$, $\nabla f(x)$
- If $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$, $A \in \mathbb{R}^{n \times m}$, find the dimensions of:

$$x^\top A, \quad x^\top Ay, \quad xx^\top$$

- If $a, x \in \mathbb{R}^n$, compute gradients of:

$$f(x) = a^\top x$$

$$f(x) = \log(a^\top x)$$

Notation and Gradients

- Define: \mathbb{R}^n , $x^\top Ax$, $\|x\|$, $\nabla f(x)$
- If $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$, $A \in \mathbb{R}^{n \times m}$, find the dimensions of:

$$x^\top A, \quad x^\top Ay, \quad xx^\top$$

- If $a, x \in \mathbb{R}^n$, compute gradients of:

$$f(x) = a^\top x$$

$$f(x) = \log(a^\top x)$$

- Sanity check: If $x \in \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $\nabla f(x) \in \mathbb{R}^n$

Finite Difference Approximations

- Recall the definition of the derivative:

$$\frac{df}{dx} = \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon) - f(x)}{\epsilon}$$

Finite Difference Approximations

- Recall the definition of the derivative:

$$\frac{df}{dx} = \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon) - f(x)}{\epsilon}$$

- We can approximate the derivative by choosing small ϵ (say $\epsilon = 1e - 4$) and ignoring the limit:

$$\frac{df}{dx} \approx \frac{f(x + \epsilon) - f(x)}{\epsilon}$$

Finite Difference Approximations

- Recall the definition of the derivative:

$$\frac{df}{dx} = \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon) - f(x)}{\epsilon}$$

- We can approximate the derivative by choosing small ϵ (say $\epsilon = 1e - 4$) and ignoring the limit:

$$\frac{df}{dx} \approx \frac{f(x + \epsilon) - f(x)}{\epsilon}$$

- This can be extended to the gradient case by computing each element of the gradient as above. If $x \in \mathbb{R}^n$:

$$(\nabla f(x))_i \approx \frac{f(x_1, \dots, x_i + \epsilon, \dots, x_n) - f(x)}{\epsilon}$$

Python and Numpy Basics (Demo)
