### CPSC 340 Tutorial 2

September 20, 2021

### Table of Contents

Probability Review

References

### Table of Contents

Probability Review

References

► An **event** is an outcome of a random process that may or may not happen

- ► An **event** is an outcome of a random process that may or may not happen
  - Notation: *A*, *B*, etc.

- ► An **event** is an outcome of a random process that may or may not happen
  - ► Notation: A, B, etc.
  - Example: Rolling a 6 on a 6-sided die

- An event is an outcome of a random process that may or may not happen
  - ▶ Notation: *A*, *B*, etc.
  - Example: Rolling a 6 on a 6-sided die
- ► The **probability** that an event occurs is the likeliness that it will happen, a value between 0 and 1

- An event is an outcome of a random process that may or may not happen
  - Notation: A. B. etc.
  - Example: Rolling a 6 on a 6-sided die
- ► The **probability** that an event occurs is the likeliness that it will happen, a value between 0 and 1
  - Notation: p(A), p(B), etc.

- An event is an outcome of a random process that may or may not happen
  - Notation: A. B. etc.
  - Example: Rolling a 6 on a 6-sided die
- ► The **probability** that an event occurs is the likeliness that it will happen, a value between 0 and 1
  - Notation: p(A), p(B), etc.
  - Other Notation: Pr[We roll a 6]

- An event is an outcome of a random process that may or may not happen
  - Notation: A. B. etc.
  - Example: Rolling a 6 on a 6-sided die
- ► The **probability** that an event occurs is the likeliness that it will happen, a value between 0 and 1
  - Notation: p(A), p(B), etc.
  - ▶ Other Notation: Pr[We roll a 6]
  - Example: The probability of rolling a 6 on a 6-sided die is  $\frac{1}{6}$

► A random variable is a variable that can take different values each with some probability

- ► A **random variable** is a variable that can take different values each with some probability
  - ▶ Notation: *X*, *Y*, etc.

A random variable is a variable that can take different values each with some probability

Notation: 
$$X$$
,  $Y$ , etc.

Example:  $X = \begin{cases} \text{Heads} & 0.5 \\ \text{Tails} & 0.5 \end{cases}$ 

A random variable is a variable that can take different values each with some probability

Notation: X, Y, etc.

Example:  $X = \begin{cases} \text{Heads} & 0.5 \\ \text{Tails} & 0.5 \end{cases}$ 

ightharpoonup Example:  $Y \sim \mathcal{N}(0,1)$ 

▶ A random variable is a variable that can take different values each with some probability

▶ Notation: *X*, *Y*, etc.

Example:  $X = \begin{cases} \text{Heads} & 0.5 \\ \text{Tails} & 0.5 \end{cases}$ 

**Example:**  $Y \sim \mathcal{N}(0,1)$ 

▶ If X is the set of all possible outcomes for random variable X, then

$$\sum_{x \in \mathcal{X}} p(x) = 1$$

▶ Joint Probability: The probability that two events occur together

- Joint Probability: The probability that two events occur together
  - ▶ Notation: p(A, B) or  $p(A \cap B)$

- Joint Probability: The probability that two events occur together
  - Notation: p(A, B) or  $p(A \cap B)$
  - Example: Probability that we flip two heads on separate coins

- Joint Probability: The probability that two events occur together
  - ▶ Notation: p(A, B) or  $p(A \cap B)$
  - Example: Probability that we flip two heads on separate coins
- Union of Events: The probability that at least one of two (or more) events occur

- Joint Probability: The probability that two events occur together
  - ▶ Notation: p(A, B) or  $p(A \cap B)$
  - Example: Probability that we flip two heads on separate coins
- Union of Events: The probability that at least one of two (or more) events occur
  - ▶ Notation:  $p(A \cup B)$

- Joint Probability: The probability that two events occur together
  - ▶ Notation: p(A, B) or  $p(A \cap B)$
  - Example: Probability that we flip two heads on separate coins
- Union of Events: The probability that at least one of two (or more) events occur
  - ▶ Notation:  $p(A \cup B)$
  - Example: Probability that we roll 1 or 2

► Two events are **disjoint** if they can never happen at the same time

- ► Two events are **disjoint** if they can never happen at the same time
  - Example: "Coin lands on heads" vs "coin lands on tails"

- ► Two events are **disjoint** if they can never happen at the same time
  - Example: "Coin lands on heads" vs "coin lands on tails"
  - Example: "We roll a 1" vs "We roll a 6"

- Two events are disjoint if they can never happen at the same time
  - Example: "Coin lands on heads" vs "coin lands on tails"
  - Example: "We roll a 1" vs "We roll a 6"
- ► Two events are **independent** if one happening does not affect the probability that the other happens

- Two events are disjoint if they can never happen at the same time
  - Example: "Coin lands on heads" vs "coin lands on tails"
  - Example: "We roll a 1" vs "We roll a 6"
- ► Two events are **independent** if one happening does not affect the probability that the other happens
  - Example: "Coin 1 lands on heads" vs "coin 2 lands on heads"

- Two events are disjoint if they can never happen at the same time
  - Example: "Coin lands on heads" vs "coin lands on tails"
  - Example: "We roll a 1" vs "We roll a 6"
- ► Two events are **independent** if one happening does not affect the probability that the other happens
  - Example: "Coin 1 lands on heads" vs "coin 2 lands on heads"
  - Notation:  $A \perp B$  vs  $A \not\perp B$

▶ We may want to express the probability of one event (A) happening given that another event (B) also happens

- We may want to express the probability of one event (A) happening given that another event (B) also happens
  - Notation:  $p(A \mid B)$

- ► We may want to express the probability of one event (A) happening **given** that another event (B) also happens
  - Notation:  $p(A \mid B)$
  - Example: Probability we roll a 1 given we rolled less than or equal to 3

- ▶ We may want to express the probability of one event (A) happening given that another event (B) also happens
  - Notation:  $p(A \mid B)$
  - Example: Probability we roll a 1 given we rolled less than or equal to 3
- Bayes rule:

$$p(A \mid B) = \frac{p(A, B)}{p(B)}$$
$$= \frac{p(B \mid A)p(A)}{p(B)}$$

For independent events A and B,

$$p(A,B)=p(A)p(B)$$

► For independent events *A* and *B*,

$$p(A, B) = p(A)p(B)$$

► For disjoint events *A* and *B*,

$$p(A \cup B) = p(A) + p(B)$$

► For independent events A and B,

$$p(A, B) = p(A)p(B)$$

For disjoint events A and B,

$$p(A \cup B) = p(A) + p(B)$$

► Total Probability: We can partition the probability space and add up joint probabilities

For independent events A and B,

$$p(A, B) = p(A)p(B)$$

For disjoint events A and B,

$$p(A \cup B) = p(A) + p(B)$$

- Total Probability: We can partition the probability space and add up joint probabilities
  - Let B, C be a partition of the probability space  $\mathcal{X}$
  - Consider the event A
  - ► Then

$$p(A) = p(A, B) + p(A, C)$$
  
=  $p(A | B)p(B) + p(A | C)p(C)$ 

### Table of Contents

Probability Review

References

### References

 $https://www.cs.ubc.ca/\ schmidtm/Courses/Notes/probability.pdf$