

Macroeconomic Forecasting (MFx)

IMFx



This training material is the property of the International Monetary Fund (IMF) and is intended for use in IMF Institute for Capacity Development (ICD) courses. Any reuse requires the permission of the ICD.

EViews® is a trademark of IHS Global Inc.

Structural Vector Autoregressive Models (SVAR)

Part 1



Session 1. Introduction to VARs



Introduction

Del Negro and Schorfheide (2011):

*“At first glance, VARs appear to be straightforward multivariate generalizations of univariate autoregressive models. At second sight, they turn out to be **one of the key empirical tools** in modern macroeconomics.”*

What are VARs ?

- multivariate *linear* time-series models
- endogenous variable in the system are functions ***of lagged values of all endogenous*** variables
- simple and flexible alternative to the traditional multiple-equations models

Historical Overview: Sims' Critique




- In the 1980s criticized the large-scale macro-econometric models of the time
- Proposed VARs as an alternative that allowed one to model macroeconomic data informatively



What Are VARs Used For?

- **Forecasting**
 - *Reduced-Form VARs*
- **Structural Analysis**
 - *Structural VARs*

Unit Plan/Roadmap



Source:Lutkepohl (2007)

Session 2. Estimation of VARs



Introduction to VARs

- Let \mathbf{y}_t be a vector with the value of n variables at time t :

$$\mathbf{y}_t = [y_{1,t} y_{2,t} \dots y_{n,t}]'$$

- A ***p-order vector autoregressive process*** generalizes a one-variable AR(p) process to n variables:

$$\mathbf{y}_t = \mathbf{G}_0 + \mathbf{G}_1 \mathbf{y}_{t-1} + \mathbf{G}_2 \mathbf{y}_{t-2} + \dots + \mathbf{G}_p \mathbf{y}_{t-p} + \mathbf{e}_t$$

Reduced form VAR

$\mathbf{G}_0 = (n \times 1)$ vector of constants

$\mathbf{G}_j = (n \times n)$ matrix of coefficients

$\mathbf{e}_t = (n \times 1)$ vector of white noise innovations

$$E[\mathbf{e}_t] = 0$$

$$E[\mathbf{e}_t \mathbf{e}_\tau'] = \begin{cases} \Omega, & \text{if } t = \tau \\ 0 & \text{otherwise} \end{cases}$$

NOT diagonal

Example: A VAR(1) in 2 Variables

$$y_{1,t} = g_{11}y_{1,t-1} + g_{12}y_{2,t-1} + e_{1,t}$$

$$y_{2,t} = g_{21}y_{1,t-1} + g_{22}y_{2,t-1} + e_{2,t}$$

In matrix notation:

$$\mathbf{y}_t = \mathbf{G}_1 \mathbf{y}_{t-1} + \mathbf{e}_t$$

where

$$\mathbf{y}_t = \begin{pmatrix} y_{1,t} \\ y_{2,t} \end{pmatrix}, \text{ for example: } \mathbf{y}_t = \begin{bmatrix} \pi_t \\ gdp_t \end{bmatrix}$$

$$\mathbf{G}_1 = \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}, \mathbf{e}_t = \begin{pmatrix} e_{1,t} \\ e_{2,t} \end{pmatrix}$$

Assumptions about the error terms:

$$E[\mathbf{e}_t \mathbf{e}_t'] = \begin{pmatrix} \sigma_{e_1}^2 & \sigma_{e_1 e_2} \\ \sigma_{e_1 e_2} & \sigma_{e_2}^2 \end{pmatrix} = \Omega$$

$$E[\mathbf{e}_t \mathbf{e}_\tau'] = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \text{ for } t \neq \tau$$

Estimation: by OLS

- Performed with OLS applied equation by equation
- Estimates are:
 - consistent
 - efficient
 - equivalent to GLS

General Specification Choices

- *Selection of variables* to be included: in accordance with economic theory, empirical evidence and/or experience
- *Exogenous variables* can be included: constant, time trends, other additional explanators
- Non-stationary level data is often *transformed* (log levels, log differences, growth rates, etc.)
- The model should be *parsimonious*

Session 4. Stationary VARs



Stationarity of a VAR: Definition

A p -th order VAR is said to be *covariance-stationary* if:

1. The expected value of \mathbf{y}_t does not depend on time

$$E[\mathbf{y}_t] = E[\mathbf{y}_{t+j}] = \boldsymbol{\mu} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \dots \\ \mu_n \end{bmatrix}$$



Finite and
time-invariant
first and
second order
moments

2. The covariance matrix of \mathbf{y}_t and \mathbf{y}_{t+j} depends on the time lapsed j and not on the reference period t

$$E[(\mathbf{y}_t - \boldsymbol{\mu})(\mathbf{y}_{t+j} - \boldsymbol{\mu})'] = E[(\mathbf{y}_s - \boldsymbol{\mu})(\mathbf{y}_{s+j} - \boldsymbol{\mu})'] = \boldsymbol{\Gamma}_j$$


Conditions for Stationarity

- The conditions for a VAR to be stationary are similar to the conditions for a univariate AR process to be stationary:

$$\mathbf{y}_t = \mathbf{G}_0 + \mathbf{G}_1 \mathbf{y}_{t-1} + \mathbf{G}_2 \mathbf{y}_{t-2} + \dots + \mathbf{G}_p \mathbf{y}_{t-p} + \mathbf{e}_t$$
$$(\mathbf{I}_n - \mathbf{G}_1 \mathbf{L} - \mathbf{G}_2 \mathbf{L}^2 - \dots - \mathbf{G}_p \mathbf{L}^p) \mathbf{y}_t = \mathbf{G}_0 + \mathbf{e}_t$$

Lag operator

$$\mathbf{G}(\mathbf{L}) \mathbf{y}_t = \mathbf{G}_0 + \mathbf{e}_t$$

Lag polynomial

- For \mathbf{y}_t to be stationary, the matrix polynomial in the lag operator $\mathbf{G}(\mathbf{L})$ must be invertible


Conditions for Stationarity

- A VAR(p) process is stationary (thus invertible) if all the np roots of the *characteristic polynomial* are (in modulus) outside the unit imaginary circle


$$\det(\mathbf{I}_n - \mathbf{G}_1\mathbf{L} - \mathbf{G}_2\mathbf{L}^2 - \dots - \mathbf{G}_p\mathbf{L}^p) = 0$$

- EViews calculates the *inverse roots* of the characteristic AR polynomial, which should then lie *within* the unit imaginary circle

Inverse Roots of AR Characteristic Polynomial



Vector Moving Average Representation of a VAR

- If a VAR is stationary, the \mathbf{y}_t vector can be expressed as a sum of all of the past white noise shocks \mathbf{e}_t (**VMA(∞) representation**)

$$\mathbf{y}_t = \boldsymbol{\mu} + \mathbf{G}(\mathbf{L})^{-1} \mathbf{e}_t, \text{ where } \boldsymbol{\mu} = \mathbf{G}(\mathbf{L})^{-1} \mathbf{G}_0$$

$$\mathbf{y}_t = \boldsymbol{\mu} + (\mathbf{I}_n + \boldsymbol{\Psi}_1 \mathbf{L} + \boldsymbol{\Psi}_2 \mathbf{L}^2 + \dots) \mathbf{e}_t$$

$$\mathbf{y}_t = \boldsymbol{\mu} + \mathbf{e}_t + \boldsymbol{\Psi}_1 \mathbf{e}_{t-1} + \boldsymbol{\Psi}_2 \mathbf{e}_{t-2} + \dots$$

$$\mathbf{y}_t = \boldsymbol{\mu} + \sum_{i=0}^{\infty} \boldsymbol{\Psi}_i \mathbf{e}_{t-i}$$

*Wold
theorem*

- where $\boldsymbol{\Psi}_i$ is a $(n \times n)$ matrix of coefficients, and $\boldsymbol{\Psi}_0$ is the identity matrix.
- From the VMA(∞) representation it is possible to obtain impulse response functions

Session 5. Lag Specification Criteria



Lags Needed for the VAR

What number is most appropriate? ...


- If p is extremely short, the model may be poorly specified
- If p is extremely long, too many degrees of freedom will be lost
- The number of lags should be sufficient for the residuals from the estimation to constitute individual white noises

The Curse of Dimensionality

- VARs are very densely parametrized
 - In a VAR(p) we have p matrices of dimension $n \times n$: G_1, \dots, G_p
 - Assume G_0 is an intercept vector (dimension: $n \times 1$)
 - The number of *total* coefficients/parameters to be estimated is:

$$n + n \times n \times p = n(1 + nxp)$$

Overfitting versus Omitted Variable Bias

- *Over-fitting* problem
 - *Omitted variable bias*
- 
- poor-quality estimates and bad forecasts
- Possible solutions:
 - Core VAR plus rotating variables
 - Bayesian analysis

Lag Length Criteria

- As for univariate models, one can use multidimensional versions of the:
 - AIC: Akaike information criterion
 - SC: Schwarz information criterion
 - HQ: Hanna-Quinn information criterion
- *Information-based criteria*: trade-off between parsimony and reduction in sum of squares

Lag Specification: Practitioner's Advice

Rules of Thumb:

- $p = 4$ when working with quarterly data
- $p = 12$ with monthly data
- The effective constraint is $np < T/3$

Example:

$$T = 100$$

$$p = 4$$

$$n \leq 7$$

Session 8. Forecasting using VARs



Forecasting Using the Estimated VAR

- Let \mathbf{Y}_{t-1} be a matrix containing all information available up to time t (before realizations of \mathbf{e}_t are known):

- Then:
$$\mathbf{Y}_{t-1} = (\mathbf{y}_{t-1}, \mathbf{y}_{t-2}, \dots, \mathbf{y}_{t-T})$$

$$E[\mathbf{y}_t | \mathbf{Y}_{t-1}] = \mathbf{G}_0 + \mathbf{G}_1 \mathbf{y}_{t-1} + \mathbf{G}_2 \mathbf{y}_{t-2} + \dots + \mathbf{G}_p \mathbf{y}_{t-p}$$

Forecasting Using the Estimated VAR

- The forecast error can be decomposed into the sum of \mathbf{e}_t , the unexpected innovation of \mathbf{y}_t , and the coefficient estimation error:

$$\mathbf{y}_t - E[\mathbf{y}_t \mid \mathbf{Y}_{t-1}] = \mathbf{e}_t + \boldsymbol{\nu}(\mathbf{Y}_{t-1})$$

- If the estimator for the coefficients is consistent and estimates are based on many data observations, the coefficient estimation error tends to be small, and:

$$\mathbf{y}_t - E[\mathbf{y}_t \mid \mathbf{Y}_{t-1}] \cong \mathbf{e}_t$$

Iterated Forecasts

- Iterating one period forward:

$$E[\mathbf{y}_{t+1} \mid \mathbf{Y}_{t-1}] = \mathbf{G}_0 + \mathbf{G}_1 E[\mathbf{y}_t \mid \mathbf{Y}_{t-1}] + \mathbf{G}_2 \mathbf{y}_{t-1} + \dots + \mathbf{G}_p \mathbf{y}_{t-p+1}$$

- Iterating j periods forward:

$$E[\mathbf{y}_{t+j} \mid \mathbf{Y}_{t-1}] = \mathbf{G}_0 + \mathbf{G}_1 E[\mathbf{y}_{t+j-1} \mid \mathbf{Y}_{t-1}] + \mathbf{G}_2 E[\mathbf{y}_{t+j-2} \mid \mathbf{Y}_{t-1}] + \dots + \mathbf{G}_p \mathbf{y}_{t-p+j}$$