

1. A blood pressure of systolic 160-179mmHg and diastolic 100-109mmHg can be classified as:

- a. Stage two hypertension
- b. Stage one hypertension
- c. Optimal hypertension
- d. Stage three hypertension

****Answer: b. Stage one hypertension****

2. The following may be possible cause of hypertension except:

- a. genetic variation in kidney sodium handling
- b. increased renal absorption of sodium chloride
- c. dysfunction of the autonomic nervous system
- d. decrease activity of the renin-angiotensin-aldosterone system

****Answer: d. decrease activity of the renin-angiotensin-aldosterone system****

3. A type of pericarditis characterized by fibrinous or effusive exudates:

- a. Chronic pericarditis
- b. Post cardiac pericarditis
- c. Acute pericarditis
- d. Autoimmune pericarditis

****Answer: c. Acute pericarditis****

4. Jugular vein distention indicates failure of:

- a. Right-sided heart failure
- b. Left-sided heart failure

****Answer: a. Right-sided heart failure****

5. The largest heart chamber with the thickest myocardium is:

- a. Right Ventricle
- b. Left atrium
- c. Left Ventricle
- d. Right atrium

****Answer: c. Left Ventricle****

6. Crackles are a sign of:

- a. Left-sided heart failure
- b. Right-sided heart failure

****Answer: a. Left-sided heart failure****

7. An early sign of anaemia is:

- a. Palpitation
- b. Weight loss
- c. Pallor
- d. Glossitis

****Answer: c. Pallor****

8. A client with sodium level 160 mEq/L should avoid:

- a. Cauliflower
- b. Processed oat cereals
- c. Blended banana
- d. Peanut

****Answer: b. Processed oat cereals****

9. Hepatomegaly indicates:

- a. Left-sided heart failure
- b. Right-sided heart failure

****Answer: b. Right-sided heart failure****

10. Focus of care for hypertension is:

- a. Ineffective airway clearance
- b. Activity intolerance
- c. Deficient knowledge
- d. Impaired physical mobility

****Answer: c. Deficient knowledge****

11. Splenomegaly indicates:

- a. Left-sided heart failure
- b. Right-sided heart failure

****Answer: b. Right-sided heart failure****

12. A normal breathing pattern is:

- a. Ventilation
- b. Bradypnea
- c. Tachypnea
- d. Eupnea

****Answer: d. Eupnea****

13. High-normal systolic blood pressure range:

- a. 120–128
- b. 130–139
- c. 140–170
- d. 130–150

****Answer: b. 130–139 mmHg****

14. Chest pain caused by cardiac ischemia is:

- a. Atheroma
- b. Ischaemia
- c. Atherosclerosis
- d. Angina pectoris

****Answer: d. Angina pectoris****

15. Painless edema of the legs that disappears at night indicates:

- a. Pulmonary oedema
- b. Lung disease
- c. MI
- d. Right ventricular heart failure

****Answer: d. Right ventricular heart failure****

16. Cardiac output formula:

- a. $PR \times SV$
- b. $BP \div PR$
- c. $HR \times SV$
- d. $HR + SV$

****Answer: c. Product of heart rate and stroke volume****

17. Severe distress with frothy sputum indicates:

- a. Pneumonia
- b. Pulmonary oedema
- c. Pneumothorax
- d. Cardiac tamponade

****Answer: b. Pulmonary oedema****

18. Drug that increases thrombosis risk:

- a. Sulphonamides
- b. Antacids
- c. Iron supplements
- d. Oral contraceptives

****Answer: d. Oral contraceptives****

19. NOT a common pulmonary embolism feature:

- a. Chest pain
- b. Tachypnea
- c. Bradypnea
- d. Dyspnea

****Answer: c. Bradypnea****

20. Kidneys release this when BP drops:

- a. Erythropoietin
- b. Aldosterone
- c. Renin
- d. Epinephrine

****Answer: c. Renin****

21. Most common atrial septal defect:

- a. Ostium Secundum
- b. Sinus Venosa
- c. Ostium Primum
- d. Sinus Venosus

****Answer: a. Ostium Secundum defect****

22. A modifiable CAD risk factor:

- a. Race
- b. Hypertension
- c. Age
- d. Gender

****Answer: b. Hypertension****

23. Dry cough indicates:

- a. Left-sided heart failure
- b. Right-sided heart failure

****Answer: a. Left-sided heart failure****

24. Type A personalities are prone to:

- a. Varicose veins
- b. Atherosclerosis
- c. Heart attack
- d. Aneurysm

****Answer: c. Heart attack****

25. Most common pericarditis symptom:

- a. Calf pain
- b. Claudication
- c. Chest pain
- d. Dyspnea

****Answer: c. Chest pain****

26. Hormones controlling fluid balance:

- a. Aldosterone & ADH
- b. Parathormone & Calcitonin
- c. Calcitonin & ADH
- d. ACE & ADH

****Answer: a. Aldosterone and ADH****

27. Daily assessment for heart failure severity:

- a. Abdominal girth
- b. Calorie count
- c. Weight
- d. Appetite

****Answer: c. Weight****

28. Best explanation of heart failure:

- a. Heart stops briefly
- b. Heart is not an efficient pump
- c. Heart muscle is dead
- d. Heart pumps too much blood

****Answer: b. Heart is not an efficient pump****

29. Dyspnoea indicates:

- a. Left-sided heart failure
- b. Right-sided heart failure

****Answer: a. Left-sided heart failure****

30. Risk for fluid volume deficit:

- a. Frequent wound irrigation
- b. CHF
- c. Colostomy
- d. Increased kidney function

****Answer: a. Frequent wound irrigation****

31. Structure separating left and right heart:

- a. Chamber
- b. Septum
- c. Valve
- d. Pericardium

****Answer: b. Septum****

32. Myocardial muscle fiber pattern:

- a. Circular
- b. Figure of eight
- c. Longitudinal
- d. Lateral

****Answer: b. Figure of eight pattern****

33. Hypertension lifestyle modification:

- a. Low-tar cigarettes
- b. Regular aerobic exercise
- c. Multivitamin
- d. 3 alcoholic drinks/day

****Answer: b. Regular aerobic exercise****

34. Blood returns to heart via:

- a. Pulmonary vein
 - b. RCA
 - c. Aorta
 - d. Vena cavae
- **Answer: d. Vena cavae****

35. True statements about VSD:

- a. I only
 - b. I, II, III
 - c. II and III
 - d. I and II
- **Answer: c. II and III****

36. Test with tourniquet and walking:

- a. Angiography
 - b. Trendelenburg test
 - c. Perthes test
 - d. Phlebography
- **Answer: c. Perthes test****

37. Peripheral oedema indicates:

- a. Right-sided HF
 - b. Left-sided HF
- **Answer: a. Right-sided HF****

38. Type A personality risk:

- a. Aneurysm
 - b. Atherosclerosis
 - c. Heart attack
 - d. Varicose veins
- **Answer: c. Heart attack****

39. NOT part of Dressler's syndrome:

- a. Fever
 - b. Leukocytosis
 - c. Endocarditis
 - d. Chest pain
- **Answer: c. Endocarditis****

40. Best advice after 12 months lifestyle modification (BP 158/98):

- a. Medication required
 - b. Stop drugs
 - c. More vigorous lifestyle changes
 - d. Monitor every 3 months
- **Answer: a. Medication required****