

experiencing which complication of a blood transfusion?

A. Septicemia ✓

B. Hyperkalemia

C. Circulatory overload

D. Delayed transfusion reaction

43. What is the primary goal of a nurse when engaging the family in care discussions for a patient with a severe injury in the Emergency Department?

A. Ensuring the family is informed about hospital policies

B. Encouraging the family to participate in the patient's care decisions

C. Providing emotional support to the family during the crisis ✓

D. Clarifying the roles of different healthcare providers

44. When evaluating a patient in the Emergency Department who exhibits unexplained injuries and is accompanied by someone who is excessively protective and controlling, what should the nurse prioritize for further assessment?

A. Substance abuse

B. Domestic violence ✓

C. Mental health crisis

D. Potential human trafficking

45. The nurse assesses a patient with ischemia and chest pain. The patient should be assigned what Emergency Severity Index (ESI) rating?

A. ESI-1

B. ESI-2 ✓

C. ESI-3

D. ESI-4

46. Which psychological factor often inhibits victims of sexual violence from disclosing their experiences during nursing assessments in emergency care settings?

A. Fear of retaliation from the perpetrator ✓

B. Belief that the incident was their fault

C. Concern about the emotional impact on family members

D. Fear of being stigmatized

47. A nurse is about to discharge a patient who has been treated for a fractured wrist. During the discharge conversation, the patient expresses concerns about increased swelling and discomfort in the wrist that was not noted before. What is the most appropriate response by the nurse?

A. The nurse should refer the patient to a specialist immediately.

B. The nurse should conduct a focused assessment of the wrist ✓

C. The nurse should reassure the patient that swelling is normal.

D. The nurse should document the patient's concerns and proceed with discharge.

48. Which assessment tool is specifically designed to prioritize patient care in emergency situations based on the severity of their condition?

A. Emergency Severity Index (ESI) ✓

B. ABCDEF

C. SALT

D. Nursing Assessment Tool

49. What assessment tool is designed to categorize patients based on the severity of their condition in the Emergency Department?

A. Triage Assessment Tool

B. Emergency Severity Index (ESI) ✓

C. Patient Acuity Scale

D. Critical Care Assessment Matrix

50. A triage system used for determining patient treatment priority during mass casualty incidents: Sort, Assess, Lifesaving interventions, Treatment/Transport. What are the sources of assessment data?

A. client, family/significant other, health care team

B. client, family/significant other, health care team, medical records

C. client, family/significant other, health care team, medical records, literature

D. client, family/significant other, health care team, medical records, literature, nurse's experience ✓

51. In which healthcare setting is the ABCDEF assessment tool primarily utilized to enhance patient outcomes?

A. Emergency Department ✓

- B. Critical Care Units
- C. Outpatient Clinics
- D. Long-term Care Facilities

52. Which intervention does the nurse implement for the active external rewarming of a patient experiencing hypothermia?

- A. Administer humidified oxygen.
- B. Remove the patient's wet clothing.
- C. Immerse the patient in warm water. ✓✓**
- D. Place the patient under radiant lighting.

53. Which effect does the nurse anticipate in a patient who experiences respiratory impairment after a submersion injury in fresh water?

- A. Extreme thirst
- B. Pulmonary edema ✓✓**
- C. Muscle contractions
- D. Altered mental status

54. A patient who attempted suicide is brought into the emergency department by a parent. The parent states, "My child consumed liquid bleach." The nurse expects which clinical manifestation?

- A. Jaundice
- B. Almond odor to breath
- C. Respiratory depression
- D. Irritation of lips, mouth, and eyes ✓✓**

55. The nurse understands that which fluid and electrolyte change occurs during the early emergent phase of burn injury?

- A. Increased excretion of urine
- B. Increased colloidal osmotic pressure

C. Return of interstitial fluid to the vascular space

D. Movement of sodium and water into interstitial spaces ✓

56. The nurse assesses a patient who is rated as an ESI-1 on the Emergency Severity Index (ESI). The nurse identifies that which procedure is preferred for securing an unprotected airway in the emergency department (ED)?

A. Rapid-sequence intubation ✓

B. Jaw-thrust maneuver

C. Tracheotomy

D. A rigid cervical collar (C collar)

57. Which type of dysrhythmia is treated with a pacemaker?

A. Atrial fibrillation

B. Sinus tachycardia

C. Ventricular fibrillation

D. Third-degree atrioventricular (AV) block ✓

58. Which action by the emergency department disaster team will be most useful in preparing hospital staff to provide care in a disaster?

A. Provide a mass casualty community drill. ✓

B. Show a video of a disaster relief group working.

C. Have a police officer talk with the staff about disasters.

D. Ask the staff to write a scenario to turn in to the manager.

59. Which intervention would the nurse expect to incorporate into the nutritional plan of care of a burn patient who has been intubated?

A. Start total parenteral nutrition.

B. Provide enteral tube feeding, starting at 20 mL/hour. ✓

C. Provide bolus enteral tube feedings four times a day.

D. Feed at least 1500 calories/day in small, frequent meals.

60. A burn patient newly arrived at the emergency department is confused, has difficulty breathing, and has nasal burns. Which condition would the nurse suspect?

- A. Pulmonary edema
- B. Lower airway injury ✓**
- C. Upper airway injury
- D. Carbon monoxide poisoning

61. On a hot summer day, a patient presents to the emergency department reporting extreme thirst and vomiting. The patient's assessment findings include profuse diaphoresis, BP of 90/50 mm Hg, pulse 98 beats/minute, and a temperature of 101° F (38.3° C). The nurse performs which action?

- A. Administers salt tablets
- B. Initiates oral fluid and electrolyte replacement
- C. Starts an IV fluid bolus of lactated Ringer's
- D. Places the patient in a cool area and removes restrictive clothing ✓**

62. The nurse observes a patient that seeks treatment for multiple bee stings. The nurse monitors the patient for which indication of a severe reaction?

- A. Fatigue
- B. Stiff neck
- C. Headache ✓**
- D. Bullseye rash

63. The nurse is conducting a primary survey in an emergency department. Which is the purpose of the survey?

- A. To assess whether the patient has any threat to life
- B. To determine the priority for treatment for patients who are in the emergency department
- C. To evaluate whether the resources in the emergency department are adequate to treat the patient
- D. To evaluate the status of airway, breathing, circulation, disability, and exposure and environmental control ✓**

64. A patient is admitted to the emergency department with a traumatic head injury. Which action does the nurse take during the primary survey?

A. Uses the Glasgow Coma Scale ✓✓

B. Removes the patient's clothing

C. Obtains a portable chest x-ray exam

D. Monitors the electrocardiogram (ECG) for heart rate and rhythm

65. Nurse Edna is assigned to telephone triage. A client called who was stung by a honeybee and is asking for help. The client reports pain and localized swelling but has no respiratory distress or other symptoms of anaphylactic shock. What is the appropriate **initial** action that the nurse should direct the client to perform?

a) Applying a cold compress to the area

b) Taking an oral antihistamine

c) Removing the stinger by scraping it ✓✓

d) Calling the doctor to address it

66. Nurse Anna is an experienced travel nurse who was recently employed and is assigned to the emergency unit. In her first week of the job, which of the following area is the **most** appropriate assignment for her?

a) Triage

b) Ambulatory section ✓✓

c) Trauma team

d) Psychiatric care

67. A client arrives at the emergency department who suffered multiple injuries from a head-on car collision. Which of the following assessments should take the **highest** priority to take?

a) Unequal pupils

b) Irregular pulse

c) Bruise in the flank

d) A deviated trachea ✓

68. An anxious female client complains of chest tightness, tingling sensations, and palpitations. Deep, rapid breathing, and carpal spasms are noted. Which of the following priority action

should the nurse do **first**?

a) Provide oxygen therapy

b) Notify the physician immediately

c) Administer anxiolytic medication as ordered

d) Have the client breathe into a brown paper bag ✓

69. Michael works as a triage nurse, and four clients arrive at the emergency department at the

same time. List the order in which he will assess these clients from first to last.

I. An ambulatory dazed 25-year-old male with a bandaged head wound

II. An irritable infant with a fever, petechiae, and nuchal rigidity

III. A 35-year-old jogger with a twisted ankle, having a pedal pulse, and no deformity

IV. A 50-year-old female with moderate abdominal pain and occasional vomiting

a) I, II, III, IV

b) II, I, IV, III ✓

c) IV, III, I, II

d) III, II, I, IV

70. In conducting a primary survey on a trauma patient, which of the following is considered one of the **priority** elements of the primary survey?

a) Brief neurologic assessment ✓

b) Client's allergy history

c) Initiation of pulse oximetry

d) Complete set of vital signs

71. A 65-year-old patient arrived at the triage area with complaints of diaphoresis, dizziness, and left-sided chest pain. This patient should be prioritized into which category?

a) Non-urgent

b) Urgent

c) Emergent ✓

d) High urgent

72. You respond to a call for help from the ED waiting room. There is an elderly patient lying on the floor. List the order for the actions that you must perform.

I. Perform the chin lift or jaw thrust maneuver

II. Initiate cardiopulmonary resuscitation (CPR)

III. Establish unresponsiveness

IV. Get the emergency cart and activate the code team

a) III, IV, I, II ✓

b) I, IV, III, I

c) II, III, I, IV

d) IV, II, I, III

73. A 15-year-old male client arrives at the emergency department. He is conscious, coherent, and ambulatory, but his shirt and pants are covered with blood. He and his hysterical friends are yelling and trying to explain that they were goofing around and he got poked in the abdomen with a stick. Which of the following comments should be given **first** consideration?

a) "The stick was really dirty and covered with mud."

b) "He pulled the stick out, just now, because it was hurting him." ✓

c) "He's a diabetic, so he needs attention right away."

d) "There was a lot of blood and we used three bandages."

74. A client arrived at the emergency department after suffering multiple physical injuries including a fractured pelvis from a vehicular accident. Upon assessment, the client is incoherent, pale, and diaphoretic. With vital signs as follows: temperature of 97°F (36.11° C), blood pressure of 60/40 mm Hg, heart rate of 143 beats/minute, and a respiratory rate of 30 breaths/minute. The client is mostly suffering from which of the following shock?

- a) Distributive
- b) Obstructive
- c) Cardiogenic
- d) Hypovolemic ✓

75. A 15-year-old male client was sent to the emergency unit following a small laceration on the forehead. The client says that he can't move his legs. Upon assessment, respiratory rate of 20, strong pulses, and capillary refill time of less than 2 seconds. Which triage category would this client be assigned to?

- a) Red
- b) Black
- c) Yellow ✓
- d) Green

76. When attending to a client with a head and neck trauma following a vehicular accident, the nurse's **initial** action is to?

- a) Provide oxygen therapy
- b) Initiate intravenous access
- c) Immobilize the cervical area ✓
- d) Do oral and nasal suctioning

77. Which of the following actions is the **priority** of care for a client exhibiting signs and symptoms of coronary artery disease?

- a) Decrease anxiety level of patient

b) Enhance myocardial oxygenation. ✓

c) Administer sublingual nitroglycerin.

d) Educate the client about his symptoms.

78. A patient with status asthmaticus is receiving oxygen via a non-rebreather mask. The patient is diaphoretic, using accessory muscles to breathe, and has a respiratory rate of 32 breaths/min. Arterial blood gas results show: pH 7.28, PaCO₂ 50 mmHg, PaO₂ 62 mmHg, HCO₃ 24 mEq/L. What does this indicate about the patient's condition?

a) Adequate oxygenation with respiratory alkalosis

b) Need to switch to a Venturi mask for controlled oxygen delivery

c) Hyperventilation requiring sedation

d) Impending respiratory failure requiring intubation ✓

79. patient is receiving oxygen via a non-rebreather mask at 15 L/min. The nurse observes that

the reservoir bag is completely deflating during inspiration. What should the nurse do FIRST?

a) Increase the oxygen flow rate ✓

b) Switch to a Venturi mask

c) Assess the patient for respiratory distress

d) Decrease the oxygen flow rate

80. A patient presents to the emergency department with acute respiratory distress. His oxygen saturation is 86% on room air. Which oxygen delivery device would be MOST appropriate

to deliver a high concentration of oxygen?

a) Simple face mask

b) Nasal cannula

c) Non-rebreather mask ✓