

d) Venturi mask

81. A nurse is preparing to administer oxygen via a nasal cannula to a patient with mild hypoxemia. What is the maximum flow rate typically recommended for nasal cannula oxygen delivery?

a) 2 L/min

b) 4L/min

c) 6 L/min ✓

d) 10 L/min

82. A nurse is administering oxygen via a simple face mask at 8 L/min. The patient complains of feeling claustrophobic and repeatedly removes the mask. Which alternative oxygen delivery method would be MOST appropriate?

a) Nasal cannula at 4 L/min ✓

b) Non-rebreather mask at 12 L/min

c) Face tent at 8 L/min

d) Venturi mask at 40%

83. A 68-year-old patient with COPD exacerbation requires oxygen therapy at 3 L/min via nasal

cannula. The patient will likely need oxygen for 3-4 days. Which type of oxygen delivery would be most appropriate?

a) Dry oxygen

b) Humidified oxygen ✓

c) Helium-oxygen mixture

d) High-flow oxygen

84. The nurse witnesses a neighbor's husband sustain a fall from the roof of his house. The nurse rushes to the victim and determines the need to open the airway. The nurse opens the airway

in this victim by using which method?

- a) Flexed position
- b) Head tilt-chin lift
- c) Jaw thrust maneuver ✓**
- d) Modified head tilt-chin lift

85. The nurse is performing rescue breathing on a 7-year-old child. The nurse delivers one breath per how many seconds to the child?

- a) 6-8 ✓**
- b) 8-10
- c) 10-12
- d) 12-14

86. The nurse is performing cardiopulmonary resuscitation (CPR) on an adult client. When performing chest compressions, the nurse should depress the sternum by how many inch(es)?

- a) 3/4 inch
- b) 1 inch
- c) 2 inches ✓**
- d) 3 inches

87. while triaging clients in a crowded emergency department, a nurse assesses a client who presents with symptoms of tuberculosis. Which action should the nurse take first?

- a) Apply oxygen via nasal cannula.
- b) Administer intravenous 0.9% saline solution.
- c) Obtain a sputum culture and sensitivity.

d) Transfer the client to a negative-pressure room ✓

88. An emergency department (ED) nurse is preparing to transfer a client to the trauma intensive

care unit. Which information should the nurse include in the nurse-to-nurse hand-off report?

a) Patient educational status

b) Diagnostic test results ✓

c) Immunizations

d) List of home medications

89. An emergency room nurse is triaging victims of a multi-casualty event. Which client should receive care first?

a) A 30-year-old distraught mother holding her crying child

b) A 65-year-old conscious male with a head laceration

c) A 26-year-old male who has pale, cool, clammy skin ✓

d) A 48-year-old with a simple fracture of the lower leg

90. An emergency room nurse assesses a client who has been raped. With which health care team member should the nurse collaborate when planning this client's care?

a) Forensic nurse examiner ✓

b) Emergency medicine physician

c) Case manager

d) Psychiatric crisis nurse

91. A patient with Parkinson's disease has a nursing diagnosis of Impaired Physical Mobility related to neuromuscular impairment. You observe an Enrolled nurse performing all of these actions. For which action must you intervene?

a) The NA assists the patient to ambulate to the bathroom and back to bed.

b) The NA reminds the patient not to look at his feet when he is walking.

c) The NA sets up the patient's tray and encourages the patient to feed himself.

d) The NA performs the patient's complete bath and oral care. ✓✓

92. A patient with a spinal cord injury at level C3-4 is being cared for in the ED. What is the priority assessment?

a) Determine the level at which the patient has intact sensation.

b) Assess the level at which the patient has retained mobility.

c) Check blood pressure and pulse for signs of spinal shock.

d) Monitor respiratory effort and oxygen saturation level. ✓✓

93. A patient is admitted with a Glasgow Coma Scale (GCS) score of 6 following a motor vehicle accident. Which nursing diagnosis takes highest priority?

a) Risk for ineffective cerebral tissue perfusion

b) Disturbed sensory perception

c) Ineffective airway clearance ✓✓

d) Risk for falls

94. A patient with a traumatic brain injury has an ICP of 26 mmHg and a mean arterial pressure (MAP) of 70 mmHg. The nurse calculates the cerebral perfusion pressure (CPP) as:

a) 44 mmHg ✓✓

b) 96 mmHg

c) 70 mmHg

d) 26 mmHg

95. A neurosurgical patient has an external ventricular drain (EVD) in place. The nurse should immediately notify the physician if:

a) The drainage is clear

b) The patient complains of a headache

c) The drainage appears cloudy or bloody ✓✓

d) The waveform on the monitor changes with patient movement

96. Following a decompressive craniectomy, which nursing intervention is most important?

a) Maintaining the patient in a flat position

b) Protecting the unprotected brain area ✓✓

c) Encouraging coughing and deep breathing

d) Administering prophylactic antibiotics routinely

97. A patient with severe traumatic brain injury is being treated with 3% hypertonic saline. The primary purpose of this treatment is to:

a) Increase serum sodium levels

b) Reduce cerebral edema ✓✓

c) Prevent seizure activity

d) Improve renal perfusion

98. A client in the emergency department presents with acute onset of dyspnea, tachycardia, and hypoxemia. The nurse suspects pulmonary embolism. Which diagnostic test should the nurse

anticipate will be ordered first?

a) Ventilation-perfusion scan

b) Pulmonary angiogram

c) Duplex ultrasound of lower extremities

d) D-dimer assay ✓✓

99. A patient is admitted with a severe head injury following a motor vehicle accident. During assessment, the nurse observes clear fluid leaking from the patient's nose. Which of the following is the priority nursing action?

a) Position the patient with the head of bed elevated at 30 degrees ✓✓

b) Test the fluid for glucose using a reagent strip

- c) Insert nasal packing to stop the drainage
- d) Administer prophylactic antibiotics as ordered

100. A nurse is assessing a patient with a head injury who has a Glasgow Coma Scale (GCS) score of 12. Which of the following statements about this patient's condition is correct?

- a) The patient has a severe head injury
- b) The patient has a moderate head injury ✓**
- c) The patient has a mild head injury
- d) The GCS cannot determine severity of head injury

101. A nurse is caring for a patient who experienced a head injury 12 hours ago. Which of the following assessment findings should the nurse report immediately to the healthcare provider?

- a) Blood pressure 152/88 mmHg
- b) Development of pupillary light reflex
- c) Confusion about the current date ✓
- d) Unilateral pupillary asymmetry ✓**

102. A nurse is monitoring a patient with increased intracranial pressure (ICP) following a traumatic brain injury. Which of the following vital sign changes would be consistent with Cushing's triad?

- a) Decreased blood pressure, increased pulse, increased respirations
- b) Increased blood pressure, decreased pulse, irregular respirations ✓**
- c) Decreased blood pressure, decreased pulse, decreased respirations
- d) Increased blood pressure, increased pulse, increased respirations

103. A patient with a head injury is exhibiting abnormal flexion (posturing). The nurse understands that this indicates:

- a) Damage to the cerebral cortex or internal capsule**

b) Damage to the midbrain or upper brainstem ✓

c) Damage to the lower brainstem or upper spinal cord

d) Damage to the cerebellum

104. A client is admitted to the ICU with massive pulmonary embolism and is experiencing hypotension. Which intervention should the nurse anticipate?

a) Administration of diuretics

b) Fluid bolus administration ✓

c) Placement of arterial line for BP monitoring

d) Immediate initiation of prone positioning

105. A client with a history of VTE is admitted to the emergency department with recurrent thrombosis despite therapeutic anticoagulation. Which condition should the nurse suspect?

a) Protein deficiency

b) Heparin-induced thrombocytopenia (HIT)

c) Antiphospholipid syndrome ✓

d) Factor V Leiden mutation

106. A nurse is preparing to initiate a heparin infusion for a client with pulmonary embolism. Which of the following should be available at the bedside before starting therapy?

a) Fresh frozen plasma

b) Protamine sulphate ✓

c) Factor VIII concentrate

d) Cryoprecipitate

107. A nurse is caring for a client in the emergency department who has been diagnosed with pulmonary embolism (PE). Which intervention should receive the highest priority?

a) Administration of ordered anticoagulant therapy

b) Initiating continuous cardiac monitoring

c) Ensuring adequate oxygenation ✓

d) Obtaining a D-dimer level

108. A nurse is preparing to administer furosemide (Lasix) 40 mg IV push to a client with acute heart failure. Which nursing intervention is most important before administering this medication?

a) Assess for allergies to sulpha drugs

b) Check the client's potassium level ✓

c) Place the client in high Fowler's position

d) Restrict fluid intake to 1,000 mL/day

109. A client presents to the emergency department with severe dyspnea, respiratory rate of 32 breaths/minute, oxygen saturation of 85% on room air, and diffuse crackles throughout all lung fields. Which immediate nursing intervention takes highest priority?

a) Administer IV furosemide as ordered

b) Place the client on continuous cardiac monitoring

c) Administer supplemental oxygen ✓

d) Obtain a 12-lead ECG

110. A nurse is caring for a client with acute heart failure who has been prescribed nitroglycerin 0.4 mg sublingual. Before administering this medication, the nurse should assess the client's:

a) Blood glucose level

b) Blood pressure ✓

c) Respiratory rate

d) Temperature

111. A nurse is implementing discharge instructions for a client with newly diagnosed heart failure. Which statement by the client indicates understanding of self-management?

- a) "I should restrict my fluids to 3,000 mL per day."
- b) "I will weigh myself weekly to monitor for fluid retention."
- c) "I should report a weight gain of 0.91-1.36 Kg in a day." ✓**
- d) "I can take over-the-counter NSAIDs for joint pain."

112. When caring for a patient with multiple trauma injuries, the nurse understands that the primary survey follows the ABCDE approach. What does the "E" represent?

- a) Examination
- b) Evaluation
- c) Exposure ✓**
- d) Emergency

113. A trauma patient is experiencing significant physiological stress. The nurse understands that the patient is most likely to develop:

- a) Reduced metabolic demands
- b) Decreased inflammatory response
- c) Stress-induced hypermetabolism ✓**
- d) Improved immune function

114. In a trauma patient experiencing shock, the earliest compensatory mechanism the nurse would observe is:

- a) Decreased respiratory rate
- b) Tachycardia ✓**
- c) Decreased urine output
- d) Reduced mental status