

- c. Metaplasia
 - d. Neoplasm
182. Cancer is not a single disease with a single cause. When it affects the nerve cells, it is called
- a. **Neuroblastoma**
 - b. Neuroma
 - c. Glioma
 - d. Glioblastoma
183. Kaposi's sarcoma is associated with:
- a. Human papilloma virus
 - b. Human T cell lymphotropic virus type 1 (HTLV-1)
 - c. **Human immunodeficiency virus**
 - d. Epstein Barr virus
184. Helicobacter pylori is believed to be associated with
- a. Prostate cancer
 - b. Lung cancer
 - c. Vaginal cancer
 - d. **Gastric cancer**
185. The following are true about HBV except
- a. HBV has a long incubation period
 - b. HBV is transmitted percutaneously or through permucosal routes
 - c. **The infection is usually transmitted from the mother to the fetus via umbilical vein**
 - d. It replicates in the liver and remains in the serum
186. Nurse Wilson must apply an elastic bandage to a client's ankle and calf. He should apply the bandage beginning at the client
- a. Knee
 - b. Angle
 - c. Lower thigh
 - d. **Foot**
187. While delivering speech on causes and effect of Down Syndrome, nurse Okpo said that down syndrome is a common
- a. X-Chromosomes disorder
 - b. Y-chromosome disorder
 - c. XO disorder of females
 - d. **Autosomal disorder**
188. A mother has recently been informed that her child has Down's syndrome. Which of the following characteristics is not associated with Down's syndrome?

- a. Simian crease
 - b. Brachycephaly
 - c. Oily skin**
 - d. Hypotonicity
189. Bariatric is a type of surgery performed for correction of
- a. Morbid obesity**
 - b. Diverticulum
 - c. Uterine prolapse
 - d. Wear and tear at the joints
190. Pulse pressure widens in normal individuals as they age because of which of the following factors?
- a. Stroke volume increases
 - b. Venous return increase
 - c. Heart rate increases
 - d. Arterial compliance increases**
191. Cognitive deficits, cerebellar atrophy and sensorimotor neuropathy all may be associated with
- a. Vitamin B12 deficiency
 - b. Multiple sclerosis
 - c. Alzheimer's disease
 - d. Alcoholism**
192. Which of the following bacteria is associated with food poisoning and myonecrosis?
- a. Clostridium difficile
 - b. Clostridium perfringens**
 - c. Clostridium botulinum
 - d. Clostridium tetani
193. A patient with fever of unknown origin has a positive blood culture for coagulase-negative staphylococcus that was drawn from the right arm and a negative culture drawn from the left arm. The patient most likely has
- a. Infective endocarditis
 - b. Pneumonia
 - c. A contaminated blood sample**
 - d. Osteomyelitis
194. Of the following disorders is caused by excessive production of somatotropin?
- a. Adrenogenital Syndrome
 - b. Cretinism
 - c. Acromegaly**

- d. Dwarfism
- 195. Diseases of thyroid gland have of recent been an area of attraction in general hospitals. Which of the following thyroid disorders is referred to as Hashimoto's disease?
 - a. Acute thyroiditis
 - b. Adrenal insufficiency
 - c. Chronic thyroiditis**
 - d. Graves' disease
- 196. A nurse knows that the chief cause of endemic goiter is
 - a. Autoimmune diseases
 - b. Goitrogenic substances
 - c. Iodine deficiency**
 - d. Thyroid gland tumors
- 197. All are characteristics of acute disease except
 - a. Sudden onset
 - b. May require intervention
 - c. Have period of remission and period of exacerbation**
 - d. Have duration
- 198. The procedure of compressing cricothyroid cartilage posteriorly against vertebra bodies during endotracheal intubation is called
 - a. Valsalva's maneuver
 - b. Sellick maneuver**
 - c. Heimlich maneuver
 - d. Leopold's maneuver
- 199. Subdiaphragmatic abdominal thrust performed to clear airway obstruction in the awake patient without ventilation is termed as
 - a. Leopold's maneuver
 - b. Pallach's maneuver
 - c. Valsalva maneuver
 - d. Heimlich maneuver**
- 200. There are a number of risk factors associated with coronary artery disease, which of the following is a modifiable risk factor?
 - a. Obesity**
 - b. Hereditary
 - c. Gender
 - d. Age
- 201. Which of the following conditions will most likely produce jaundice?
 - a. Crohn's disease

- b. **Gilbert's syndrome**
 - c. Hashimoto's disease
 - d. Zollinger-Ellison Syndrome
- 202. Scrotal hydrocele is the accumulation of serous fluid in the
 - a. **Tunica vaginalis**
 - b. Tunica vasculosa
 - c. Tunica albuginea
 - d. Tunica tunicalis
- 203. The most commonly isolated organism in prostatitis is
 - a. Staphylococcus aureus
 - b. Pyrogens
 - c. Mycoplasma
 - d. **E.Coli**
- 204. Cullen's sign is usually observed in a patient most likely to be diagnosed of
 - a. Peritonitis
 - b. Pericarditis
 - c. **Pancreatitis**
 - d. Diabetes mellitus
- 205. _____ are classified as unicellular, non-photosynthetic protists
 - a. Algae
 - b. Fungi
 - c. **Protozoa**
 - d. None of the above
- 206. Pertaining to fracture healing, which of the following is in correct order of the healing process?
 - a. Angiogenesis; cartilage calcification, Hematoma; cartilage removal
 - b. Cartilage calcification, Hematoma, cartilage removal, angiogenesis
 - c. **Hematoma, Angiogenesis, cartilage calcification, cartilage removal**
 - d. Hematoma, cartilage calcification; Angiogenesis, cartilage removal
- 207. Two major complications of portal hypertension a nurse should know are
 - a. Shock and paralysis
 - b. Generalized body oedema and hypertension
 - c. **Ascites and varices**
 - d. Liver damage and obstructive jaundice
- 208. A patient presents with hypertension, hypokalaemia and hypernatremia. Which of the following disorders could be responsible for these symptoms?
 - a. **Conn syndrome**
 - b. Syndrome of inappropriate antidiuretic hormone

- c. Excessive secretion of atrial natriuretic
 - d. Addison disease
209. Severe head trauma that results in interruption of the hypothal-amohypophyseal portal vessels would most likely lead to which of the following conditions?
- a. Hyperpigmentation
 - b. Hyperprolactinemia**
 - c. Acromegaly
 - d. Insulin resistance
210. Which of the following would least likely be a direct result of portal hypertension
- a. Ascites
 - b. Hyperestrinism**
 - c. Hypersplenism
 - d. Oesophageal varices
211. A sedentary man who spends most of his day sitting at a desk is developing increased girth around his waistline. He was not overweight as a child or during his adolescence. His obesity is most likely caused by
- a. Hyperplasia
 - b. Metaplasia
 - c. Hypertrophy**
 - d. Anaplasia
212. Michael shows the best adaptation with the new colostomy if he shows which of the following?
- a. Look at the ostomy site
 - b. Participate with the nurse in his daily ostomy care**
 - c. Ask for leaflets and contact members of ostomy support groups
 - d. Talk about his ostomy openly to the nurse and friends
213. Secondary anaemia is caused by
- a. Proliferative disorder of the myeloid stem cell**
 - b. Excessive iron absorption
 - c. Excessive production of erythropoietin
 - d. Decreased bone marrow activities
214. Signs of circulatory overload most likely would not include
- a. Tachycardia
 - b. Orthopnoea**
 - c. Bradycardia
 - d. Jugular vein distension
215. Morbidity is a term applied to people whose body mass index exceeds

- a. 24kg/m²
- b. 28kg/m²
- c. **30kg/m²**
- d. 40kg/m²

Situation: Match the items in column A with items in column B

Column A	Column B	Answers
216. Female hypogonadism	a. Down syndrome	C
217. 47XXY	b. Edward's syndrome	E
218. Trisomy 18	c. Turner's syndrome	B
219. Trisomy 13	d. Patau's syndrome	D
220. Trisomy 21	e. Klinefelter's syndrome	A
	f. Ewing's Syndrome	

From questions 221 to 237, tick to indicate statement is true or false. "Wrong answer attracts a penalty

- 221. Congenital lactase deficiency is an autosomal recessive genetic disorder.
 - a. **True**
 - b. False
- 222. Ice cream has more lactose content than yogurt
 - a. **True**
 - b. False
- 223. Regionalis most commonly occurs in the proximal ileum
 - a. True
 - b. **False**
- 224. The crampy pains in regional enteritis occur after meals
 - a. **True**
 - b. False
- 225. The prevalence of ulcerative colitis is highest in Caucasians
 - a. **True**
 - b. False
- 226. Erythrocyte sedimentation rate (ESR) is elevated in diverticulitis
 - a. **True**
 - b. False
- 227. Estimation of fluid intake and output remains the best method for ascertaining fluid balance and function of kidney
 - a. True
 - b. **False**
- 228. Cervical smear is a tool to detect pre-cancer and not metastasis

a. **True**

b. False

229. Colposcope is a microscope designed to allow examination of the cervix

a. **True**

b. False

230. Hepatitis A virus is a DNA virus

a. True

b. **False**

231. While females reach menopause at average of 45 years old,males' menopause occurs at average of 70 years

a. True

b. **False**

232. Using palm method of estimating the extent of burn, the patient's palm is approximately 1%

a. **True**

b. False

233. Obesity may increase the risk of back pain.

a. **True**

b. False

234. Lordosis is an increase in lumber curvature of the spine.

a. **True**

b. False

235. Neck retraction is one of the major signs of neonatal tetanus

a. **True**

b. False

236. A well approximated wound (a cut) heals by secondary intention

a. True

b. **False**

237. Iron deficiency anemia is the most common form of anaemia in many parts of the world

a. **True**

b. False

238. The primary site for hematopoiesis in adults is the **Red Bone marrow.**

239. Leukocytes are divided into two general categories ; **Granulocyte** and **Agranulocytes.**

240. The four clinical phases of acute renal failure are **Initial, Oligouria, Diuresis** and **Recovery.**

241. Surgery is a unique experience of planned physical alteration encompassing three phases, namely **Preoperative**, **Intraoperative** and **Postoperative**.

242. Four dimensions of preoperative teaching have been identified as important to clients, they are **information; including what will happen to the client**, **Psychosocial support to reduce anxiety** and **the role of the client & support people; skill training**.

243. Four common types of fevers are **intermittent**, **remittant**, **relapsing** and **Constant**.

244. Inflammatory bowel disease (IBD) refers to two chronic inflammatory GI disorder are **Regional enteritis** and **ulcerative colitis**.

245. Two types of gall bladder stones are **Pigment stone** and **cholesterol stone**.

246. Types of jaundice one can encounter in clinical practice include **hemolytic** and **hepatocellular obstructive**.

247. Abnormal respiratory rhythms create an irregular breathing pattern. Two abnormal respiratory rhythms are **Cheyne stroke** and **Biot's respiration**.

248. Hearing impairments are grouped into two **Conductive** and **Sensorineural**

249. DOTSC stands for **Directly Observed Therapy Short Course**.

250. State the full meaning of the following diagnostic procedures.

- i. **MRI – Magnetic Resonance Imaging**
- ii. **CBC – Complete Blood Count**
- iii. **CT scan – Computerized Tomography Scan**
- iv. **ELISA – Enzyme Linked Immunosorbent Essay**
- v. **ECG – Electrocardiography**
- vi. **EKG – Electroencephalography**
- vii. **BUN – Blood Urea Nitrogen**
- viii. **PCV – Pack Cell Volume**
- ix. **GFR – Glomerular Filtration Rate**
- x. **EDAA – Ethylene Diaminetetra Acetic Acid**
- xi. **IP – Intravenous Pyelography**
- xii. **ECR – Erythrocyte Sedimentation Rate**
- xiii. **ABG – Arterial Blood Gas**

- xiv. **EDD – Expected Date of Delivery**
- xv. **MCS – Microscopic Culture and Sensitivity**
- xvi. **MCV – Mean Corpuscular Volume**
- xvii. **MAP- Mean Arterial Pressure**
- xviii. **LMP – Last Menstrual Period**
- xix. **VDRL – Veneral Disease Research Laboratory**
- xx. **MVA – Manual Vacuum Aspiration**

PHARMACOLOGY

1. Which of the following estimate is more accurate for calculation of paediatric dose of drugs?
 - a. Age
 - b. Body weight
 - c. Body surface area**
 - d. Body mass index
2. The choice of appropriate route of a drug in a given situation depends on the following except
 - a. Physician's preference
 - b. Nature of drug
 - c. Patient related factors
 - d. Method of the disease transmission**
3. Mrs.Comfort, 60 year old woman who developed brain injury following RTA was admitted into a medical ward where doctor prescribed paracetamol P.R.N meaning
 - a. Three times daily
 - b. At once
 - c. As required**
 - d. As directed
4. One of the following is not a fat soluble vitamin
 - a. Vitamin A
 - b. Vitamin C**
 - c. Vitamin E
 - d. Vitamin K
5. The three most commonly used types of syringes a nurse must be familiar with are

- a. Tuberculin syringe, BCG syringe, & HBV syringe
 - b. Hypodermic syringe, insulin syringe, & tuberculin syringe**
 - c. DPT syringe, HBV syringe, & BCG syringe
 - d. Insulin syringe, BCG syringe, & HBV syringe
6. In buccal administration of medication, a drug is
- a. Placed under the tongue
 - b. Swallowed through the mouth
 - c. Held in the mouth against the mucous membranes of the cheek**
 - d. D. Injected into the muscles of the mouth
7. The home care nurse provides medication instructions to an older hypertensive client who is taking lisinopril (prinivil), 20mg orally daily. Which statement, if made by the client, indicates that further teaching is necessary?
- a. I can skip a dose once a week**
 - b. I need to change my position slowly
 - c. I take the pill after breakfast each day
 - d. If I get hard headache, I should call my doctor immediately
8. Cloxacillin, 200mg orally every 8hrs is prescribed for a child with an elevated temperature who is suspected of having a respiratory tract infection. The child weighs 17kg. The safe paediatric dosage is 50mg/kg/ day. The nurse determines that:
- a. The dose prescribed is safe**
 - b. The dose prescribed is too low
 - c. The dose prescribed is too high
 - d. There is not enough information to determine the safe dose
9. A Nurse knows that the use of standard dose in the marketing literature suggests that standard responses are the rule but in reality, there is considerable variation in drug response. Reasons for these variations include the following except
- a. Adherence
 - b. Drug formulation
 - c. Weight and age
 - d. Height**
10. A nurse is administering haloperidol(haldol) to a patient who have been admitted to the psychiatric unit. The nurse should know that an important use of this medication is to;
- a. Lift a long term depressive mood
 - b. Improve the patients' self concept
 - c. Provide a substitute for opioid abuse
 - d. Sedate agitated psychotic patient**
11. Children doses of a drug may be calculated from adult doses by using the following variables except