

Situation: Match the items in column A to those in column B

Column A	Column B	Answers
82. Biguanide	a. Tramadol	C
83. Aminoglycoside	b. Cefetazidime	F
84. Cephalosporins	c. Metformin	B
85. Sulfonylureas	d. Aspirin	E
86. Opioids	e. Glibenclamide	A
87. NSAID	f. Gentamicin	D
	g. Clozapine	
	h. Erythromycin	
88. Desipramine	a. Corticosteroid	C
89. Indapamide	b. Angiotensin Antagonist	D
90. Dexamethasone	c. Tricyclic Antidepressant	A
91. Losartan	d. Diuretic	B
92. Lisinopril	e. ACE Inhibitors	E
	f. Macrolide	

Situation: Baby Abigail, a 2 week old baby with infection of the umbilical cord (Omphalitis) is to have 62.5mg of i.m Ampiclox 6 hourly for 5 days. The vial comes in 250mg/2mls.

93. Calculate the volume of drug you will give baby Abigail for each dose

- a. 1ml
- b. 0.5ml**
- c. 2.5mls
- d. 0.1ml

94. How many vials of ampiclox will be needed for the total treatment?

- a. 2 vials
- b. 3 vials
- c. 4 vials
- d. 5 vials**

95. A patient has recently experienced a myocardial infarction (MI) within the last 4 hours. Which of the following medications would most likely be administered?

- a. Streptokinase**
- b. Atropine
- c. Acetaminophen
- d. Metformin

96. If not contraindicated, the nurse knows that a patient with diagnosis of Chlamydia can be treated with

- a. Amoxicillin
- b. Ampicillin
- c. Penicillin
- d. Tetracycline**

97. B-Lactam antibiotics include

- a. Nalidixic acid
- b. Chloramphenicol
- c. Streptomycin
- d. Penicillins**

98. Cephalosporins are effective broad-spectrum antibiotics which act by

- a. Inhibiting protein synthesis
- b. Causing leakage from cell membranes
- c. Inhibiting DNA gyrase
- d. Inhibiting cell wall synthesis**

Situation: Match the items in column A with those in column B

Column A	Column B	Answers
99. Acyclovir	a. Antifungi	C
100. Retinol	b. Quinolones	E
101. Amphotericin B	c. Anti Herpes	A
102. Norfloxacin	d. Oral hypoglycemia	B
103. Glipizide	e. Vitamin A	D
	f. Antianginal	

104. When administering i.m injection to an infant, the nurse in charge should use which site?

- a. Deltoid
- b. Dorsogluteal
- c. Ventrogluteal
- d. Vastus lateralis**

105. The first effective antipsychotic drug was

- a. Chlorpromazine**
- b. Thioridazine
- c. Haloperidol
- d. Trifluoperazine

106. Levamisole is used as an anticancer drug in the treatment of

- a. Breast Cancer
- b. Colorectal cancer**
- c. Hodgkin's disease

- d. Leukemia
- 107. Which of the following is not anti-metabolic group of chemotherapeutic agent?
 - a. 6-Mercaptopurine
 - b. Methotrexate
 - c. 5-fluorouracil
 - d. **Vindesine**
- 108. Aspirin, NSAIDS and Acetaminophen act by
 - a. Inhibiting impulses to brain
 - b. **Inhibiting the synthesis of prostaglandins**
 - c. Competitively blocking the neuronal terminal
 - d. Releasing endorphin
- 109. Which of the following drugs on over dosage is known to cause liver failure?
 - a. **Acetaminophen**
 - b. Ibuprofen
 - c. Piroxicam
 - d. Ketorelac
- 110. The enzyme involved in the therapeutic action of NSAIDS is
 - a. **Cyclo-oxygenase**
 - b. Mono amine oxidase
 - c. Cytochrome
 - d. Lipocortin-1
- 111. Inadequate the pregnant woman of which vitamin may cause neural tube defects?
 - a. Niacin
 - b. Riboflavin
 - c. **Folic acid**
 - d. Thiamine
- 112. Which of the following antitubercular drugs is bacteriostatic?
 - a. Rifampin
 - b. Pyrazinamide
 - c. Streptomycin
 - d. **Ethambutol**
- 113. Drugs affecting coagulation include the following except
 - a. Enoxaparin
 - b. Warfarin sodium
 - c. **Hydroxocobalamin**
 - d. Heparin
 - e.

Match the items in column A with those in column B

Column A	Column B	Answers
114.Precursor of B1	a.Pyridoxine	B
115. Vitamin E	b.Sulbutiamine	E
116. Vitamin B6	c.Riboflavin	A
117. Vitamin B2	d.Phytomenadione	C
118. Vitamin K1	e.Tocopherol	D
	f.Thiamine	

119. Which of the following is among the first line antitubercular drugs

- a. Ethionamide
- b. Clarithromycin
- c. Rifabutin
- d. Pyrazinamide**

120. One of these drugs is not a first line antitubercular drug

- a. Ethambutol
- b. Rifampin
- c. Streptomycin
- d. Thiacetazone**

121. Directly observed treatment, short course should take minimum of

- a. 4 months
- b. 6 months**
- c. 1 year
- d. 2 months

122. How should the nurse prepare a suspension before administration?

- a. By diluting it with normal saline solution?
- b. By diluting it with 5% dextrose solution
- c. By shaking it so that all the drug particles are dispersed uniformly**
- d. By crushing remaining particles with a mortar and pestle.

123. Dr. Ukwé prescribes corticosteroids for a child with nephrotic syndrome. The primary purpose of administering corticosteroids to this child is to

- a. Increase blood pressure
- b. Reduce inflammation
- c. Decrease proteinuria**
- d. Prevent infection

124. A commonly used NSAID 'ibuprofen' belongs to the subclass of

- a. Salicylates
- b. Pyrazolone derivatives**

- c. Indole derivatives
- d. Propionic acid derivatives**

125. Ciprofloxacin is not recommended for use in children less than

- a. 10 years
- b. 12 years
- c. 6 years
- d. 18 years**

126. The adult recommended dose of Omeprazole for treatment of Acute gastritis is

- a. 20mg once daily**
- b. 40mg every 12 hours
- c. 60mg daily
- d. 10mg 6 hourly

127. The vitamin necessary for the production of Prothrombin in the liver is

- a. Vitamin A
- b. Vitamin B₁₂
- c. Vitamin D
- d. Vitamin K**

128. Ciprofloxacin is a drug belonging to which group of antibiotics?

- a. Fluoroquinolone**
- b. Macrolide
- c. Cephalosporin
- d. Aminoglycoside

129. Nitrofurantoin is a bactericidal agent used for the treatment of

- a. Respiratory tract infections
- b. Skin infections
- c. Staphylococcal infections
- d. Urinary tract infections**

130. Which of the following is not in macrolide group of antibiotics?

- a. Erythromycin
- b. Azithromycin
- c. Roxithromycin
- d. Gentamicin**

131. Which of the following is a bacteriostatic, which acts by reversible inhibition of ribosomal protein synthesis?

- a. Vancomycin
- b. Tetracyclines**
- c. Cephalosporins
- d. Isoniazid

132. Which of the following antibiotics is specifically effective against anerobic bacteria?

- a. Amphotericin B
- b. Erythromycin
- c. Metronidazole**
- d. Gentamicin

133. An example of H₂receptor antagonist is

- a. Ranitidine**
- b. Omeprazol
- c. Metronidazole
- d. Ribavirin

134. Which of the following drugs could best be used for open angle glaucoma?

- a. Timolol**
- b. Catapril
- c. Omeprazole
- d. Propranolol

135. The formula: Age in Months x Adult Dose for calculation of drug dosages for Children was developed by

- a. Fried**
- b. Young
- c. Clarke
- d. Matthews

136.Which of the following is not true about a drug?

- a. Every drug is capable of producing adverse effect
- b. Whenever a drug is given,a risk is taken
- c. There is no difference between a drug and poison**
- d. Side effect is an unavoidable pharmacodynamic effects at therapeutic doses

137.One of the following is not included among the type A (Aug-mented) adverse drug reactions?

- a. Side effects
- b. Consequences of drug withdrawal
- c. Idiosyncrasy**
- d. Toxic effects

138. Unwanted but often unavoidable but predictable pharmacody-namics effects that occur at therapeutic doses is called

- a. Side effects**
- b. Toxic effects
- c. Secondary effects
- d. Idiosyncratic effects

139. Suppression of bacterial flora by tetracyclines paves way for super-infections. This is an illustration of _____ of a drug

- a. Side effects
- b. Toxic effects
- c. Secondary effects**
- d. Allergic effects

140. Emesis/gastric lavage is not recommended if the patient presents more than

- a. 4 hours after ingestion of poison
- b. 2 hours after ingestion of poison**
- c. 12 hours after ingestion of poison
- d. 8 hours after ingestion of poison

141. An increased response of the body system to a particular drug is called drug

- a. Hypersensitivity
- b. Supersensitivity**
- c. Cross sensitivity
- d. Idiosyncrasy

142. Which of the following drugs is associated with feminizing effects in men during chronic therapy?

- a. Cimetidine**
- b. Ranitidine
- c. Famotidine
- d. Omeprazole

143. An otherwise healthy, pregnant woman with the recent onset of urinary frequency, urgency and dysuria, and evidence of bacteriuria should be treated with

- a. Ciprofloxacin
- b. Norfloxacin
- c. Trimethoprim-sulfamethoxazole
- d. Cephalexin**

144. The drug 'Ondansetron' can best be described as an

- a. Analgesic
- b. Antiemetic**
- c. Antibiotic
- d. Antineoplastic

145. Terbinafine is an

- a. Antiinflammatory
- b. Antibiotic
- c. Antiviral drug
- d. Antifungal drug**

146.A needle has three discerning parts which are the hub, canulla or shaft and bevel.

147.The three variable characteristics of needles for injection are length of the bevel, length of the shaft and diameter (guage) of the shaft.

148.An adverse drug reaction may be defined as Unwanted or unintended effects of medicine.

149. Two systems of measurement currently in use for calculation of dosages are Metric and Apothecary.

150. Gentamin I.M.was prescribed to be given 20mgbd x 8 doses to a patient; if the ampoule available is 80mg/2ml, calculate (i) the amount for each dose (ii) How many ampoules will be required

0.5ml: 2 ampoules.

COMMUNITY HEALTH NURSING

1. Which of the following plasmodium species is capable of causing malignant tertian fever?
 - a. **Plasmodium falciparum**
 - b. Plasmodium vivax
 - c. Plasmodium malariae
 - d. Plasmodium oval
2. Oralydration fluid is better used within
 - a. 12hours
 - b. 18 hours
 - c. 36 hours
 - d. **24 hours**
3. The concept in I.E.C. includes the following except
 - a. Information
 - b. Education
 - c. Communication
 - d. **Counseling**
4. _____ is concerned with the study of factors that influence the occurrence and distribution of disease, defects, disability or death which occurs in groups or aggregation of individuals

- a. **Epidemiology**
 - b. Demographics
 - c. Vital signs
 - d. Health statistics
5. Which of the following is the backbone in disease prevention?
- a. **Epidemiology**
 - b. Demographics
 - c. Vital statistics
 - d. Health statistics
6. The following are all functions of the nurse in vital statistics except
- a. **Consolidate data**
 - b. Collects data
 - c. Analyze data
 - d. Tabulate data
7. The following are notifiable diseases that need to have a tally sheet in data reporting except?
- a. Hypertension
 - b. Bronchiolitis
 - c. Chemical poisoning
 - d. **Accidents**
8. Which of the following requires reporting within 24 hrs?
- a. Neonatal tetanus
 - b. **Measles**
 - c. Hypertension
 - d. Tetanus
9. Which of the following is a point source epidemic?
- a. Dengue
 - b. Malaria
 - c. **Contaminated water source**
 - d. Tuberculosis
10. All but one is a characteristic of a point source epidemic
- a. The spread of the disease is caused by a common vehicle
 - b. The disease is usually caused by contaminated food
 - c. **There is a gradual increase of cases**
 - d. Epidemic is usually sudden
11. The microorganism monitored in cases of contaminated water is
- a. Vibrio cholera
 - b. **Escherichia coli**

- c. Entamoeba histolytica
 - d. Coliform test
12. Dengue increase in number during June, July and August. This pattern is called
- a. Epidemic
 - b. Endemic
 - c. Cyclical**
 - d. Secular
13. Mang Ambo approaches you for counseling. You are an effective counselor if you
- a. Give good advice to Mang Ambldento
 - b. Ambo's problems
 - c. Convince Mang Ambo to follow your advice
 - d. Help Mang Ambo identify his problems**
14. Preparatory phase is the first phase in organizing the community. Which of the following is the initial step in the preparatory phase?
- a. Area selection**
 - b. Community profiling
 - c. Entry in the community
 - d. Integration with the people**
15. In choosing a leader in the community during the organizational phase, which among these people will you choose?
- a. Miguel Zobel, 50 years old, rich and famous**
 - b. Rustom, 27 year old actor
 - c. Mang Ambo, 70, willing to work for the desired change
 - d. Ricky 30 years old, influential, and willing to work for the desired change.
16. Which of the following is the correct sequence in health education? (1) Information (2) Communication (3) Education
- a. 1,2,3**
 - b. 3,2,1
 - c. 1,3,2
 - d. 3,1,2
17. The health status of the people is greatly affected and determined by which of the following?
- a. Behavioural factors
 - b. Socio economic factors**
 - c. Political factors
 - d. Health statistics
18. One of these refers to systematic study of vital events such as births, illnesses, marriages, divorces and deaths