

- a. 2
 - b. 4
 - c. 5**
 - d. 10
49. The term foetus papyraceous refers to death of one foetus in the
- a. First trimester of pregnancy
 - b. Second trimester of pregnancy**
 - c. Third trimester of pregnancy
 - d. First stage labour
50. Infant mortality refers to babies who die
- a. An unexpected cot death
 - b. In the first month of life
 - c. In the first year of life**
 - d. As a result of infection
51. _____ is the term used for twins conceived as a result of two coital acts in different menstrual cycles
- a. Superfecundation
 - b. Superfetation**
 - c. Superovulation
 - d. Superfertilization
52. _____ is the term used when twins are conceived from sperm from different men
- a. Superfecundation**
 - b. Superfetation
 - c. Superovulation
 - d. Superfertilization

Situation: Mrs. Aliyu, an unbooked primipara at 36weeks of gestation was rushed to your ward with the history of convulsions at home. On examination, temperature, pulse & respiration were all normal. Blood pressure was 160/110mmHg and bleeding per vaginum. Questions 53-55 refer to this.

53. What is the possible diagnosis of Mrs. Aliyu?
- a. Pregnancy induced hypertension
 - b. Pre-eclampsia
 - c. Transient ischaemia of pregnancy
 - d. Eclampsia**
54. The following are possible complications of Mrs. Aliyu condition except
- a. Cardiac failure
 - b. Renal failure
 - c. Eclampsia**

- d. Disseminated intravascular coagulation
55. The proper drug to be used in controlling Mrs. Aliyu's blood pressure is
- a. Frusemmide
 - b. Hydrochlorothiazide
 - c. Hydralazine**
 - d. Verapamil
56. The last secondary sexual characteristic to develop in the female adolescent is
- a. Menstruation**
 - b. Breast enlargement
 - c. Ovulation
 - d. Pubic hair growth
57. The preconception period refers to time span of anything from before conception
- a. 3 months to 1 year**
 - b. Puberty to 1 year
 - c. 6 months to 1 year
 - d. Puberty to 3 months
58. The purplish streaks that sometimes appear on the breasts and abdomen during pregnancy are called
- a. Hegar line
 - b. Striae terminalis
 - c. Linear nigra
 - d. Striae gravidarum**
59. Growth monitoring programme is for children
- a. Below 2 years
 - b. From birth to 5 years**
 - c. Between 3-5 years
 - d. Above 5 years
60. Factors that contribute to ectopic gestation include the following except
- a. Infection of the fallopian tube
 - b. Severe malaria in pregnancy**
 - c. Peritubal adhesions
 - d. History of previous abortion
61. The term aspermia simply means
- a. Absence of spermatozoa from the ejaculate**
 - b. Reduced sperm density
 - c. Absence of semen
 - d. Decreased sperm motility

62. Pathological causes of primary amenorrhea include the following except

- a. **Pregnancy**
- b. Testicular feminization syndrome
- c. Vaginal agenesis
- d. Imperforate hymen

63. A 50 year old woman with diagnosis of Ca of cervix was admitted in a ward where Nurse Josephine is in-charge. The nurse trying to teach the junior colleagues about the cause of Ca of cervix should mention

- a. HIV
- b. **HPV**
- c. EBV
- d. Herpes simplex

Situation: In a home visit done by the nurse, she suspects that her client (a wife) and her child are victims of abuse

64. Which of the following is the most appropriate for the nurse to ask?

- a. **Are you being threatened or hurt by your partner?**
- b. Are you frightened of your partner?
- c. Is something bothering you?
- d. What happens when you and your partner argue?

65. The wife admits that she is a victim of abuse and opens up about her persistent distaste for sex. This sexual disorder is:

- a. **Sexual desire disorder**
- b. Sexual arousal disorder
- c. Orgasm disorder
- d. Sexual pain disorder

66. What would be the best approach for a wife who is still living with her abusive husband?

- a. **Here's the number of a crisis centre that you can call for help**
- b. It is best to leave your husband
- c. Discuss this with your family
- d. Do not allow yourself to be treated this way

67. The primary nursing intervention for a victim of child abuse is

- a. Assess the scope of the problem
- b. Analyze the family dynamics
- c. **Ensure the safety of the victim**
- d. Teach the victim coping skill

68. The commonest causes of miscarriage are

- a. Maternal illness
- b. **Chromosome abnormalities in the foetus**

- c. Uterine abnormalities
 - d. Hard labour
69. The term ectopic pregnancy refers to
- a. Any pregnancy taking place in the fallopian tube
 - b. Any pregnancy occurring outside the uterine cavity**
 - c. One in which after fertilization, the foetus develops in the ovary
 - d. Any pregnancy taking place outside the abdominal cavity
70. The commonest site of extrauterine implantation is the
- a. Ovary
 - b. Cervical canal
 - c. Uterine tube**
 - d. Abdominal cavity
71. The normal discharge from the vaginal on 6th day post delivery
- a. Lochia rubra
 - b. Lochia serosa**
 - c. Lochia alba
 - d. Lochia exudates
72. The posterior fontanelle closes at
- a. 18 months
 - b. 6 months
 - c. 18 weeks
 - d. 6 weeks**
73. The reproductive organs return to the pre-pregnant state by the
- a. 12th week after birth
 - b. 10th week after birth
 - c. 8th week after birth
 - d. 6th week after birth**
74. Pseudomenses may occur in female newborn due to withdrawal of
- a. Maternal progesterone
 - b. Maternal oestrogen**
 - c. Maternal lactogen
 - d. Human chorionic gonadotrophin
75. The colour of babies with pigmented skin is best assessed by inspecting the colour of
- a. Mucous membrane**
 - b. Pinna
 - c. Rectum
 - d. Elbow
76. The following are obstetric emergencies except

- a. Cord prolapse
 - b. Uterine rupture
 - c. Amniotic fluid embolism
 - d. Rupture of the membrane**
77. Painless sore on the genital is diagnostic of
- a. Syphilis**
 - b. Gonorrhoea
 - c. Chlamydia infection
 - d. Herpes genitalis
78. Puerperal pyrexia could result from the following except
- a. Breast engorgement
 - b. Urinary tract infection
 - c. Varicose vein**
 - d. Genital tract infection
79. Which of the following is a commonest cause of maternal mortality in Nigeria
- a. Amniotic fluid embolism
 - b. Abortion**
 - c. Breast cancer
 - d. Pregnancy induced hypertension
80. Which of the following could be a complication of pregnancy induced hypertension?
- a. Placenta praevia
 - b. Abortion
 - c. Dystocia
 - d. Abruptio placentae**
81. How many cubes of sugar would you use in preparing two (2) cocacola 35cl bottle filled water for oral rehydration solution?
- a. 10 cubes
 - b. 8cubes
 - c. 5cubes**
 - d. 3cubes
82. Family planning was identified as key to the control of population in the world conference held at
- a. United Kingdom in 1964**
 - b. Bucharest in 1974
 - c. Mexico in 1984
 - d. Cairo (Egypt) in 1994
83. The first to function in an embryo is the
- a. Respiratory system

- b. Gastrointestinal system
 - c. Cardiovascular system**
 - d. Nervous system
84. The fetal erythrocytes have a life span of
- a. 120 days
 - b. 110 days
 - c. 100 days
 - d. 90 days**
85. Connecting the umbilical vein to the inferior vena cava is the
- a. Foramen ovale
 - b. Ductus venosus**
 - c. Ductus arteriosus
 - d. Hypogastric arteries
86. A newborn's respiratory rate may be described as tachypnoea when it is above
- a. 80 b/m
 - b. 60 b/m**
 - c. 40 b/m
 - d. 20b/m
87. The placenta is completely formed and functioning weeks after fertilization
- a. 10**
 - b. 12
 - c. 15
 - d. 20
88. Contraceptive method(s) most suitable for unmarried adolescent is
- a. Total abstinence**
 - b. Oral contraceptive pills
 - c. Barrier methods and spermicides
 - d. Lippes loop
89. In Billings method of contraception, one of the following is used
- a. The pills
 - b. Cervical mucus**
 - c. Condom
 - d. IUCD
90. In type 2 placenta praevia, the placenta is
- a. At the upper uterine segment
 - b. Partially located in the lower segment near the internal os**
 - c. Located over the internal os but not centrally
 - d. Located centrally over the internal os

91. An 18-year-old woman comes to the physician's office for a routine prenatal checking at 34 weeks gestation. Abdominal palpation reveals the fetal position as right occipitoanterior (ROA). At which of the following sites would the nurse expect to find the fetal heart tone?
- a. Below the umbilicus, on the mother's left side
 - b. Below the umbilicus, on the mother's right side**
 - c. Above the umbilicus, on the mother's left side
 - d. Above the umbilicus, on the mother's right side.
92. Sudden infant death syndrome (SIDS) is one of the most common causes of death in infants. At what age is the diagnosis of SIDS most likely?
- a. At 1 to 2 years of age
 - b. At 1 week to 1 year of age peaking at 2 to 4 months**
 - c. At 6 to 8 weeks of age
 - d. At 6 to 1 year, peaking at 10 months
93. The trophoblasts begin to make human chorionic gonadotropin (hcg) by
- a. 8 days**
 - b. 16 days
 - c. 24 days
 - d. 32 days
94. In most female adolescent, the time from the appearance of breast buds to the first menstruation is approximately
- a. 2 months
 - b. 6 months
 - c. 12 months
 - d. 24 months**
95. A newborn infant who extends its large toe in response to stroking of the lateral surface of the sole of its foot is demonstrating
- a. Babinski reflex**
 - b. Rooting reflex
 - c. Grasp reflex
 - d. Moro reflex
96. A pregnancy is defined as being postterm if it exceeds
- a. 280 days
 - b. 287 days
 - c. 294 days**
 - d. 301 days
97. Which of the following disorders is associated with dysfunctional uterine bleeding?
- a. Endometriosis

- b. Secondary dysmenorrhoea
 - c. Cervical cancer
 - d. Anovulatory cycles**
98. Which of the following is not a cause of secondary amenorrhoea?
- a. Polycystic ovarian syndrome
 - b. Hyperprolactinaemia
 - c. Delay puberty**
 - d. Premature menopause
99. The presence of fragments of endometrial tissue at sites in pelvis outside the uterus is called a
- a. Dyspareunia
 - b. Adenomyosis
 - c. Endometritis
 - d. Endometriosis**
100. Surgical removal of one or more fibroids from the uterus is called
- a. Hysterectomy
 - b. Myomectomy**
 - c. Cervical sweep
 - d. Oophorectomy
101. An accumulation of menstrual blood in the uterus is called
- a. Haematometria
 - b. Haemavesicus
 - c. Haemamyoma
 - d. Haematocolpos**
102. The combined oral contraceptive pill
- a. Is taken continuously for 28 days
 - b. Has a pearl index of 10
 - c. Can be used as a form of post-coital contraception
 - d. Can be used to reduce menorrhagia and dysmenorrhoea effectively**
103. A female sterilization
- a. Has a lower failure rate than a vasectomy
 - b. Can never be reversed
 - c. Can be done at any point during the patient's cycle**
 - d. Can be carried out on multiparous women
104. On assessment of semen fluid analysis, which of the following is not correct?
- a. Azoospermia indicates a sperm count more than 5 million/ml**
 - b. Asthenozoospermia indicates abnormal motility of >60%
 - c. Teratozoospermia indicates abnormal forms of >30%

- d. Abstinence should be maintained for 3 days prior to productivity of the sample
105. In the presence of a positive pregnancy test and vaginal bleeding, the following are possible differential diagnoses if the cervix is closed except
- a. Ectopic pregnancy
 - b. Incomplete miscarriage**
 - c. Threatened miscarriage
 - d. Complete miscarriage
106. Cations released from copper-releasing intrauterine devices(IUDs)
- a. Prevent ovulation
 - b. Make uterus unsuitable for implantation
 - c. Increase phagocytosis of sperms
 - d. Suppress sperm motility**
107. The permissible use of the technique 'amniocentesis' is for
- a. Detecting any genetic abnormality**
 - b. Detecting sex of the unborn foetus
 - c. Artificial insemination
 - d. Transfer of embryo into the uterus of a surrogate mother
108. The most significant risk factor for breast cancer is
- a. Age the patient**
 - b. Number of pregnancies
 - c. Age at onset of menarche
 - d. Family history of breast cancer
109. Gynecomastia in a 25-year-old man would most likely be associated with a primary cancer located in the
- a. Testicle**
 - b. Lung
 - c. Small bowel
 - d. Thyroid gland
110. Which of the following factors most clearly rules out an organic cause of male impotence?
- a. Normal serum thyroxine
 - b. Nocturnal erections**
 - c. Normal serum testosterone
 - d. Normal serum gonadotropins
111. Which of the following disease causing organisms associated with anal intercourse causes visible lesions with no pain or discomfort in the anus or anal canal?
- a. Human papillomavirus
 - b. Treponema pallidum**

- c. Herpes genitalis
 - d. Neisseria gonorrhoea
112. Which of the following conditions would least likely be associated with dysmenorrhoea?
- a. **Oral contraceptive use**
 - b. Increased uterine prostaglandins
 - c. Adenomyosis
 - d. Endometriosis
113. Sperm pass through each of the following structures except the
- a. Epididymis
 - b. Seminiferous tubule
 - c. **Seminal vesicle**
 - d. Ampulla of ductus deferens
114. Which of the following site-breast disease relationships is not correct?
- a. Nipple-paget disease
 - b. Lactiferous duct-intraductal papilloma
 - c. **Large ducts-sclerosing adenosis**
 - d. Stroma-fibroadenoma
115. Which one of the following clinical situations would be least likely to result in a newborn that is small for gestational age?
- a. Premature rupture of the membrane
 - b. Abruptio placenta
 - c. Pre-eclampsia
 - d. **Gestational diabetes mellitus**
116. The most common benign tumour in women is most commonly located in the
- a. Ovary
 - b. Breast
 - c. **Uterus**
 - d. Stomach
117. In a normal newborn infant, the palmar grasp reflex disappears at approximately
- a. **2 months**
 - b. 4 months
 - c. 6 months
 - d. 9 months
118. I.M.C.I means **Integrated Management of Childhood Illness.**
119. The two types of menstruation are **Ovular** and **Anovular** .
120. The average blood loss during menstruation is **50** to **150ml.**