

194. Two types of liver biopsy a nurse may encounter in the course of practice are Open liver biopsy and Closed liver biopsy.

195. Percutaneous liver biopsy is also called Closed liver biopsy.

196. Florence Nightingale's achievements in improving the standards for the care of war casualties in the Crimean war earned her the title : Lady with the lamp.

197. The two kinds of body temperature are Core and surface.

198. In determining the infusion rate, the number of drops in 1 milliliter is called the Drip factor.

199. The following abbreviations represent the various care givers associations. State the full meaning of each

- i. **NANNM – National Association of Nigerian Nurses and Midwives.**
- ii. **NMCN – Nursing and Midwifery Council of Nigeria.**
- iii. **GNAN – Graduate Nurses Association of Nigeria.**
- iv. **NAFDAC – National Agency for Food and Drug Administration.**
- v. **NDLEA – National Drug Law Enforcement Agency.**
- vi. **MDCN – Medical and Dental Council of Nigeria.**
- vii. **NANDA – North American Nursing Diagnosis Association.**

MEDICAL-SURGICAL NURSING

Situation: A 30 year old man was brought to medical outpatient department where a nurse measured the BP and observed that when the blood pressure cuff on the arm is inflated 20mmHg above his sys-tolic pressure for 2 minutes, carpal spasm occurs. Use this information to answer questions 1 and 2


1. The sign observed by the nurse is best described as
 - a. Chvostek's sign
 - b. Rovsing's sign
 - c. Trousseau's sign**
 - d. Koplik's sign
2. The observation made is mainly seen in patients suffering from
 - a. Hypernatremia
 - b. Meningitis
 - c. Hypokalaemia

d. Hypocalcemia

3. An 80 year old male client is admitted to the hospital with a diagnosis of pneumonia. Nurse Oliver learns that the client lives alone and hasn't been eating nor drinking. When assessing him for dehydration, nurse Oliver would expect to find

- a. Hypothermia
- b. Hypertension
- c. Distended neck veins

d. Tachycardia



Situation: Use the sketch provided to answer the next three questions

4. The sketch above represents
- a. **Negative feedback**
 - b. Positive feedback
 - c. A and b
 - d. None of the above
5. The hormone x is the
- a. ACTH
 - b. Vasopressin**
 - c. Calcitonin
 - d. Insulin
6. In addition to water re-absorption, there is also increased
- a. K⁺
 - b. Na⁺**
 - c. Ca⁺
 - d. Cl⁻

7. An effusion of blood into a joint is called
- Haematemesis
 - Haemarthrosis**
 - Haematocolpos
 - Haemachromatosis
8. The following are true causes of oedema except
- Increased venous hydrostatic (blood) pressure
 - Increased plasma osmotic pressure**
 - Impaired lymphatic drainage
 - Increased small vessel permeability
9. A patient with extensive third degree burns begins to look ill. Test results indicate inadequately circulating blood volume due to fluid shifts and a loss of plasma proteins. The nurse alerts a doctor because she suspects
- Cardiogenic shock
 - Distributive shock
 - Hypovolemic shock**
 - Obstructive shock
10. A patient's chart indicates a history of hyperkalemia. Which of the following would you not expect to see with this patient if this condition was acute?
- Decreased heart rate
 - Parathesias
 - Muscle weakness of the extremities
 - Migraines**
11. The nurse would expect to find an improvement in which of the blood values as a result of dialysis treatment?
- High serum creatinine levels**
 - Low haemoglobin
 - Hypocalcaemia
 - Hypokalaemia
12. Haemodialysis is the most common method of dialysis. Which of the following is not among the principle on which haemodialysis is based?
- Tubal secretion**
 - Ultra filtration
 - Diffusion
 - Osmosis
13. Ultrafiltration occurs in peritoneal dialysis through an osmotic gradient created by using a dialysate fluid with a
- Lower salt concentration

- b. **Higher glucose concentration**
 - c. Lower glucose concentration
 - d. Higher molecule (water) concentration
- 14. A nurse understands that _____ is a type of diarrhoea caused by increased production and secretion of water and electrolytes by the intestinal mucosa into the intestinal lumen.
 - a. Osmotic diarrhoea
 - b. Mixed diarrhoea
 - c. **Secretory diarrhoea**
 - d. Pressure disease
- 15. A patient's chart indicates a history of ketoacidosis, which of the following would you not expect to see with this patient if this condition was acute?
 - a. Vomiting
 - b. Extreme thirst,
 - c. **Weight gain**
 - d. Acetone sbreathes smell.
- 16. The most common site for a pressure ulcer to develop is the
 - a. Occipital area
 - b. Sacrum
 - c. **Shoulder**
 - d. Ankle
- 17. The key factor contributing to the cause of a pressure ulcer is
 - a. Moisture
 - b. Incontinence
 - c. **Pressure**
 - d. Malnutrition
- 18. The following factors may contribute to formation of pressure ulcer except
 - a. Decreased mental status
 - b. **Strenuous exercise**
 - c. Immobility
 - d. Inadequate nutrition.
- 19. An older patient sits and slumps in her chair most of the day. She is most likely to develop a pressure ulcer because of
 - a. Malnutrition
 - b. **Shearing force**
 - c. Edema
 - d. A chronic disease
- 20. A priority nursing action to prevent a patient from developing a pressure ulcer is

- a. Use waterproof material on the bed
 - b. Massage any reddened area frequently
 - c. Use an air inflated ring to relieve pressure on an area
 - d. Use a mild soap when cleansing the skin**
21. After an initial assessment, the nurse documents the presence of a reddened area that has abraded and blistered. According to recognized staging systems, this ulcer is classified as
- a. Stage I
 - b. Stage II**
 - c. Stage III
 - d. Stage IV
22. Nurse Bridget assesses a stage III pressure ulcer. She has observed
- a. Redness that persists when pressure is relieved
 - b. An open lesion with subcutaneous tissue exposed**
 - c. A necrotic area extending through the fascias to bone
 - d. A reddened area with an abrasion
23. A student nurse was to assess a patient for increased risk for developing pressure sore. She should use
- a. Glasgow scale
 - b. Apgar scoring system
 - c. Braden Scale**
 - d. Bishop's score.
24. A patient was assessed for the risk of pressure sore using the scale/ system in question 23. If the nurse elicits that the patient is constantly moist, slightly limited in sensory perception, chair fast, completely mobile and probably adequately fed with no apparent problem of friction and shear, then the total score of the patient is
- a. 23
 - b. 18
 - c. 15**
 - d. 21
25. Oluchi, a nurse in female surgical ward was reporting on possible complications of pressure ulcers, she reports the most serious complications of pressure ulcers as:
- a. Potential Infection**
 - b. Haemorrhage
 - c. Disability
 - d. Ischaemia
26. Which stage of pressure ulcer development does the ulcer extend into the subcutaneous tissue?

- a. Stage I
 - b. Stage II
 - c. Stage III**
 - d. Stage IV
27. Which of the following actions can the nurse institute independently when caring for a male client with a 3cm stage I pressure ulcer on the coccyx?
- a. Massaging the area with an astringent every 2 hours
 - b. Applying an antibiotic crecream to the area three times per day
 - c. Using normal saline solution to clean the ulcer and applying a protective dressing as necessary**
 - d. Using a providone iodine wash on the ulceration three times per day
28. An autoimmune disease of which autoantibodies destroy intrinsic factor and parietal cells in the stomach is the
- a. Aplastic Anaemia
 - b. Pernicious anaemia**
 - c. Iron deficiency anaemia
 - d. Hypoplastic anaemia
29. Impaired erythrocyte production may lead to the following types of anaemia except
- a. Iron deficiency anaemia
 - b. Hypoplastic anaemia**
 - c. Megaloblastic anaemia
 - d. Haemolytic anaemia

Match the items in column A with those in column B

Column A	Column B	Answers
30. Normocytic	a.Cells smaller than normal	E
31. Microcytic	b. Cells paler than normal	A
32. Macrocytic	c.Rate of cell destruction raised	D
33. Hypochromic	d. Cells bigger than normal	B
34. Haemolytic	e.Cells normal sized	C
	f. Rate of cell formation raised	

35. The following are associated with hypoproliferative anaemia except
- a. Iron deficiency
 - b. Sickle cell**
 - c. Folate deficiency
 - d. Megaloblastic anaemia

36. The nurse is assessing a patient complaining of weakness and fatigue. Diagnostic laboratory studies indicate a decrease in vitamin B12 absorption. The nurse understands that anaemia related to a decrease in vitamin B12 absorption is
- Hemolytic anaemia
 - Iron deficiency anaemia
 - Megaloblastic anaemia
 - d. Pernicious anaemia**
37. In normal blood, the total leukocyte count per cubic millimeter is between
- a. 5,000-10,000 cells**
 - 20,000-24,000 cells
 - 23,000-40,000 cells
 - 32,000-38,000 cells
38. The red bone marrow located within bone cavities produces red blood cells, white blood cells and platelets through a process referred to as
- Erythropoiesis
 - Neucopoiesis
 - c. Hematopoiesis**
 - Hematogenesis.
39. If plasma protein levels fall, because of either reduced production or loss from blood vessel
- Osmotic pressure of the blood will increase
 - b. Oedema will result**
 - Fluid moves from tissue into the blood
 - The kidney will increase reabsorption rate.
40. Which of the following blood products for transfusion does not need to be typed and cross matched?
- Whole blood
 - b. Fresh Frozen plasma**
 - Packed red blood cells
 - None of the above
41. Following a heavy bleeding, it was discovered that a patient has lost a large amount of plasma protein. Which of the following will be used in restoring the blood volume?
- Packed red blood cells
 - Crystalloids
 - Normal saline
 - d. Colloids**
42. During a dressing change, the nurse observed what to be reddish pink tissue in the wound. This is most likely

- a. A sign of infection
 - b. Eschar
 - c. Exudate
 - d. Granulation tissue**
43. Which of the following terms should be used to document wound drainage that is thick, odorous, and green?
- a. Serous
 - b. Sanguineous
 - c. Serosanguineous
 - d. Purulent**
44. A client has a pressure ulcer with a shallow, partial skin thickness, eroded area but no necrotic areas. The nurse would treat the area with which of the following dressing?
- a. Alginate
 - b. Dry gauze
 - c. Hydrocolloid**
 - d. No dressing is indicated
45. The RYB as colour code of wounds stand respectively for ____, ____ and ____
- a. Red, Yellow & Blue
 - b. Red, Yellow & Black
 - c. Rainbow, Yellow & Blue**
 - d. Rainbow, Yellow & Black
46. Rose decided to use the RYB colour code of wounds to care for Mrs. Jones wound which is yellow. Which of the following should best be done on the wound?
- a. Change the dressing as infrequently as possible
 - b. Irrigate the wound**
 - c. Debridement
 - d. Gentle cleansing.
47. While managing Mrs. Ogbu's wound, Rose decided to use a scalpel to separate or remove dead tissue. Which type of debridement involves the use of scalpel or scissors?
- a. Sharp debridement**
 - b. Chemical debridement
 - c. Mechanical debridement
 - d. Autolytic debridement.
48. When the method of wound healing is one in which wound edges are not surgically approximated and integumentary continuity is restored by granulations, the wound healing is termed
- a. Secondary intention healing**

- b. Primary intention healing
 - c. Third intention healing
 - d. First intention healing
49. A nursing care plan for a client with suprapubic cystostomy would include
- a. Placing a urinal bag around the tube insertion to collect the urine
 - b. Clamping the tube and allowing the client to void through the urinary meatus before removing the tube**
 - c. Catheter irrigations every 4 hours to prevent formation of uri-nary stones
 - d. Limiting fluid intake to 1500ml per day.
50. The physician has just completed a liver biopsy. Immediately fol-lowing the procedure, the nurse will position the client
- a. On his right side to promote haemostasis**
 - b. In fowler's position to facilitate ventilation
 - c. Supine to maintain blood pressure
 - d. In Sim's position to prevent aspiration
51. A female client has orders for a cholecystogram. Prior to the test, the nursing intervention would be to
- a. Provide fat diet for dinner, then NPO
 - b. Explain that diarrhoea may result from the dye tablets**
 - c. Administer the dye tablets following a regular diet for dinner
 - d. Administer enemas until clear.
52. Myasthenic crisis and cholinergic crisis are the major complica-tions of myasthenia gravis. Which of the following is essential nursing knowledge when caring for a client in crisis?
- a. Weakness and paralysis of the muscles for swallowing and breathing occur in either crisis**
 - b. Cholinergic drugs should be administered to prevent further complications associated with the crisis
 - c. The clinical condition of the client usually improves after sev-eral days of treatment
 - d. Loss of body function creates high levels of anxiety and fear
53. Nurse Ada is preparing to ambulate a female client. The best and the safest position for the nurse in assisting the client is to stand:
- a. On the unaffected side of the client
 - b. On the affected side of the client**
 - c. In front of the client
 - d. Behind the client

54. Nurse Eke hears the alarm sound on the telemetry monitor. The nurse quickly looks at the monitor and notes that a client is in a ventricular tachycardia. The nurse rushes to the client's room upon reaching the client's bedside, the nurse would take which action first?
- a. Prepare for cardioversion
 - b. Prepare to defibrillate the client
 - c. Call a code
 - d. Check the client's level of consciousness**
55. Moon face, truncal obesity, "buffalo hump" and proximal muscle weakness are important clues to diagnosis of
- a. Kwashiorkor disease
 - b. Cushing syndrome**
 - c. Addison disease
 - d. Graves disease
56. Intracranial pressure is said to have increased beyond normal range when it is above
- a. 30mmHg
 - b. 25mmHg
 - c. 20mmHg
 - d. 15mmHg**
57. The Physician inserts a chest tube into a female client to treat a pneumothorax. The tube is connected to underwater sealed drainage. The nurse in charge can prevent chest tube air leaks by
- a. Checking and taping all connections**
 - b. Checking patency of the chest tube
 - c. Keeping the head of the bed slightly elevated
 - d. Keeping the chest drainage system below the level of the chest
58. A male client receiving total parenteral nutrition suddenly demonstrates signs and symptoms of an air embolism. What is the priority action by the nurse?
- a. Notify the physician
 - b. Place the client on the left side in the Trendelenburg position**
 - c. Place the client in high Fowler's position
 - d. Stop the total parenteral nutrition
59. A male client is being transferred to the nursing unit for admission after receiving a radium implant for bladder cancer. The nurse in charge would take which priority action in the care of this client
- a. Place client on reverse isolation
 - b. Admit the client into a private room**
 - c. Encourage the client to take frequent rest periods