

SELECT THE MOST APPROPRIATE OPTION

11. When a nurse is able to perform comprehensive and systematic patient assessment to deliver effective nursing care to the patient, the nurse is said to be competent in...

- a. Management of care.
- b. Professionalism
- c. Legal issue
- d. Nursing ethics

SELECT THE MOST APPROPRIATE OPTION

12. A client is undergoing a comprehensive admission assessment, the nurse knows that, the single most significant indicator of cerebral perfusion and functioning is called

- a. Level of consciousness
- b. Glasgow coma scale
- c. Intra cranial pressure
- d. Pupils dilatation

SELECT THE MOST APPROPRIATE OPTION

13. It is a vigorous rubbing together of lathered hands for 15secs followed by a thorough rinsing under running stream of water:

- a. Hand sanitization
- b. Hand rubbing
- c. Hand washing
- d. Hand sanity

SELECT THE MOST APPROPRIATE OPTION

14. The nurse delivers a D.C. counter shock to a cardiac patient in a critical incidence, the nurse knows that the **MOST** appropriate equipment to use in this situation is:

- a. Arterial line
- b. Defibrillator
- c. Glucometer
- d. Spiro metery

SELECT THE MOST APPROPRIATE OPTION

15. A patient has a diagnosis of ineffective airway clearance. He has sustained an inhalation burn injury. Amongst the following nursing interventions which of the following is the most rational?

- a. Monitor respiratory rate and breathing 4 hours
- b. Elevate the head of the bed
- c. Encourage cough exercise
- d. Monitor oxygen levels every 4 hours

SELECT THE MOST APPROPRIATE OPTION

16. A client who has an inhalation injury comes to emergency unit. On the initial assessment, the nurse notes that the client is very confused and physically defensive. What do you think the client may be experiencing from?

- a. Fear
- b. Anxiety
- c. Hypoxia
- d. Pain

SELECT THE MOST APPROPRIATE OPTION

- 17.** Which nursing intervention can help prevent sharp spikes in ICP and surgical haemorrhage?
- a. Give fluid restriction
 - b. Help with breath exercises
 - c. Keep the head of the bed elevated
 - d. Administer an antiemetic to prevent vomiting

SELECT THE MOST APPROPRIATE OPTION

18. A patient presents to the surgical ward after an exploratory laparotomy related to trauma. The patient is hypotensive and tachycardic. This is most likely due to which of the following ?

- a. Hypovolaemia
- b. Pain
- c. Hypothermia
- d. hyperthermia

SELECT THE MOST APPROPRIATE OPTION

19. A 16 year-old is admitted to the floor after a foot ball injury. The patient has a femur fracture. An external fixation device was applied in the operating room. The patient is at an increased risk for development of which of the following complications:

- a. Post traumatic stress disorder.
- b. Deep venous thrombosis.
- c. Pulmonary embolism.
- d. Pressure sore.

SELECT THE MOST APPROPRIATE OPTION

20. A 40 year-old woman is involved in a road traffic accident. She presents with chest pain, shortness of breath, and difficulty breathing. Her arterial oxygen saturation is 85% and her respiratory rate is 40b/minute. A chest tube is placed and the nurse observes a return of 300 mls blood into the system. What is the nurse's next line of action ?

- a. Clamp the tube and notify surgeon immediately.
- b. Call for a chest x-ray.
- c. Place the patient on oxygen.
- d. Explain to the patient what is going on to allay anxiety.