

NMTC - ATRHE
MID - SEMESTER EXAMINATION, NOVEMBER 2024
FIRST AID, EMERGENCY AND DISASTER MANAGEMENT
DURATION: 30 MINUTES

SECTION A

INSTRUCTION: CIRCLE THE LETTER OF THE MOST APPROPRIATE ANSWER ON THE QUESTION PAPER

1. What is the primary aim of first aid?
 - a) To inform bystanders
 - b) To provide transportation
 - c) To reduce pain
 - ☒ d) To save life
2. Which of the following actions should NOT be taken for an unconscious, vomiting casualty?
 - a) Ensuring they are lying on their side
 - ☒ b) Giving them water to prevent dehydration
 - c) Placing them in the recovery position
 - d) Turning their head to one side
3. First aid aims to prepare for further medical treatment by:
 - a) Allowing bystanders to apply treatments
 - ☒ b) Arranging suitable transportation to the hospital
 - c) Avoiding the use of medical terms
 - d) Completing the full treatment
4. In an emergency where a victim is unconscious, which doctrine allows a first aider to proceed with treatment?
 - a) Consent Doctrine
 - ☒ b) Emergency Doctrine
 - c) Mandatory Reporting
 - d) Privacy and Confidentiality Doctrine
5. Which of the following statements is TRUE regarding informed consent in first aid?
 - a) Informed consent can be obtained even if the victim does not understand the language used.
 - b) Informed consent is only necessary when the victim is a minor.
 - c) Informed consent is optional for adults and required only for children.
 - ☒ d) Informed consent is valid as long as the victim is an adult and of sound mind.
6. Under the principles of confidentiality, a first aider must:
 - a) Disclose the victim's condition to anyone who inquires.

- b) Maintain confidentiality and not disclose information without the victim's permission.
- c) Only disclose information to other healthcare providers without restrictions.
- d) Share the victim's information only with close family members.

7. A first aider suspects that an injury is related to a violent crime. How should they handle any clothing removed from the victim?

- a) Discard the clothing after treatment.
- b) Give it directly to the victim's family.
- c) Place it in a paper bag to preserve the evidence. ✓
- d) Place it in a plastic bag to prevent damage.

8. Which of the following situations would likely require the first aider to follow mandatory reporting laws?

- a) A minor burn with no criminal implications
- b) A minor injury from a sports accident
- c) A non-emergency medical condition
- d) A road traffic accident with serious injuries

9. Which medication is primarily included in first aid kits for treating central chest pain due to its anti-platelet properties?

- a) Aspirin ✓
- b) Diphenhydramine
- c) Ibuprofen
- d) Paracetamol

10. Which of the following medications would be used in a first aid kit to treat anaphylactic shock in emergency situations?

- a) Activated charcoal
- b) Anti-inflammatory painkiller
- c) Epinephrine auto-injector ✓
- d) Loperamide

11. Which medication in the first aid kit serves as both a painkiller and an anti-diarrheal?

- a) Codeine ✓
- b) Hydrocortisone cream
- c) Ibuprofen
- d) Paracetamol

12. Which of the following would likely be the initial response of the body in hypovolemic shock due to severe dehydration?

- a) Decreased breathing rate to stabilize blood oxygen levels
- b) Increased blood volume by retaining water ✓

- ✓ c) Increased heart rate to compensate for low blood volume
- d) Slower heart rate to conserve energy

13. A patient experiences a sudden rise in blood pressure shortly after an injury, causing bleeding to resume after it had previously stopped. What type of haemorrhage is this?

- a) Capillary haemorrhage
- b) Primary haemorrhage
- ✓ c) Relapsing haemorrhage
- d) Secondary haemorrhage

14. During an assessment, blood is observed oozing evenly from all parts of a wound and has a regular red colour. Which type of haemorrhage is most likely being observed?

- a) Arterial haemorrhage
- ✓ b) Capillary haemorrhage
- c) Primary haemorrhage
- d) Venous haemorrhage

15. A patient arrives with a nail embedded in their leg, causing bleeding. What should you do first?

- a) Apply a tight tourniquet around the object to stop bleeding
- ✓ b) Apply pressure around the nail and secure it with a donut bandage
- c) Pull out the nail and apply direct pressure to the wound
- d) Remove any dressings and allow the wound to bleed for 1-2 minutes

16. You are providing first aid to someone with an embedded object in the chest. What is a critical next step to minimize movement of the object?

- a) Apply pressure directly on the object
- b) Bandage the wound loosely around the object
- ✓ c) Loosen all tight clothing to ease breathing
- d) Remove any clothing around the wound

17. A patient with a deep laceration is experiencing extreme thirst and pallor. What is the most likely reason for these symptoms?

- a) Blood loss causing shock
- ✓ b) Dehydration from fluid loss
- c) Excessive caffeine intake
- d) High blood pressure due to anxiety

18. What is the color and nature of the blood seen in haemoptysis?

- ✓ a) Bright red and frothy
- b) Dark brown and gritty
- c) Dark red and thick
- d) Pale yellow and watery

19. When treating a patient with a tourniquet, how often should the tourniquet be released?

- a) Every 10 minutes
- b) Every 20 minutes
- c) Every 30 minutes ✓
- d) Never

20. During first aid treatment for hemorrhage, when is it appropriate to give the victim something to drink?

- a) After bleeding has been stopped completely
- b) Never, as it may cause choking ✓
- c) Only after medical personnel arrive
- d) Only if the victim complains of thirst

21. Which of the following is a typical sign of a fracture in the lower limb?

- a) Pain that decreases with movement
- b) Shortening of the limb compared to the opposite side ✓
- c) Swelling without bruising
- d) Excessive blood loss at the site of injury

22. What is the first priority when treating a suspected fracture?

- a) Apply an ice pack to the injury
- b) Immobilize the fractured area ✓
- c) Check for a pulse
- d) Apply a bandage around the site of injury

23. Which of the following is a common symptom of a fracture caused by excessive muscular contraction?

- a) Loss of consciousness
- b) Sudden pain without any visible injury ✓
- c) Pain and a sudden snapping sound
- d) Numbness in the affected area

24. What should a first aider do when they detect unnatural movement at the site of a suspected fracture?

- a) Attempt to realign the bone to its normal position
- b) Test for crepitus to confirm the fracture
- c) Immobilize the affected limb and avoid moving it ✓
- d) Apply heat to the affected area to reduce swelling

25. Which of the following factors could predispose someone to a fracture?

- a) Increased physical activity
- b) Being underweight
- c) Extremes of age (very young or very old) ✓
- d) Healthy bones with no history of injury

26. A casualty has suffered a compound fracture of the leg with heavy bleeding. What is the immediate first aid priority?

- a) Immobilize the leg to prevent further movement and bleeding ✓
- b) Provide pain relief and wait for medical help
- c) Attempt to realign the bones to their original position

d) Cover the wound with a bandage and wait for the swelling to subside

27. Which of the following complications can occur from a badly applied splint?

- a) Paralysis due to nerve compression
- ☒ b) Sepsis due to infection from an open fracture
- c) Pressure sore due to improper splinting technique ✓
- d) Peritonitis in cases of pelvic fractures

28. In the case of a fracture with visible blood loss, what is the first aid aim?

- a) To reduce pain and make the casualty comfortable
- ☒ b) To stop the bleeding and prevent shock
- c) To transport the casualty as quickly as possible
- d) To apply a bandage to the injured part only

29. What is the best method of immobilizing a fractured arm when no mechanical splints are available?

- ☒ a) Apply a body splint by bandaging the arm to the chest
- b) Use a piece of cloth to tie the arm to the torso
- c) Place the arm in an elevated position and apply pressure to reduce swelling
- d) Allow the casualty to move the arm to reduce stiffness

30. A patient with a pelvic fracture is at risk of developing which of the following complications?

- a) Gangrene due to artery injury
- b) Peritonitis due to injury to abdominal organs
- ☒ c) Sepsis from a compound fracture wound
- ☒ d) Paralysis due to compression of the spine

31. Which action should be taken to prevent infection in a compound fracture?

- ☒ a) Apply a clean dressing to cover the wound and stop any bleeding
- b) Expose the fracture to air for faster healing
- c) Apply a hot compress to the wound to promote circulation
- d) Try to remove any foreign object from the wound

32. A person has just twisted their ankle during a game, and they are experiencing severe pain and swelling. What is the first step in treating this injury?

- a) Apply a heat pad to the ankle to relax the muscles
- ☒ b) Apply a cold compress to the ankle to prevent swelling
- c) Bandage the ankle tightly to stop the swelling
- d) Encourage the person to walk on the ankle to check for mobility

33. A casualty has sustained a sprained wrist. After applying a cold compress and resting the joint, you should next:
- a) ☒ Expose the joint and wrap it with a cotton band, applying firm pressure with a bandage
 - b) Attempt to move the wrist gently to check the range of motion
 - c) Apply a warm compress to ease the muscle tension
 - d) Encourage the casualty to use the wrist lightly to strengthen it
34. A person is found unconscious on the floor. You check for signs of life, and they are not responding to stimuli. What is the first step you should take?
- a) Try to wake them up by shaking them
 - b) ☒ Immediately call for medical help or ambulance
 - c) Place the person in a standing position to help them regain consciousness
 - d) Give them water to drink
35. A casualty is unconscious and lying on their back. You are unable to find any external injuries but notice shallow breathing. What should you do next?
- a) Leave them in the current position, as they are breathing
 - b) ☒ Put them in the recovery position to ensure their airway remains clear
 - c) Try to wake them by pinching their ear or arm
 - d) Give the patient a few sips of water to hydrate them
36. You find a person unconscious, breathing slowly, and with a sweet smell to their breath. What does this likely indicate and what should you do?
- a) The person may have a head injury; keep the person warm and wait for help
 - b) The person may be diabetic and in a hypoglycemic state; give them food or drink immediately
 - c) ☒ The person may have alcohol poisoning; ensure they are in the recovery position and call for medical help
 - d) The person is dehydrated; try to give them water
37. A patient has been unconscious for several minutes after a fall. You have called for help, and the ambulance is on the way. What is the most important action to take while waiting for medical assistance?
- a) Keep the patient sitting upright to help them breathe
 - b) ☒ Keep the patient still and place them in the recovery position to maintain airway and prevent choking
 - c) Give the patient food or drink to help them regain consciousness
 - d) Attempt to revive the person by shaking them or tapping their face
38. You find an unconscious person lying on the ground with cyanosis (blue coloring) around their lips. What should you do first?

- ☒ a) Check their pulse and breathing
- b) Immediately give them water to drink
- ☒ c) Check for wounds on the head
- d) open windows and put him in the recovery position

39. While examining an unconscious patient, you notice a deep wound on the skull. The person is breathing but is not responding to stimuli. What is the best course of action?

- ☒ a) Apply direct pressure to the wound and wait for medical help
- b) Keep the person lying on their back with their head tilted forward
- ☒ c) Keep in the recovery position and call for immediate medical assistance
- d) Try to wake the patient by slapping their face

40. A person suddenly starts having a seizure, with their body becoming rigid and then jerking uncontrollably. The first thing you should do is:

- a) Try to hold them still to prevent injury
- ☒ b) Make sure the area around the person is clear of dangerous objects
- c) Give them something to drink to help them recover
- d) Force something into their mouth to prevent them from biting their tongue